

# Psy 613

## Assignment 2

**1**

For the matrices below, obtain the following matrices, and state their dimensions: (1)  $\mathbf{A} + \mathbf{B}$ , (2)  $\mathbf{A} - \mathbf{B}$ , (3)  $\mathbf{AC}$ , (4)  $\mathbf{AB}'$ , (5)  $\mathbf{B}'\mathbf{A}$

$$\mathbf{A} = \begin{bmatrix} 1 & 4 \\ 2 & 6 \\ 3 & 8 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 1 & 3 \\ 1 & 4 \\ 2 & 5 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 3 & 8 & 1 \\ 5 & 4 & 0 \end{bmatrix}$$

**2**

Let  $\mathbf{B}$  be defined as follows:

$$\mathbf{B} = \begin{bmatrix} 1 & 5 & 0 \\ 1 & 0 & 5 \\ 1 & 0 & 5 \end{bmatrix}$$

- a Are the column vectors of  $\mathbf{B}$  linearly dependent?
- b What is the rank of  $\mathbf{B}$ ?
- c What must be the determinant of  $\mathbf{B}$ ?

**3**

- a Find the inverse of  $\mathbf{A}$ :

$$\mathbf{A} = \begin{bmatrix} 2 & 4 \\ 3 & 1 \end{bmatrix}$$

- b Check that your resulting matrix is indeed the inverse.

## 4

Diagonal matrices have non-zero elements on the main diagonal, but zeroes in the off-diagonals. Show that the following is true:

- a If all diagonal elements  $d_{ij}$  of  $\mathbf{D}$  are the *same*, then pre- or postmultiplying any matrix  $\mathbf{X}$  with  $\mathbf{D}$  gives the same result, namely, a scalar multiplication of  $\mathbf{X}$  with  $d$ .
- b If the diagonal elements  $d_{ij}$  of a diagonal matrix  $\mathbf{D}$  are *different* from each other, then premultiplying a matrix  $\mathbf{X}$  with  $\mathbf{D}$  gives a different result than post-multiplying  $\mathbf{X}$  with with  $\mathbf{D}$ .
- c Try to explain why both (a) and (b) have to be true.

## 5

Compute the results of the following operations by hand, if they have a solution.

$$\mathbf{a} = \begin{bmatrix} 1 \\ 8 \\ 4 \\ 2 \end{bmatrix} \quad \mathbf{y} = \begin{bmatrix} 7 \\ 5 \\ 10 \\ 2 \end{bmatrix} \quad \mathbf{X} = \begin{bmatrix} 7 & 3 \\ 5 & 3 \\ 10 & 3 \\ 2 & 1 \end{bmatrix} \quad \underline{\mathbf{1}} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

$$\frac{1}{4}\mathbf{a} =$$

$$\mathbf{a}' =$$

$$\mathbf{a}'\mathbf{y} =$$

$$\underline{\mathbf{1}}'\mathbf{a} =$$

$$\mathbf{a}'\mathbf{a} =$$

$$\mathbf{a} - \frac{1}{4}(\mathbf{a}'\underline{\mathbf{1}}) =$$

$$\mathbf{a}'\mathbf{X} =$$

$$\mathbf{X}'\mathbf{a} =$$

$$\mathbf{X}'\mathbf{X} =$$

$$\mathbf{X}\mathbf{X}^{-1} =$$

## 6

Using matrix algebra, derive a correlation matrix for the following data:

Cases	X1	X2
1	8	4
2	7	3
3	2	6
4	2	7
5	6	0

Show your work at each of the following steps:

- a** Pre-multiply your  $n \times p$  data matrix by the transpose of an appropriate unit vector ( $\mathbf{1}'$ ); then multiply by the scalar  $n^{-1}$  to compute a vector of means. Thus you compute  $\mathbf{1}'\mathbf{X}n^{-1}$ .
- b** Premultiply the above  $1 \times p$  vector of means by an appropriate unit vector such that you expand the means vector to  $n \times p$  dimensionality. Then subtract this expanded matrix of means from the raw data matrix to produce mean-deviated (centered) scores:  $\mathbf{X} - \mathbf{1}\mathbf{1}'\mathbf{X}n^{-1}$ , often written as  $\mathbf{Y}$ .
- c** Generate a  $p \times p$  Sums-of-Squares and Cross-Products matrix from your  $\mathbf{Y}$  matrix. Then multiply this SSQ/CP matrix by the scalar  $(n-1)^{-1}$  to compute  $\mathbf{S}_{\mathbf{y}\mathbf{y}}$ , the variance-covariance matrix of the mean-deviated scores, which is equivalent to  $\mathbf{S}_{\mathbf{x}\mathbf{x}}$ .
- d** Pre- and post-multiply  $\mathbf{S}_{\mathbf{y}\mathbf{y}}$  with an appropriate diagonal matrix that contains the reciprocals of the corresponding standard deviations,  $s_{x1}$  and  $s_{x2}$ . Briefly explain what this step accomplishes.