

Summary on Lecture 1, April 2, 2018

**Recurrence Relations**

**Warm-up: linear recurrence relations.**

(1) **Geometric progression.** Define a sequence  $\{a_n\}$  as follows:  $a_0 = A$ ,  $a_{n+1} = da_n$ ,  $n \geq 1$ . Then we have:

$$a_1 = dA, \quad a_2 = d^2A, \quad a_3 = d^3A, \quad \dots \quad a_n = d^nA, \quad \dots$$

Thus we have a general formula:  $a_n = d^nA$ . This is a *geometric progression*.

**Exercise.** Prove formula  $a_n = d^nA$  by induction.

**Definition.** A recurrence relation  $a_{n+1} - da_n = 0$ , where  $d$  is a constant, is called *linear relation*. More general, a recurrence relation  $a_{n+1} - da_n = f(n)$ , where  $d$  is a constant, and  $f(n)$  is a function, is called a *first order relation*.

(2) **Example: Bubble Sort algorithm.** Let  $x_1, \dots, x_n$  be  $n$  real numbers. We would like to sort them out into ascending order. Here is an algorithm known as **BubbleSort**:

```

begin(BubbleSort)
  for i := 1 to n - 1 do
    for j := n down to i + 1 do
      if  $x_j < x_{j-1}$  then
        begin(Interchange)
           $t := x_{j-1}$ 
           $x_{j-1} := x_j$ 
           $x_j := t$ 
        end(Interchange)
      end(BubbleSort)

```

First, we would like to understand how does it work. Let us start with the sequence  $(x_1, x_2, x_3, x_4, x_5) = (7, 9, 2, 5, 8)$ .

$i = 1$		$j = 5$	$j = 4$	$j = 3$	$j = 2$	
$x_1$		7	7	7	2	2
$x_2$		9	9	2	7	7
$x_3$	:=	2	2	9	9	9
$x_4$		5	5	5	5	5
$x_5$		8	8	8	8	8

$i = 2$		$j = 5$	$j = 4$	$j = 3$	
$x_1$		2	2	2	2
$x_2$		7	7	5	5
$x_3$	:=	9	5	7	7
$x_4$		5	9	9	9
$x_5$		8	8	8	8

$i = 3$		$j = 5$	$j = 4$	
$x_1$		2	2	2
$x_2$		5	5	5
$x_3$	:=	7	7	7
$x_4$		8	8	8
$x_5$		9	9	9

$i = 4$		$j = 5$	
$x_1$		2	2
$x_2$		5	5
$x_3$	:=	7	7
$x_4$		8	8
$x_5$		9	9

Here we have: for  $i = 1$ , 4 comparisons and 2 interchanges, for  $i = 2$ , 3 comparisons and 2 interchanges, for  $i = 3$ , 2 comparisons and 1 interchange, for  $i = 4$ , 1 comparison and no interchanges.

Now we denote by  $a_n$  a total number of comparisons to sort out a sequence  $(x_1, \dots, x_n)$ . First, we can identify the smallest number: this is done when we run the algorithm for  $i = 1$ . Clearly, we use  $(n - 1)$  comparisons for that. Then we obtain the recursion:

$$a_1 = 0, \quad a_n = a_{n-1} + (n - 1).$$

We have:

$$\begin{aligned}
 a_1 &= 0 \\
 a_2 &= a_1 + (2 - 1) = 1 \\
 a_3 &= a_2 + (3 - 1) = 1 + 2 \\
 a_4 &= a_3 + (4 - 1) = 1 + 2 + 3 \\
 \dots &\quad \dots \quad \dots \\
 a_n &= a_{n-1} + (n - 1) = 1 + 2 + 3 + \dots + (n - 1)
 \end{aligned}$$

The answer:

$$a_n = 1 + 2 + 3 + \dots + (n - 1) = \frac{(n - 1)n}{2} = \frac{1}{2}(n^2 - n).$$

In that case we say that the time-complexity function of that algorithm is  $O(n^2)$ .

**Second Order Recurrence Relations.** Let  $\{a_n\}$  be a Fibonacci sequence, i.e.  $a_0 = 0$ ,  $a_1 = 1$ , and  $a_n = a_{n-1} + a_{n-2}$  for all  $n \geq 2$ . We would like to find a *closed formula* for  $a_n$ 's. Let us try  $a_n = c \cdot r^n$ , where  $c \neq 0$  and  $r$  some real numbers. Then the relation  $a_n = a_{n-1} + a_{n-2}$  gives:

$$cr^n = cr^{n-1} + cr^{n-2}, \quad n \geq 2.$$

We cancel  $cr^{n-2}$  and get the equation  $r^2 = r + 1$  or  $r^2 - r - 1 = 0$ . We find the solutions:

$$r = \frac{1 \pm \sqrt{5}}{2}, \quad \text{or} \quad r_1 = \frac{1 + \sqrt{5}}{2}, \quad r_2 = \frac{1 - \sqrt{5}}{2}.$$

Then both sequences  $c_1 r_1^n$  and  $c_2 r_2^n$  will satisfy the relation  $a_n = a_{n-1} + a_{n-2}$ . Moreover, the sequence  $c_1 r_1^n + c_2 r_2^n$  will satisfy the same relation. Then we can find  $c_1$  and  $c_2$ .

We have for  $n = 0$  and  $n = 1$ :

$$\begin{cases} 0 = c_1 + c_2 \\ 1 = c_1 r_1 + c_2 r_2 \end{cases} \quad \begin{cases} c_2 = -c_1 \\ 1 = c_1 r_1 - c_1 r_2 \end{cases} \quad \begin{cases} c_2 = -\frac{1}{\frac{r_1 - r_2}{r_1 - r_2}} \\ c_1 = \frac{1}{r_1 - r_2} \end{cases}$$

Since  $r_1 - r_2 = \sqrt{5}$ , we obtain a formula for  $a_n$ :

$$a_n = c_1 r_1^n + c_2 r_2^n = \frac{1}{\sqrt{5}} \left[ \left( \frac{1 + \sqrt{5}}{2} \right)^n - \left( \frac{1 - \sqrt{5}}{2} \right)^n \right].$$

**Theorem 1.** Let  $a_0$  and  $a_1$  are given, and  $a_n = Aa_{n-1} + Ba_{n-2}$  be a second order recurrence relation,  $n \geq 2$ , where  $A, B$  are non-zero constants. Assume that the characteristic equation  $r^2 - Ar - B = 0$  has two real different real solutions  $r_1$  and  $r_2$ . Then  $a_n = c_1 r_1^n + c_2 r_2^n$ , where the constants  $c_1$  and  $c_2$  are determined by solving the system

$$\begin{cases} a_0 = c_1 + c_2 \\ a_1 = c_1 r_1 + c_2 r_2 \end{cases}$$

**Proof.** Indeed, we look for a solution  $a_n = cr^n$ , then the recurrence relation  $a_n = Aa_{n-1} + Ba_{n-2}$  gives the characteristic equation  $r^2 - Ar - B = 0$ . By assumption, there are two two different real solutions,  $r_1$  and  $r_2$  of  $r^2 - Ar - B = 0$ . Then the sum  $c_1 r_1^n + c_2 r_2^n$  will satisfy the recurrence. Finally, we notice that the system

$$\begin{cases} a_0 = c_1 + c_2 \\ a_1 = c_1 r_1 + c_2 r_2 \end{cases} \quad \text{always have a unique solution if } r_1 \neq r_2 \text{ (Explain why).}$$