

Summary on Lecture 1, January 5, 2015

Recurrence Relations

Warm-up: linear recurrence relations.

(1) **Geometric progression.** Define a sequence $\{a_n\}$ as follows: $a_0 = A$, $a_{n+1} = da_n$, $n \geq 1$. Then we have:

$$a_1 = dA, \quad a_2 = d^2A, \quad a_3 = d^3A, \quad \dots \quad a_n = d^nA, \quad \dots$$

Thus we have a general formula: $a_n = d^nA$. This is a *geometric progression*.

Exercise. Prove formula $a_n = d^nA$ by induction.

Definition. A recurrence relation $a_{n+1} - da_n = 0$, where d is a constant, is called *linear relation*. More general, a recurrence relation $a_{n+1} - da_n = f(n)$, where c is a constant, and $f(n)$ is a function, is called a *first order relation*.

(2) **Example: Bubble Sort algorithm.** Let x_1, \dots, x_n be n real numbers. We would like to sort them out into ascending order. Here is an algorithm known as **BubbleSort**:

```

begin(BubbleSort)
  for i := 1 to n - 1 do
    for j := n down to i + 1 do
      if  $x_j < x_{j-1}$  then
        begin(Interchange)
           $t := x_{j-1}$ 
           $x_{j-1} := x_j$ 
           $x_j := t$ 
        end(Interchange)
    end(BubbleSort)

```

First, we would like to understand how does it work. Let us start with the sequence $(x_1, x_2, x_3, x_4, x_5) = (7, 9, 2, 5, 8)$.

$i = 1$		$j = 5$	$j = 4$	$j = 3$	$j = 2$	
x_1		7	7	7	2	2
x_2		9	9	2	7	7
x_3	:=	2	2	9	9	9
x_4		5	5	5	5	5
x_5		8	8	8	8	8

$i = 2$		$j = 5$	$j = 4$	$j = 3$	
x_1		2	2	2	2
x_2		7	7	5	5
x_3	:=	9	5	7	7
x_4		5	9	9	9
x_5		8	8	8	8

$i = 3$		$j = 5$	$j = 4$	
x_1		2	2	2
x_2		5	5	5
x_3	:=	7	7	7
x_4		8	8	8
x_5		9	9	9

$i = 4$		$j = 5$	
x_1		2	2
x_2		5	5
x_3	:=	7	7
x_4		8	8
x_5		9	9

Here we have: for $i = 1$, 4 comparisons and 2 interchanges, for $i = 2$, 3 comparisons and 2 interchanges, for $i = 3$, 2 comparisons and 1 interchange, for $i = 4$, 1 comparison and no interchanges.

Now we denote by a_n a total number of comparisons to sort out a sequence (x_1, \dots, x_n) . First, we can identify the smallest number: this is done when we run the algorithm for $i = 1$. Clearly, we use $(n - 1)$ comparisons for that. Then we obtain the recursion:

$$a_1 = 0, \quad a_n = a_{n-1} + (n - 1).$$

We have:

$$\begin{aligned}
 a_1 &= 0 \\
 a_2 &= a_1 + (2 - 1) = 1 \\
 a_3 &= a_2 + (3 - 1) = 1 + 2 \\
 a_4 &= a_3 + (4 - 1) = 1 + 2 + 3 \\
 \dots &\quad \dots \quad \dots \\
 a_n &= a_{n-1} + (n - 1) = 1 + 2 + 3 + \dots + (n - 1)
 \end{aligned}$$

The answer:

$$a_n = 1 + 2 + 3 + \dots + (n - 1) = \frac{(n - 1)n}{2} = \frac{1}{2}(n^2 - n).$$

In that case we say that the time-complexity function of that algorithm is $O(n^2)$.

Second Order Recurrence Relations. Let $\{a_n\}$ be a Fibonacci sequence, i.e. $a_0 = 0$, $a_1 = 1$, and $a_n = a_{n-1} + a_{n-2}$ for all $n \geq 2$. We would like to find a *closed formula* for a_n 's. Let us try $a_n = c \cdot r^n$, where $c \neq 0$ and r some real numbers. Then the relation $a_n = a_{n-1} + a_{n-2}$ gives:

$$cr^n = cr^{n-1} + cr^{n-2}, \quad n \geq 2.$$

We cancel cr^{n-2} and get the equation $r^2 = r + 1$ or $r^2 - r - 1 = 0$. We find the solutions:

$$r = \frac{1 \pm \sqrt{5}}{2}, \quad \text{or} \quad r_1 = \frac{1 + \sqrt{5}}{2}, \quad r_2 = \frac{1 - \sqrt{5}}{2}.$$

Then both sequences $c_1 r_1^n$ and $c_2 r_2^n$ will satisfy the relation $a_n = a_{n-1} + a_{n-2}$. Moreover, the sequence $c_1 r_1^n + c_2 r_2^n$ will satisfy the same relation. Then we can find c_1 and c_2 .

We have for $n = 0$ and $n = 1$:

$$\begin{cases} 0 = c_1 + c_2 \\ 1 = c_1 r_1 + c_2 r_2 \end{cases} \quad \begin{cases} c_2 = -c_1 \\ 1 = c_1 r_1 - c_1 r_2 \end{cases} \quad \begin{cases} c_2 = -\frac{1}{\frac{r_1 - r_2}{1}} \\ c_1 = \frac{1}{r_1 - r_2} \end{cases}$$

Since $r_1 - r_2 = \sqrt{5}$, we obtain a formula for a_n :

$$a_n = c_1 r_1^n + c_2 r_2^n = \frac{1}{\sqrt{5}} \left[\left(\frac{1 + \sqrt{5}}{2} \right)^n - \left(\frac{1 - \sqrt{5}}{2} \right)^n \right].$$

Let $a_n = Aa_{n-1} + Ba_{n-2}$ be a second order recurrence relation. Then the equation $r^2 - Ar - B = 0$ is called a *characteristic equation* of that relation.

Theorem 1. Let a_0 and a_1 are given, and $a_n = Aa_{n-1} + Ba_{n-2}$ be a recurrence relation, $n \geq 2$, where A, B are non-zero constants. Assume that the characteristic equation $r^2 - Ar - B = 0$ has two real different real solutions r_1 and r_2 . Then $a_n = c_1 r_1^n + c_2 r_2^n$, where the constants c_1 and c_2 are determined by solving the

$$\text{system} \begin{cases} a_0 = c_1 + c_2 \\ a_1 = c_1 r_1 + c_2 r_2 \end{cases}$$

Proof. Indeed, we look for a solution $a_n = cr^n$, then the recurrence relation $a_n = Aa_{n-1} + Ba_{n-2}$ gives the characteristic equation $r^2 - Ar - B = 0$. By assumption, there are two two different real solutions, r_1 and r_2 of $r^2 - Ar - B = 0$. Then the sum $c_1 r_1^n + c_2 r_2^n$ will satisfy the recurrence. Finally, we notice that the system

$$\begin{cases} a_0 = c_1 + c_2 \\ a_1 = c_1 r_1 + c_2 r_2 \end{cases} \quad \text{always have a unique solution if } r_1 \neq r_2 \text{ (Explain why).}$$

Concluding question: How to solve this problem if $r_1 = r_2$?