

Summary on Lecture 8, January 27, 2017

**Introduction to Graph Theory: more.**

Let  $n$  be a positive integer. Then a *complete graph*  $K_n$  is a graph with  $n$  vertices  $v_1, \dots, v_n$  and  $\binom{n}{2}$  edges  $e_{i,j} = \{v_i, v_j\}$ , where  $i \neq j$ .

A complete graph  $K_n$ , contains subgraphs isomorphic to the graphs  $K_m$  for  $m = 1, 2, \dots, n$ . Such a subgraph can be obtained by selecting any  $m$  of the  $n$  vertices and using all the edges in  $K_n$  joining them. Thus  $K_5$  contains  $\binom{5}{2} = 10$  subgraphs isomorphic to  $K_2$ ,  $\binom{5}{3} = 10$  subgraphs isomorphic to  $K_3$  [i.e., triangles], and  $\binom{5}{4} = 5$  subgraphs isomorphic to  $K_4$ . In fact, every graph with  $n$  or fewer vertices and with no loops or parallel edges is isomorphic to a subgraph of  $K_n$ ; just delete the unneeded edges from  $K_n$ .




Fig. 6. Complete graphs

Complete graphs have a high degree of symmetry. Each permutation  $\alpha$  of the vertices of a complete graph gives an isomorphism of the graph onto itself, since both  $\{u, v\}$  and  $\{\alpha(u), \alpha(v)\}$  are edges whenever  $u \neq v$ . The next theorem relates the degrees of vertices to the number of edges of the graph.

**Theorem 2.** *The sum of the degrees of the vertices of a graph  $G = (V, E)$  is twice the number of edges, i.e.,*

$$\sum_{v \in V} \deg(v) = 2 \cdot |E(G)|.$$

**Proof.** Each edge, whether a loop or not, contributes 2 to the degree sum. This is a place where our convention that each loop contributes 2 to the degree of a vertex pays off.  $\square$ .

**Euler Trails and circuits**

**The Seven Bridges of Königsberg.** *The Seven Bridges of Königsberg Problem* is a historically important problem in mathematics. Its negative resolution by Leonhard Euler in 1736 laid the foundations of *graph theory* and prefigured the idea of *topology*.




Fig. 7. The Seven Bridges of Königsberg<sup>1</sup>

Here is The Seven Bridges of Königsberg Problem: find a walk through the city that would cross each bridge once and only once, with the conditions that the islands could only be reached by the bridges and every bridge once

<sup>1</sup>These pictures are taken from Wikipedia

accessed must be crossed to its other end.

Since only the connection information is relevant, the shape of pictorial representations of a graph may be distorted in any way, without changing the graph itself. Thus it is enough to analyze the corresponding graph (on the left of Fig. 7). A closed walk which uses each edge only once is called an *Euler circuit*.

A key observation due to Euler is that whenever one enters a vertex by a bridge, one leaves the vertex by a bridge. In our terms, it means that if a graph has an Euler circuit, then a degree of every vertex has to be even. It sounds too easy, however, there is a remarkable result that this is the only condition for existence of an Euler circuit:

**Theorem 3.** (Leonard Euler, 1736)

*Let  $G$  be a finite connected graph. Then  $G$  has an Euler circuit if and only if all vertices of  $G$  have even degrees.*

We prove Theorem 3 later. We say that a walk in a graph  $G$  is an *Euler trail* if it uses every edge only once.

**Corollary 4.** *Let  $G$  be a finite connected graph. Then  $G$  has an Euler trail if and only if it has either two vertices of odd degree or no vertices of odd degree.*

**Proof.** Suppose that  $G$  has an Euler trail starting at  $v$  and ending at  $v'$ . If  $v = v'$ , the path is closed and Theorem 3 says that all vertices have even degree. If  $v \neq v'$ , we create a new edge  $e$  joining  $v$  and  $v'$ . The new graph  $G \cup \{e\}$  has an Euler circuit consisting of the Euler trail for  $G$  followed by  $e$ , so all vertices of  $G \cup \{e\}$  have even degree. Then we remove the edge  $e$ . Then  $v$  and  $v'$  are the only vertices of  $G = (G \cup \{e\}) \setminus \{e\}$  of odd degree.  $\square$

**Remark.** Returning to The Seven Bridges of Königsberg Problem, we see that there is no an Euler trail for the graph from Fig. 7. Indeed, all four vertices have odd degree.

### Finding an Euler Circuit

In order to prove Theorem 3, we would like to describe an algorithm how to find an Euler circuit if all vertices of  $G$  have even degrees. We start with an algorithm which finds a circuit which is not necessarily an Euler circuit, i.e. it may visit only once some of edges.

Let  $H = (V(H), E(H))$  be a graph with all vertices of even degree and let  $v \in V(H)$  be a vertex with positive even degree. For a graph  $G$  and an edge  $e$ , we define a graph  $G \setminus \{e\}$  which has exactly the same vertices as  $G$  and the same edges except given edge  $e$ . We say that the graph  $G \setminus \{e\}$  is given by removing  $e$  from  $E(G)$ . Here is the algorithm:

**Circuit** ( $H, v$ )

```
Choose an edge  $e$  with endpoint  $v$ 
Let  $P := (e)$  and remove  $e$  from  $E(H)$ 
while there is an edge at the terminal vertex of  $P$  do
    Choose such an edge  $e$  and add it to the path:
     $P := (P, e)$  and remove it from  $E(H)$ ,
return  $P$ 
```

**Exercise:** Analyze the algorithm **Circuit** ( $H, v$ ). Why does it produce a circuit?

Now we are ready for an algorithm which produces an Euler circuit.

**EulerCircuit**  $G = (V, E)$  ( $\deg v$  is even for each  $v \in V$ )

```
Choose a vertex  $v \in V(G)$ 
Let  $C := \mathbf{Circuit}(G, v)$ 
while  $\text{length}(C) < E(G)$  do
    Choose a vertex  $w$  in  $C$  of positive degree in  $G \setminus C$ .
    Attach  $\mathbf{Circuit}(G \setminus C, w)$  to  $C$  at  $w$  to obtain a longer circuit  $C$ .
return  $C$ 
```

**Exercise:** Analyze the algorithm **EulerCircuit** ( $G$ ). Why does it produce an Euler circuit?