



Summary on Lecture 6, April 8th, 2015

**Zero-one matrices and graphs.**

Let  $G = (V, E)$  be a directed graph (digraph), where  $|V| = n$ . Then a vertex  $e = (v, v') \in E$  is an ordered pair of vertices. Thus we can describe vertices as a binary relation on the set of vertices. Let us consider the following directed graphs:



(a) Digraph  $G_1$



(b) Digraph  $G_2$


Here we have the adjacency matrices  $M(G_1)$  and  $M(G_2)$  of these digraphs:

$$M(G_1) = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \end{bmatrix} \quad M(G_2) = \begin{bmatrix} 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \end{bmatrix}$$


**Partial order and Hasse diagrams.**

Let  $A$  be a set. We recall that a *partial order on  $A$*  is a binary relation  $\mathcal{R} \subset A \times A$  which is reflexive, antisymmetric and transitive. A useful tool to work with a partial order is a *Hasse diagram*. We give several examples.

**Examples.** (a) Let  $S$  be a set, and  $\mathcal{P}(S)$  be a set of all subsets of  $S$ . Then we define the relation: for  $A, B \in \mathcal{P}(S)$ ,  $A \leq B$  iff  $A \subseteq B$ . Here is a *Hasse diagram* of this partial order if  $S = \{1, 2, 3, 4\}$ .



(b) Let  $n = p_1^{e_1} \cdots p_k^{e_k}$  be a prime decomposition of a positive integer  $n$ . Let  $D(n)$  be the set of all divisors of  $n$ . Then every  $d \in D(n)$  has a form  $d = p_1^{a_1} \cdots p_k^{a_k}$ , where  $0 \leq a_i \leq e_i$  for each  $i = 1, \dots, k$ . We already have considered the following partial order on  $D(n)$ :  $d \leq d'$  iff  $d$  is a divisor of  $d'$ . Below is a Hasse diagram for this partial order if  $n = 2015$ .




Let  $\mathcal{R}$  be a partial order on  $A$ . We say that  $\mathcal{R}$  is a *linear (or total) order* on  $A$  if for any  $x, y \in A$  either  $(x, y) \in \mathcal{R}$  or  $(y, x) \in \mathcal{R}$ .

**Example.** Let  $A = \{1, 2, 2^2, \dots, 2^k\}$ , and  $(x \leq y)$  iff  $x$  is a divisor of  $y$ , i.e.,  $x|y$ . This is a linear order on  $A$ .

There are some practical applications of these concepts.

**Example.** Assume we would like to manufacture a product  $X$  (say, a toy). In order to do that there are several operations we have to perform according to the following Hasse diagram:



Nevertheless we have to organize the production in linear order since those operations could not be done at the same time. We select (from right to left) a “highest” leaf in that diagram, and we delete it. Then we recur. We obtain the following linear order:  $A < B < C < D < G < F < E$ . This new linear order provides a “linear” process to manufacture our product.

**Definition.** Let  $A$  a poset (i.e., a partial ordered set). We say that  $x \in A$  is maximal if  $x \leq a$  implies  $x = a$ . Similarly,  $y \in A$  is minimal if  $a \leq y$  implies  $a = y$ .

**Theorem 1.** Let  $A$  be a finite poset. Then there exists a maximal (minimal) element in  $A$ .

**Exercise.** Prove Theorem 1.

**Definition.** Let  $A$  a poset. An element  $x \in A$  is a greatest element if  $a \leq x$  for all  $a \in A$ . Similarly, an element  $y \in A$  is a least element if  $y \leq a$  for all  $a \in A$ .

**Theorem 2.** Let  $A$  be a poset. Assume there exists a greatest (least) element in  $A$ . Then it is unique.

**Exercise.** Prove Theorem 2.