

Summary on Lecture 7, April 10th, 2015

**Equivalence relations and partitions.**

Let  $A$  be a set. A family of subsets  $\{A_i\}_{i \in I}$ ,  $A_i \subseteq A$ , is called a partition if

$$A = \bigcup_{i \in I} A_i, \quad A_i \cap A_{i'} = \emptyset \text{ if } i \neq i'.$$

**Important example.** Let  $n \in \mathbf{Z}_+$  be a positive integer. We define an equivalence relation on  $\mathbf{Z}$  as follows:  $m \sim m'$  iff  $m - m'$  is divisible by  $n$ . Then we have  $n$  different classes of equivalent integers:

$$\begin{aligned} \mathbf{0} &:= \{0, \pm n, \pm 2 \cdot n, \dots\}, \\ \mathbf{1} &:= \{1, 1 \pm n, 1 \pm 2 \cdot n, \dots\}, \\ \mathbf{2} &:= \{2, 2 \pm n, 2 \pm 2 \cdot n, \dots\}, \\ \dots & \dots \dots \dots \\ \mathbf{n-1} &:= \{n-1, n-1 \pm n, n-1 \pm 2 \cdot n, \dots\}. \end{aligned}$$

We obtain that  $\mathbf{Z} = \bigcup_{i=0}^{n-1} \mathbf{i}$ , and clearly the sets  $\mathbf{i}$  and  $\mathbf{i}'$  do not intersect if  $i \neq i'$ . The set of equivalent classes  $\{\mathbf{0}, \mathbf{1}, \dots, \mathbf{n-1}\}$  is denoted by  $\mathbf{Z}/n$ . There are well-defined sum and product operations on  $\mathbf{Z}/n$ :

$$\mathbf{i} + \mathbf{i}' \quad \text{and} \quad \mathbf{i} \cdot \mathbf{i}'$$

Here are the addition and multiplication tables in  $\mathbf{Z}/5$ :

+	0	1	2	3	4
0	0	1	2	3	4
1	1	2	3	4	0
2	2	3	4	0	1
3	3	4	0	1	2
4	4	0	1	2	3

×	0	1	2	3	4
0	0	0	0	0	0
1	0	1	2	3	4
2	0	2	4	1	3
3	0	3	1	4	2
4	0	4	3	2	1

Next, we have the following addition and multiplication tables in  $\mathbf{Z}/6$ :

+	0	1	2	3	4	5
0	0	1	2	3	4	5
1	1	2	3	4	5	0
2	2	3	4	5	0	1
3	3	4	5	0	1	2
4	4	5	0	1	2	3
5	5	0	1	2	3	4

×	0	1	2	3	4	5
0	0	0	0	0	0	0
1	0	1	2	3	4	5
2	0	2	4	0	2	4
3	0	3	0	3	0	3
4	0	4	2	0	4	2
5	0	5	4	3	2	1

We notice that  $\mathbf{2} \cdot \mathbf{3} = \mathbf{0}$ ,  $\mathbf{4} \cdot \mathbf{3} = \mathbf{0}$ , and  $\mathbf{3} \cdot \mathbf{3} = \mathbf{3}$ .

Let  $\mathcal{R} \subset A \times A$  be an equivalence relation. For each element  $x \in A$  we define a subset

$$[x] = \{ y \in A \mid (x, y) \in \mathcal{R} \}$$

We notice that either  $[x] = [x']$  or  $[x] \cap [x'] = \emptyset$ . Indeed, assume that  $[x] \cap [x'] \neq \emptyset$ , and  $z \in [x] \cap [x']$ . Then  $(x, z) \in \mathcal{R}$  and  $(x', z) \in \mathcal{R}$ , and this implies that  $(x, x') \in \mathcal{R}$ , and thus  $[x] = [x']$ . We obtain that the family of sets  $\{[x]\}$  is a partition of  $A$ .

Now let  $\{A_i\}_{i \in I}$  be a partition of  $A$ . Then we define a relation  $\mathcal{R} \subset A \times A$  as follows:

$$(x, x') \in \mathcal{R} \text{ iff there exists } i \in I \text{ such that } x, x' \in A_i.$$

It is easy to check that  $\mathcal{R} \subset A \times A$  is an equivalence relation.

**Theorem 1.** Let  $A$  be a set. Then there is one-to-one correspondence between equivalence relations on  $A$  and partitions of  $A$ .

Now we would like to count all possible partitions of a finite set  $A = \{a_1, \dots, a_m\}$ . We'll say that a partition  $A = \bigcup_{i=1}^k A_i$  has a size  $k$ . Clearly,  $1 \leq k \leq m$ . We fix such  $k$  and count how many partitions of size  $k$  are there.

To get started, we can count how many onto functions are there  $f : A \rightarrow B$ , where  $B = \{b_1, \dots, b_k\}$ . Then we can think of  $b_i$  as a box to collect elements for  $A_i$ , thus we should forget the order of those boxes. By using the inclusion-exclusion principle, we obtain

$$\sum_{i=0}^{k-1} (-1)^i \binom{k}{i} (k-i)^m$$

onto functions are there  $f : A \rightarrow B$ . We divide by  $k!$  to obtain the Stirling number

$$S(m, k) = \frac{1}{k!} \sum_{i=0}^{k-1} (-1)^i \binom{k}{i} (k-i)^m.$$

Now we sum up partitions of  $A$  of all sizes. We obtain that there are

$$\sum_{k=1}^m S(m, k) = \sum_{k=1}^m \left( \frac{1}{k!} \sum_{i=0}^{k-1} (-1)^i \binom{k}{i} (k-i)^m \right)$$

partitions of  $A = \{a_1, \dots, a_m\}$ .