

**CORRIGENDA TO ‘HECKE-CLIFFORD SUPERALGEBRAS,
CRYSTALS OF TYPE $A_{2\ell}^{(2)}$ AND MODULAR BRANCHING
RULES FOR \widehat{S}_n ’**

JONATHAN BRUNDAN AND ALEXANDER KLESHCHEV

We are grateful to Steffen Koenig and Steffen Oppermann for pointing out that there is a gap in the proof of Lemma 5.22 of [1]. We do not know at the moment whether Lemma 5.22 is correct or not. However, we claim that it is not needed anywhere in the paper if the following changes are made.

- 1) Drop Lemma 5.22.
- 2) Amend Lemmas 5.23 and 6.16 as follows.

Lemma 5.23. *Take $i, j \in I$ with $i \neq j$ and set $k = -\langle h_i, \alpha_j \rangle$. Let M be an irreducible module in $\text{Rep}_I \mathcal{H}_n$.*

- (i) *There exists a unique integer a with $0 \leq a \leq k$ such that for every $m \geq 0$ we have*

$$\varepsilon_i(\tilde{f}_i^m \tilde{f}_j M) = m + \varepsilon_i(M) - a.$$

- (ii) *Assume $m \geq k$. Then a copy of $\tilde{f}_i^m \tilde{f}_j M$ appears in the cosocle of*

$$\text{ind } \tilde{f}_i^{m-k} M \otimes L(i^a j i^{k-a}).$$

- (iii) *Assume $0 \leq m < k \leq m + \varepsilon$. Then a copy of $\tilde{f}_i^m \tilde{f}_j M$ appears in the cosocle of*

$$\text{ind } \tilde{e}_i^{k-m} M \otimes L(i^a j i^{k-a}).$$

Proof. Let $\varepsilon = \varepsilon_i(M)$ and write $M = \tilde{f}_i^\varepsilon N$ for irreducible $N \in \text{Rep}_I \mathcal{H}_{n-\varepsilon}$ with $\varepsilon_i(N) = 0$. It suffices to prove (i) for any fixed choice of m , the conclusion for all other $m \geq 0$ then follows immediately by (5.11). So take $m \geq 0$ with $k \leq m + \varepsilon$. Note that $\tilde{f}_i^m \tilde{f}_j M = \tilde{f}_i^m \tilde{f}_j \tilde{f}_i^\varepsilon N$ is a quotient of

$$\begin{cases} \text{ind } N \otimes L(i^\varepsilon) \otimes L(j) \otimes L(i)^{\otimes k} \otimes L(i^{m-k}) & \text{if } m \geq k, \\ \text{ind } N \otimes L(i^{m+\varepsilon-k}) \otimes L(i)^{\otimes (k-m)} \otimes L(j) \otimes L(i^m) & \text{if } m < k, \end{cases}$$

which by Lemma 5.19 has a filtration with factors isomorphic to

$$\begin{cases} F_a := \text{ind } N \otimes L(i^\varepsilon) \otimes L(i^a j i^{k-a}) \otimes L(i^{m-k}) & \text{if } m \geq k, \\ F_a := \text{ind } N \otimes L(i^{m+\varepsilon-k}) \otimes L(i^a j i^{k-a}) & \text{if } m < k, \end{cases}$$

for $0 \leq a \leq k$, each appearing with some multiplicity. So $\tilde{f}_i^m \tilde{f}_j M$ is a quotient of some such factor, and to prove (i) it remains to show that $\varepsilon_i(L) = \varepsilon + m - a$ for any irreducible quotient L of F_a . The inequality $\varepsilon_i(L) \leq \varepsilon + m - a$ is clear from the Shuffle Lemma. On the other hand, by transitivity of induction and Lemma 5.21, $F_a \cong \text{ind } N \otimes (\text{ind } L(i^a j i^{k-a}) \otimes L(i^{\varepsilon+m-k}))$.

So by Frobenius reciprocity, the irreducible module $N \otimes (\text{ind } L(i^a j i^{k-a}) \otimes L(i^{\varepsilon+m-k}))$ is contained in $\text{res}_{n-\varepsilon, m+1+\varepsilon} L$. Hence $\varepsilon_i(L) \geq \varepsilon + m - a$.

To complete the proof of (ii) and (iii), by Lemma 5.21, we also have $F_a \cong \text{ind } N \otimes L(i^{\varepsilon+m-k}) \otimes L(i^a j i^{k-a})$, and by the Shuffle Lemma, the only composition factors K of F_a with $\varepsilon_i(K) = \varepsilon + m - a$ come from its quotient

$$\text{ind } \tilde{f}_i^{m-k+\varepsilon} N \otimes L(i^a j i^{k-a}).$$

The latter is $\text{ind } \tilde{f}_i^{m-k} M \otimes L(i^a j i^{k-a})$ if $m \geq k$ and $\text{ind } \tilde{e}_i^{k-m} M \otimes L(i^a j i^{k-a})$ otherwise. \square

Lemma 6.16 *Let $i, j \in I$ with $i \neq j$. Let M be an irreducible module in $\text{Rep } \mathcal{H}_n^\lambda$ such that $\varphi_j(M) > 0$. Then, $\varphi_i(\tilde{f}_j M) - \varepsilon_i(\tilde{f}_j M) \leq \varphi_i(M) - \varepsilon_i(M) - \langle h_i, \alpha_j \rangle$.*

Proof. Let $\varepsilon = \varepsilon_i(M)$, $\varphi = \varphi_i(M)$ and $k = -\langle h_i, \alpha_j \rangle$. By Lemma 5.23, there exist unique $a, b \geq 0$ with $a + b = k$ such that $\varepsilon_i(\tilde{f}_j M) = \varepsilon - a$. We need to show that $\varphi_i(\tilde{f}_j M) \leq \varphi + b$, which follows if we can show that $\text{pr}^\lambda \tilde{f}_i^m \tilde{f}_j M = 0$ for all $m > \varphi + b$. We claim that

$$\varepsilon_i^*(\tilde{f}_i^m \tilde{f}_j M) \geq \varepsilon_i^*(\tilde{f}_i^{m-b} M)$$

for all $m > \varphi + b$. Given the claim, we know by the definition of φ , Corollary 6.13 and Lemma 6.15 that $\varepsilon_i^*(\tilde{f}_i^{m-b} M) > \langle h_i, \lambda \rangle$ for all $m > \varphi + b$. So the claim implies that $\varepsilon_i^*(\tilde{f}_i^m \tilde{f}_j M) > \langle h_i, \lambda \rangle$ for all $m > \varphi + b$, hence by Corollary 6.13 once more, $\text{pr}^\lambda \tilde{f}_i^m \tilde{f}_j M = 0$ as required.

To prove the claim, note that $k \leq m + \varepsilon$, so by Lemma 5.23(ii),(iii) that there is a surjection

$$\text{ind}_{n-\varepsilon, \varepsilon+m-k, k+1}^{n+m+1} N \otimes L(i^{\varepsilon+m-k}) \otimes L(i^a j i^b) \twoheadrightarrow \tilde{f}_i^m \tilde{f}_j M,$$

where $N = \tilde{e}_i^\varepsilon M$. By Lemma 5.19, $\text{res}_{a, b+1}^{a+b+1} L(i^a j i^b) \cong L(i^a) \otimes L(j i^b)$. Hence there is a surjection $\text{ind}_{a, b+1}^{a+b+1} L(i^a) \otimes L(j i^b) \twoheadrightarrow L(i^a j i^b)$. Combining, we have proved existence of a surjection

$$\text{ind}_{n-\varepsilon, \varepsilon+m-b, b+1}^{n+m+1} N \otimes L(i^{\varepsilon+m-b}) \otimes L(j i^b) \twoheadrightarrow \tilde{f}_i^m \tilde{f}_j M.$$

Hence by Frobenius reciprocity there is a non-zero map

$$(\text{ind}_{n-\varepsilon, \varepsilon+m-b}^{n+m-b} N \otimes L(i^{\varepsilon+m-b})) \otimes L(j i^b) \rightarrow \text{res}_{n+m-b, b+1}^{n+m+1} \tilde{f}_i^m \tilde{f}_j M.$$

Since the left-hand module has irreducible cosocle $\tilde{f}_i^{m-b} M \otimes L(j i^b)$, we deduce that $\tilde{f}_i^m \tilde{f}_j M$ has a constituent isomorphic to $\tilde{f}_i^{m-b} M$ on restriction to the subalgebra $\mathcal{H}_{n+m-b} \subseteq \mathcal{H}_{n+m+1}$. This implies the claim. \square

REFERENCES

- [1] J. Brundan and A. Kleshchev, Hecke-Clifford superalgebras, crystals of type $A_{2\ell}^{(2)}$ and modular branching rules for \hat{S}_n , *Representation Theory* **5** (2001), 317–403.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403, USA
E-mail address: klesh@uoregon.edu, brundan@uoregon.edu