

CYCLOTOMIC NIL-BRAUER AND SINGULAR SOERGEL BIMODULES OF TYPE D

ELIJAH BODISH, JONATHAN BRUNDAN, AND BEN ELIAS

ABSTRACT. We introduce a new family of monoidal categories which are cyclotomic quotients of the nil-Brauer category. We construct a monoidal functor from the cyclotomic nil-Brauer category of level l to another monoidal category constructed from singular Soergel bimodules of type $A_{l-1} \setminus D_l / A_{l-1}$. We conjecture that our functor is an equivalence of categories. Although we can prove neither fullness nor faithfulness at this point, we are able to show that the functor induces an isomorphism at the level of Grothendieck rings. We compute these rings and their canonical bases over any field of characteristic different from 2, and give diagrammatic descriptions of the corresponding primitive idempotents.

CONTENTS

1. Introduction	2
2. The cyclotomic nil-Brauer category	6
2.1. Symmetric functions	6
2.2. The nil-Brauer category	7
2.3. Identification of the Grothendieck ring of \mathbf{NB}_l	9
2.4. Bubble slides	11
2.5. The cyclotomic nil-Brauer category	11
2.6. The module $V(l)$ and its l -canonical basis	13
2.7. The special cases $l = 0$ and $l = 1$	15
3. Reminders about singular Soergel bimodules	16
3.1. Coxeter group realization	16
3.2. Chains of Frobenius extensions	17
3.3. Diagrammatics for singular Bott-Samelson bimodules	19
3.4. The Soergel-Williamson categorification theorem	22
4. Extended singular Soergel bimodules in type D_l	27
4.1. Realization of the root system of type D_l	27
4.2. Motivational interlude	28
4.3. The extended q -Schur algebra	28
4.4. The subalgebra V and its Kazhdan-Lusztig basis	29
4.5. Extended singular Soergel bimodules	32
4.6. Diagrammatics for extended singular Bott-Samelson bimodules	35
4.7. Calculations	37
5. Main results	48
5.1. Construction of the monoidal functor Θ	48
5.2. Characters of indecomposables	51
5.3. Identification of the Grothendieck ring of \mathbf{cNB}_l	53
5.4. Diagrams for primitive idempotents	53
References	53

2020 *Mathematics Subject Classification.* 17B37, 18N25, 16D20.

Key words and phrases. nil-Brauer category, singular Soergel bimodules, categorification.

E.B. is partially supported by NSF grant MSPRF-2202897. J.B. is supported in part by NSF grant DMS-2348840. B.E. is supported in part by NSF grant DMS-2201387, and his research group is supported by DMS-2039316.

1. INTRODUCTION

Let \mathbb{k} be a field with $\text{char } \mathbb{k} \neq 2$. The *nil-Brauer category* \mathbf{NB}_t is a graded \mathbb{k} -linear monoidal category introduced in [BWW24] (see Definition 2.3). In [BWW23], it was shown to categorify the split t -quantum group of rank one, which is the simplest of the coideal subalgebras of quantized enveloping algebras introduced by Letzter [Let99] corresponding to symmetric pairs. There are two admissible choices for the parameter t in the definition of \mathbf{NB}_t , 0 or 1, corresponding to the two possible $\mathbb{Z}[q, q^{-1}]$ -forms, U'_0 or U'_1 , for this particular t -quantum group according to the theory developed by Bao and Wang [BW18a]. In this paper, we take the next step in the study of \mathbf{NB}_t , introducing monoidal categories \mathbf{cNB}_l for $l \in \mathbb{N}$ with $l \equiv t \pmod{2}$, which are the *cyclotomic quotients* of \mathbf{NB}_t .

Consider the irreducible $U_q(\mathfrak{sl}_2)$ module of highest weight l , with its usual $\mathbb{Z}[q, q^{-1}]$ -form $V(l)$. It is a cyclic U'_t -module generated by its highest weight vector, hence, it is isomorphic to a quotient of U'_t . Since U'_t is itself commutative, this gives $V(l)$ structure both as a U'_t -module and as a commutative $\mathbb{Z}[q, q^{-1}]$ -algebra. These considerations also equip $V(l)$ with a distinguished basis, the *t -canonical basis*, which was computed explicitly by Berman and Wang in [BW18c].

The starting point for our work was the observation that the formulae in [BW18c] match the Kazhdan-Lusztig combinatorics in a piece of the q -Schur algebra of type D_l , associated to its maximal parabolics of type A_{l-1} . Categorifying this observation, we construct a monoidal functor from \mathbf{cNB}_l to a monoidal category consisting of singular Soergel bimodules of type $A_{l-1} \setminus D_l / A_{l-1}$. This functor is a powerful tool enabling the computation of the Grothendieck group of (the graded Karoubi envelope of) \mathbf{cNB}_l . We show that it is isomorphic to $V(l)$ as a $\mathbb{Z}[q, q^{-1}]$ -algebra, and match the canonical basis arising from indecomposable objects with the t -canonical basis.

Ordinary quantum groups are categorified by the Kac-Moody 2-categories of Khovanov, Lauda and Rouquier [KL10, Rou08]. In that setting, cyclotomic quotients are certain universal 2-representations which categorify the integrable highest weight modules of the underlying quantum group. They play a central role in the theory of categorical Kac-Moody actions developed in [Rou08]. The cyclotomic quotients of \mathbf{NB}_t are expected to play a similar role in the study of categorical actions of the split t -quantum group of rank one. From this perspective, it may seem surprising that the cyclotomic quotients of \mathbf{NB}_t are monoidal categories, rather than merely being module categories, but this reflects the commutativity of the t -quantum group in the split rank one case.

In order to formulate our results in more detail, we need to recall some elementary aspects of the theory of singular Soergel bimodules. To set conventions, here is the Dynkin diagram (with its diagram automorphism γ) for the Weyl group W of type D_l , alongside the Dynkin diagram of the slightly larger Weyl group of type B_l :



We include the cases $l = 2$ and $l = 3$, identifying D_2 with $A_1 \times A_1$ and D_3 with A_3 . The Weyl groups of type D_l and B_l both act by automorphisms on the polynomial algebra $\mathbb{k}[x_1, \dots, x_l]$, which is the coordinate algebra of the standard reflection representation. Explicitly, s_i for $i = 1, \dots, l-1$ switches x_i and x_{i+1} while fixing the other generators, s_0 maps $x_1 \mapsto -x_1$ while fixing the other generators, and $s_{-1} = s_0 s_1 s_0$. We view $\mathbb{k}[x_1, \dots, x_l]$ as a graded algebra so that each x_i is in degree 2.

Let \mathfrak{gBim} be the graded \mathbb{k} -linear bicategory with objects, 1-morphisms and 2-morphisms that are graded \mathbb{k} -algebras, graded bimodules and graded bimodule homomorphisms, respectively. There is

a full sub-bicategory $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}$ of $\mathfrak{g}\mathfrak{B}\mathfrak{i}\mathfrak{m}$ consisting of *singular Bott-Samelson bimodules* (see Definition 3.3). The objects of $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}$ are the algebras of invariants $\mathbb{k}[x_1, \dots, x_l]^{W_I}$ for all parabolic subgroups $W_I \leq W$, and its 1-morphisms are tensor-generated by induction and restriction bimodules. The graded Karoubi envelope $\mathfrak{g}\mathfrak{K}\mathfrak{a}\mathfrak{r}(\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m})$ is the bicategory of singular Soergel bimodules of type D in the title. By general results of Soergel and Williamson [Wil11] recalled in Theorems 3.10 to 3.12 below, the Grothendieck ring of $\mathfrak{g}\mathfrak{K}\mathfrak{a}\mathfrak{r}(\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m})$ may be identified with the natural integral form S of the q -Schur algebra of type D_l . In fact, we will work in this paper with a mildly extended version of $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}$, denoted $\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}$, which is obtained by incorporating additional bimodules which allow for twisting by the graph automorphism γ (see Definition 4.10).

Now we focus on the monoidal category $\mathbf{End}_{\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}}(A)$ which is the endomorphism category of a particular object A of the bicategory $\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}$, namely, the algebra

$$A := \mathbb{k}[x_1, \dots, x_l]^{S_l}$$

of symmetric polynomials viewed as an object of $\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}$ by identifying S_l with the maximal parabolic subgroup of W obtained by deleting $\{-1\}$ from the Dynkin diagram. Thus, $\mathbf{End}_{\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}}(A)$ is a full monoidal subcategory of the graded \mathbb{k} -linear monoidal category of graded (A, A) -bimodules. Let $S_1 \times S_{l-1} < S_l$ denote the parabolic subgroup obtained by deleting $\{-1, 1\}$ from the Dynkin diagram, and $B \in \mathbf{ob} \mathbf{End}_{\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}}(A)$ be the graded (A, A) -bimodule $\mathbb{k}[x_1, \dots, x_l]^{S_1 \times S_{l-1}}$, with grading down-shifted so that the identity is in degree $(1 - l)$. The right action of A on B is the obvious one arising from the inclusion of algebras

$$\mathbb{k}[x_1, \dots, x_l]^{S_l} \subset \mathbb{k}[x_1, \dots, x_l]^{S_1 \times S_{l-1}},$$

while the left action is defined by restriction along the *twisted* embedding $f \mapsto s_0(f)$ induced by the graph automorphism γ . The bimodule B generates $\mathbf{End}_{\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}}(A)$ in the sense that any bimodule in $\mathbf{End}_{\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}}(A)$ is isomorphic (up to grading shift) to a summand of a tensor power

$$B^{\otimes n} = B \otimes_A \cdots \otimes_A B \quad (0 \leq n \leq l).$$

We note that $\mathbf{End}_{\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}}(A)$ is actually a C -linear graded monoidal category, where

$$C := \mathbb{k}[x_1^2, \dots, x_l^2]^{S_l} \subset A$$

is the full algebra of invariants of the Weyl group of type B_l .

The cyclotomic nil-Brauer category \mathbf{cNB}_l is another graded C -linear monoidal category defined as the quotient of the nil-Brauer category \mathbf{NB}_l by the two-sided tensor ideal generated by some explicit *cyclotomic relations* (see Definition 2.10 and Theorem 2.11). It has one generating object, also denoted B , and generating morphisms that are represented string-diagrammatically by the dot $\downarrow : B \rightarrow B$, the crossing $\times : B \star B \rightarrow B \star B$ (which squares to 0), the cup $\cup : \mathbb{1} \rightarrow B \star B$ and the cap $\cap : B \star B \rightarrow \mathbb{1}$ (these satisfy the zig-zag relations).

Main Theorem. *There is a C -linear graded monoidal functor $\Theta : \mathbf{cNB}_l \rightarrow \mathbf{End}_{\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}}(A)$ mapping the generating object B of \mathbf{cNB}_l to the (A, A) -bimodule B . Moreover:*

- (1) *The functor Θ induces an isomorphism $K_0(\mathfrak{g}\mathfrak{K}\mathfrak{a}\mathfrak{r}(\mathbf{cNB}_l)) \xrightarrow{\sim} K_0(\mathfrak{g}\mathfrak{K}\mathfrak{a}\mathfrak{r}(\mathbf{End}_{\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}}(A)))$ between Grothendieck rings of graded Karoubi envelopes of these monoidal categories.*
- (2) *Both Grothendieck rings are isomorphic to the $\mathbb{Z}[q, q^{-1}]$ -form $\mathcal{V}(l)$ of the $(l + 1)$ -dimensional irreducible $U_q(\mathfrak{sl}_2)$ -module equipped with an explicit algebra structure.*
- (3) *Up to grading shift, isomorphism classes both of indecomposable objects of $\mathfrak{g}\mathfrak{K}\mathfrak{a}\mathfrak{r}(\mathbf{cNB}_l)$ and of indecomposable bimodules in $\mathfrak{g}\mathfrak{K}\mathfrak{a}\mathfrak{r}(\mathbf{End}_{\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}}(A))$ recover the t -canonical basis of $\mathcal{V}(l)$.*

The construction of the monoidal functor Θ is naive—we simply write down bimodule homomorphisms in $\mathbf{End}_{\mathfrak{e}\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}}(A)$ that are the images of the generating morphisms of \mathbf{cNB}_l then check that they

satisfy the defining relations. The endomorphism $\Theta(\downarrow) : B \rightarrow B$ is defined by right multiplication by x_1 , and $\Theta(\cup)$ and $\Theta(\cap)$ are homomorphisms which are also quite easy to describe explicitly:

$$\begin{aligned} \Theta(\cup) : A \rightarrow B \otimes_A B, & & 1 \mapsto \sum_{r=0}^{l-1} e_r(x_2, \dots, x_l) \otimes x_1^{l-1-r}, \\ \Theta(\cap) : B \otimes_A B \rightarrow A, & & f \otimes g \mapsto \partial_{l-1} \cdots \partial_2 \partial_1 (s_0(f)g), \end{aligned}$$

where $e_r(x_2, \dots, x_l)$ is the r th elementary symmetric polynomial in these variables and ∂_i is the Demazure operator $f \mapsto \frac{f - s_i(f)}{x_i - x_{i+1}}$. However, $\Theta(\times)$, which is an endomorphism of the bimodule $B \otimes_A B$, is more difficult and we are only able to write it down using the diagrammatic calculus for $\mathfrak{B} \subseteq \mathfrak{Bim}$ from [ESW17], extended in an obvious way to $e\mathfrak{B} \subseteq \mathfrak{Bim}$. The picture for the crossing, using conventions which will be explained fully in the main body of the text, is

$$\Theta(\times) = + \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} + = + \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} + .$$


When $l = 2$, the circle and all labels $\hat{2}$ should be omitted from this picture. In Theorem 5.1, we also draw the much simpler pictures representing the images of the other generators.

Our Main Theorem gives complete information about the structure of the (A, A) -bimodules $B^{\otimes n}$. Up to isomorphism and grading shift, the indecomposable graded (A, A) -bimodules that arise as summands are parametrized by the set $\{0, 1, \dots, l\}$. The n th one, denoted $B^{[n]}$ for $0 \leq n \leq l$, is uniquely determined by the property that it appears as a summand of $B^{\otimes n}$ with graded multiplicity $[n]_q!$, and all other indecomposable summands of $B^{\otimes n}$ are isomorphic to degree shifts of $B^{[m]}$ for $m < n$ with $m \equiv n \pmod{2}$. Moreover, the character of $B^{[n]}$ is independent of the characteristic of the ground field.

We also give an explicit diagrammatic description of the homogeneous primitive idempotent \mathbf{f}_n in $\text{End}_{A-A}(B^{\otimes n})$ that is the projection of $B^{\otimes n}$ onto the unique summand that is $B^{[n]}$ shifted up in degree by $\binom{n}{2}$. For example, assuming that n is even with $2 \leq n \leq l-1$, the string diagram for \mathbf{f}_n is

$$+ \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} + \dots + \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} +$$

In this picture, there are $(n - 1)$ cups at the top and $(n - 1)$ caps at the bottom. There is a similar picture when n is odd. When $n = l - 1$, the big bubble can be “popped”—the penultimate \mathbf{f}_n is also equal to



which has been drawn again assuming that n is even. A similar bubble-free diagram also represents the primitive idempotent \mathbf{f}_n in the ultimate case $n = l$. These statements are proven using the functor Θ at the same time as establishing a parallel diagrammatic description of the corresponding primitive idempotents \mathbf{e}_n ($0 \leq n \leq l$) in \mathbf{cNB}_l . An essential ingredient is the description of primitive idempotents in \mathbf{NB}_l discovered in [BWW23] (see Theorem 2.6).

Conjecture. *The monoidal functor Θ from the Main Theorem is fully faithful, hence, it induces an equivalence between the graded Karoubi envelopes of \mathbf{cNB}_l and $\mathbf{End}_{\mathbf{eB} \otimes \mathbf{Bim}}(A)$.*

Despite the elementary nature of these categories, we do not have a proof of this conjecture at this point. Via the Soergel-Williamson categorification theorem, the graded dimensions of morphism spaces in $\mathbf{End}_{\mathbf{eB} \otimes \mathbf{Bim}}(A)$ can be computed using Lusztig’s symmetric bilinear form on $V(l)$. However, we are not able to compute these dimensions on the \mathbf{cNB}_l side (although of course this would follow from the truth of our conjecture). One problem is that $V(l)$ is not irreducible as a U_l^+ -module, so bilinear forms on $V(l)$ with the appropriate adjunction properties are not uniquely determined. In subsequent work, we hope to be able to extend the results here from the split ι -quantum group of rank one to categorifications of all quasi-split ι -quantum groups of type AIII with an odd number of nodes in the Satake diagram; the ones with an even number of nodes were categorified already in [BSWW18]. This may allow us to make progress towards proving the conjecture. It seems likely that any proof of the conjecture will also produce explicit bases for the morphism spaces $\mathrm{Hom}_{A \cdot A}(A, B^{[2n]})$ ($0 \leq n \leq \lfloor l/2 \rfloor$) (see Corollary 5.4 for the graded dimension of this space). We expect such bases will be essential for any future developments related to categorification of braid group actions of the split ι -quantum group of rank one.

General conventions. We use the word “graded” to denote a category (or an object of a category) that is enriched in \mathbf{gVec} , the symmetric monoidal category of \mathbb{Z} -graded vector spaces over the ground field \mathbb{k} . We denote the graded bicategory of graded algebras, graded bimodules and graded natural transformations by \mathbf{gBim} . In a graded category, we use the symbol \cong to denote isomorphism of degree 0. We use \otimes to denote tensor product over the field \mathbb{k} and \star to denote the tensor product in other graded monoidal categories or bicategories. For graded vector spaces V and W , we have by definition that $\mathrm{Hom}_{\mathbb{k}}(V, W) = \bigoplus_{n \in \mathbb{Z}} \mathrm{Hom}_{\mathbb{k}}(V, W)_n$, with $f \in \mathrm{Hom}(V, W)_n$ mapping V_i into W_{i+n} . The downward grading shift functor on \mathbf{gVec} is denoted q , i.e., $(qV)_n = V_{n+1}$. Assuming all V_n are finite-dimensional, its *graded dimension* is

$$\dim_q V := \sum_{n \in \mathbb{Z}} q^{-n} \dim V_n.$$

Graded rank of a free graded module over a graded algebra is defined similarly. For a series $f = \sum_{n \in \mathbb{Z}} a_n q^n$ with each $a_n \in \mathbb{N}$, fV denotes $\bigoplus_{n \in \mathbb{Z}} q^n V^{\oplus a_n}$. An additive map between $\mathbb{Z}[q, q^{-1}]$ -modules is said to be *anti-linear* if it twists scalars by the *bar involution* $- : \mathbb{Z}[q, q^{-1}] \rightarrow \mathbb{Z}[q, q^{-1}], q \mapsto q^{-1}$. Let $[n]_q$ be the quantum integer $\frac{q^n - q^{-n}}{q - q^{-1}}$, and $[n]_q!$ and $\begin{bmatrix} n \\ r \end{bmatrix}_q$ be the corresponding quantum factorial and quantum binomial coefficient.

Acknowledgements. The second author thanks Weiqiang Wang for helpful discussions about the underlying t -Schur-Weyl duality. All authors thank the Okinawa Institute of Science and Technology for its hospitality in June 2023, when this project was initiated.

2. THE CYCLOTOMIC NIL-BRAUER CATEGORY

Throughout the section, $t \in \{0, 1\}$ is a fixed parameter. We begin by defining a modified version \mathbf{NB}_t of the nil-Brauer category from [BWW24], and explaining its relationship to a $\mathbb{Z}[q, q^{-1}]$ -form U_t^l for the split t -quantum group of rank one, following [BWW23]. Then, we pass to the quotient of \mathbf{NB}_t by a certain two-sided tensor ideal \mathbf{I}_l for $l \in \mathbb{N}$ with $l \equiv t \pmod{2}$, thereby defining the cyclotomic nil-Brauer category of level l , denoted \mathbf{cNB}_l . Finally, we discuss the combinatorics of the t -canonical basis for a $\mathbb{Z}[q, q^{-1}]$ -form $V(l)$ for the $(l+1)$ -dimensional irreducible $U_q(\mathfrak{sl}_2)$ -module, which was worked out originally in [BW18c].

2.1. Symmetric functions. Let Λ be the algebra of symmetric functions over \mathbb{k} . It is freely generated by either the elementary symmetric polynomials e_r ($r \geq 1$) or the complete symmetric polynomials h_r ($r \geq 1$), and $e_0 = h_0 := 1$. We work with the grading on Λ in which e_r and h_r are of degree $2r$. As usual when working with symmetric functions, we use generating functions, setting

$$e(u) := \sum_{r \geq 0} e_r u^{-r}, \quad h(u) := \sum_{r \geq 0} h_r u^{-r}, \quad (2.1)$$

viewed as elements of $1 + u^{-1}\Lambda[[u^{-1}]]$ for a formal variable u . Then we have that $e(-u)h(u) = 1$. Let Γ be the subalgebra of Λ generated by Schur's q -functions

$$q_r := \sum_{s=0}^r h_s e_{r-s} \quad (2.2)$$

for $r \geq 0$. We have that

$$q(u) := \sum_{r \geq 0} q_r u^{-r} = e(u)h(u). \quad (2.3)$$

Hence, $q(u)q(-u) = 1$. Recall also that $2(-1)^{r-1}q_{2r} = q_r^2 + 2\sum_{s=1}^{r-1}(-1)^{r-s}q_s q_{2r-s}$ for $r \geq 1$; cf. [Mac15, (III.8.2')]. Using this, it follows that Γ is freely generated by q_{2r-1} ($r \geq 1$).

Let $e^{[2]}(u) := e(u)e(-u)$. Since $e^{[2]}(u) = e^{[2]}(-u)$, its expansion as a formal power series only involves even powers of u , so we have that

$$e^{[2]}(u) = e(u)e(-u) = \sum_{r \geq 0} (-1)^r e_r^{[2]} u^{-2r}. \quad (2.4)$$

Equating coefficients gives that $e_r^{[2]} = 2(-1)^r e_{2r} + e_r^2 + 2\sum_{s=1}^{r-1}(-1)^{r-s} e_s e_{2r-s}$. Let $\Lambda^{[2]}$ be the subalgebra of Λ generated by $e_r^{[2]}$ ($r \geq 1$).

Lemma 2.1. *The symmetric functions $e_r^{[2]}$ ($r \geq 1$) are algebraically independent. Moreover, multiplication defines an algebra isomorphism $\Gamma \otimes \Lambda^{[2]} \xrightarrow{\sim} \Lambda$.*

Proof. The first statement follows because $e_r^{[2]} = 2(-1)^r e_{2r} + (\text{a linear combination of monomials in } e_s \text{ for } s < 2r)$. Also, because $h_r = (-1)^{r-1} e_r + (\text{a linear combination of monomials in } e_s \text{ for } s < r)$, the definition (2.2) implies that $q_{2r-1} = 2e_{2r-1} + (\text{a linear combination of monomials in } e_s \text{ for } s < 2r - 1)$. Everything else now follows because Λ is freely generated by e_r ($r \geq 1$). \square

Remark 2.2. The symmetric functions $e_r^{[2]}$ are natural to consider from the point of view of symmetric polynomials. To explain what we mean, let A denote the algebra of symmetric polynomials $\mathbb{k}[x_1, \dots, x_l]^{S_l}$ for some fixed $l \geq 0$, and

$$\text{ev}_l : \Lambda \rightarrow A \quad (2.5)$$

be the homomorphism sending e_r to the r th elementary symmetric polynomial $e_r(x_1, \dots, x_l)$ (which is 0 for $r > l$). Then we have that

$$\text{ev}_l(e_r^{[2]}) = e_r(x_1^2, \dots, x_l^2). \quad (2.6)$$

To prove this, note that $\text{ev}_l(e(u)) = u^{-l}(u + x_1) \cdots (u + x_l)$ and $\text{ev}_l(e(-u)) = u^{-l}(u - x_1) \cdots (u - x_l)$. Hence, we have that $\text{ev}_l(e(u)e(-u)) = u^{-2l}(u^2 - x_1^2) \cdots (u^2 - x_l^2)$. The u^{-2r} -coefficient of this expression is $(-1)^r e_r(x_1^2, \dots, x_l^2)$.

2.2. The nil-Brauer category. In [BWW24, Def. 2.1], the nil-Brauer category was defined over the ground field \mathbb{k} . In this article, we will work with a modified version which is the $\Lambda^{[2]}$ -linear monoidal category obtained by extending scalars to the ground ring $\Lambda^{[2]}$. To explain the definition, we use the string calculus for monoidal categories, adopting the standard conventions from most of the recent categorification literature, as in [BWW24].

Definition 2.3. Let $\Lambda^{[2]}$ be the subalgebra of Λ defined in Lemma 2.1. Let $t \in \{0, 1\}$. The *nil-Brauer category* \mathbf{NB}_t is the strict graded $\Lambda^{[2]}$ -linear monoidal category with one generating object B , whose identity endomorphism id_B is denoted by the string $|$, and four generating morphisms

$$\begin{array}{cccc} \downarrow : B \rightarrow B, & \times : B \star B \rightarrow B \star B, & \cap : B \star B \rightarrow \mathbb{1}, & \cup : \mathbb{1} \rightarrow B \star B, \\ \text{(degree 2)} & \text{(degree -2)} & \text{(degree 0)} & \text{(degree 0)} \end{array}$$

subject to the following relations:

$$\downarrow = | = \uparrow, \quad \bigcirc = t \text{id}_1, \quad (2.7)$$

$$\downarrow \downarrow = \downarrow \downarrow, \quad \bigcirc = 0, \quad (2.8)$$

$$\downarrow \downarrow = - \downarrow \downarrow, \quad \times - \times = | | - \cup \cup, \quad (2.9)$$

$$\downarrow \downarrow = 0, \quad \downarrow \downarrow = \downarrow \downarrow. \quad (2.10)$$

We will denote the r th power of \downarrow by labeling the dot with r . Using the zig-zag relations from (2.7), the following are easily deduced from (2.8) and (2.9):

$$\downarrow \downarrow = \downarrow \downarrow, \quad \downarrow \downarrow = 0, \quad (2.11)$$

$$\downarrow \downarrow = - \downarrow \downarrow, \quad \times - \times = | | - \cup \cup. \quad (2.12)$$

Now it follows that there are strict graded $\Lambda^{[2]}$ -linear monoidal functors

$$\mathbb{R} : \mathbf{NB}_t \rightarrow \mathbf{NB}_t^{\text{rev}}, \quad s \mapsto (-1)^{\bullet(s)} s^{\leftrightarrow}, \quad (2.13)$$

$$\mathbb{T} : \mathbf{NB}_t \rightarrow \mathbf{NB}_t^{\text{op}}, \quad s \mapsto s^\dagger. \quad (2.14)$$

Here, the op denotes the opposite category with the same monoidal product, and rev denotes the same category with the reverse monoidal product. Also, for a string diagram s we are using s^\dagger and s^{\leftrightarrow} to denote its reflection in a horizontal or vertical axis, and $\bullet(s)$ denotes the total number of dots in the diagram. It follows that the category \mathbf{NB}_t has a strict pivotal structure. The underlying duality functor is $\mathbb{R} \circ \mathbb{T} = \mathbb{T} \circ \mathbb{R}$, which rotates a string diagram s through 180° then scales by $(-1)^{\bullet(s)}$. For more details about all of this, see [BWW24, Sec. 2].

Lemma 2.4. *Using the defining relations (2.7) to (2.9) but neither of the relations (2.10), it follows that*

$$\begin{array}{c} \text{X} \\ \text{X} \end{array} = \begin{array}{c} \text{X} \\ \text{X} \end{array}, \quad \begin{array}{c} \text{X} \\ \text{X} \end{array} = \begin{array}{c} \text{X} \\ \text{X} \end{array}. \quad (2.15)$$

Using (2.7) to (2.9) and the first relation from (2.10) but not the second (the braid relation), it follows that

$$\begin{array}{c} \text{X} \\ \text{X} \end{array} - \begin{array}{c} \text{X} \\ \text{X} \end{array} = \begin{array}{c} \text{X} \\ \text{X} \end{array} - \begin{array}{c} \text{X} \\ \text{X} \end{array}, \quad \begin{array}{c} \text{X} \\ \text{X} \end{array} - \begin{array}{c} \text{X} \\ \text{X} \end{array} = \begin{array}{c} \text{X} \\ \text{X} \end{array} - \begin{array}{c} \text{X} \\ \text{X} \end{array}, \quad \begin{array}{c} \text{X} \\ \text{X} \end{array} - \begin{array}{c} \text{X} \\ \text{X} \end{array} = \begin{array}{c} \text{X} \\ \text{X} \end{array} - \begin{array}{c} \text{X} \\ \text{X} \end{array}. \quad (2.16)$$

Proof. In view of the symmetry \mathbb{R} , it suffices to check the first relation from (2.15) and the first two relations from (2.16). They are all quite easy. For example, here are the details for the second relation from (2.16). We may use (2.11) and (2.12) since they are consequences of (2.7) to (2.9). We have that

$$\begin{array}{c} \text{X} \\ \text{X} \end{array} = \begin{array}{c} \text{X} \\ \text{X} \end{array} - \begin{array}{c} \text{X} \\ \text{X} \end{array} = \begin{array}{c} \text{X} \\ \text{X} \end{array} - \begin{array}{c} \text{X} \\ \text{X} \end{array} + \begin{array}{c} \text{X} \\ \text{X} \end{array},$$

$$\begin{array}{c} \text{X} \\ \text{X} \end{array} = \begin{array}{c} \text{X} \\ \text{X} \end{array} + \begin{array}{c} \text{X} \\ \text{X} \end{array} = \begin{array}{c} \text{X} \\ \text{X} \end{array} + \begin{array}{c} \text{X} \\ \text{X} \end{array} - \begin{array}{c} \text{X} \\ \text{X} \end{array}.$$

Subtracting gives the result. In this calculation we have omitted several diagrams which are 0 due to the first relation from (2.10). \square

Next we exploit the isomorphism $\Lambda \cong \Gamma \otimes \Lambda^{[2]}$ from Lemma 2.1 and freeness of $\Gamma = \mathbb{k}[q_1, q_3, \dots]$ to see that there is a well-defined $\Lambda^{[2]}$ -algebra homomorphism

$$\zeta : \Lambda \rightarrow \text{End}_{\mathbf{NB}_t}(\mathbb{1}) \quad (2.17)$$

mapping $q_{2r-1} \in \Gamma$ to $-2(-1)^t \bigcirc_{2r-1}$ for each $r \geq 1$. By [BWW23, Cor. 2.5], it follows that ζ maps q_r to $-2(-1)^t \bigcirc_r$ for every $r \geq 1$. However, we warn the reader that $q_0 = 1$ does not agree with $-2(-1)^t \bigcirc$.

We denote the image of any $a \in \Lambda$ under ζ simply by the labelled bubble $@$. The morphism space $\text{Hom}_{\mathbf{NB}_t}(B^{*n}, B^{*m})$ is naturally a (Λ, Λ) -bimodule so that the left and right actions $a \in \Lambda$ are by horizontal composition on the left or right with $@$, respectively. The left and right actions by elements $\Lambda^{[2]}$ coincide since we are considering a $\Lambda^{[2]}$ -linear monoidal category, but this is seldom the case for elements of Γ . Note also that the symmetries \mathbb{R} and \mathbb{T} from (2.13) and (2.14) fix all of the bubbles $@$ for any $a \in \Lambda$.

Theorem 2.5. *For $m, n \geq 0$, the morphism space $\text{Hom}_{\mathbf{NB}_t}(B^{*n}, B^{*m})$ is free as a right Λ -module with basis given by a certain combinatorially-defined set $\mathbb{D}(m, n)$ of dotted reduced $m \times n$ string diagrams.*

Proof. In [BWW24, Th. 5.3], it is proved that morphism spaces in the nil-Brauer category considered there are free as right Γ -modules with basis $D(m, n)$. In view of Lemma 2.1, the result here follows since our nil-Brauer category is the one from [BWW24] base-changed from \mathbb{k} to $\Lambda^{[2]}$. \square

2.3. Identification of the Grothendieck ring of \mathbf{NB}_t . A thick string $\mathbf{1}_n$ labelled by n indicates n parallel thin strings, i.e., it is the identity endomorphism of B^{*n} . The crossing of two thick strings denotes the minimal length composition of crossings of thin strings:

$$\begin{array}{c} \diagup \\ \diagdown \\ a \quad b \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ a \quad b \end{array}.$$

We use a cross on a string of thickness n to indicate the composition of thin crossings according to a reduced expression for the longest permutation in S_n . On a string of thickness one, this is just the identity. In general, by the braid relation, we have that

$$\begin{array}{c} \downarrow \\ \downarrow \\ a+b \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ a \quad b \end{array}$$

for any $a + b = n$. Given $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$, let

$$\begin{array}{c} \downarrow \\ \downarrow \\ n \end{array} \alpha := \begin{array}{c} \downarrow \\ \downarrow \\ \alpha_1 \end{array} \begin{array}{c} \downarrow \\ \downarrow \\ \alpha_2 \end{array} \cdots \begin{array}{c} \downarrow \\ \downarrow \\ \alpha_n \end{array}$$

Using this notation, we define

$$\mathbf{e}_n := \begin{array}{c} \downarrow \\ \downarrow \\ n \end{array} \rho_n \in \text{End}_{\mathbf{NB}_t}(B^{*n}), \quad (2.18)$$

where $\rho_n := (n-1, \dots, 2, 1, 0) \in \mathbb{N}^n$. For example:

$$\mathbf{e}_0 = \text{id}_{\mathbb{1}}, \quad \mathbf{e}_1 = \begin{array}{c} \downarrow \\ \downarrow \\ 1 \end{array}, \quad \mathbf{e}_2 = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}, \quad \mathbf{e}_3 = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}, \quad \mathbf{e}_4 = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array}.$$

The significance of these endomorphisms is explained by the following result:

Theorem 2.6. *For $n \geq 0$, \mathbf{e}_n is a primitive homogeneous idempotent, and any primitive homogeneous idempotent in \mathbf{NB}_t is equivalent to \mathbf{e}_n for a unique $n \geq 0$.*

Proof. This is proved in [BWW23, Cor. 4.24], but working over the ground field \mathbb{k} rather than the ground ring $\Lambda^{[2]}$ here. This change has no effect on homogeneous idempotents, since they necessarily have degree 0 and $\Lambda^{[2]}$ is a connected positively graded algebra. \square

For a graded category \mathbf{C} , let $\text{gKar}(\mathbf{C})$ be its graded Karoubi envelope. This is obtained by enlarging the category by formally adjoining an invertible grading shift functor q in such a way that

$$\text{Hom}(X, Y)_n = \text{Hom}(X, q^n Y)_0 = \text{Hom}(q^{-n} X, Y)_0,$$

then passing to the usual additive Karoubi envelope. In particular, objects are pairs $[X, \mathbf{e}]$ consisting of a formal finite direct sum X of grading shifts of objects of \mathbf{C} and a matrix \mathbf{e} of endomorphisms defining a homogeneous idempotent $\mathbf{e} : X \rightarrow X$. We write $K_0(\text{gKar}(\mathbf{C}))$ for the split Grothendieck group consisting of degree 0 isomorphism classes of objects in $\text{gKar}(\mathbf{C})$. It is a $\mathbb{Z}[q, q^{-1}]$ -module with action of q induced by the grading shift functor. If \mathbf{C} is monoidal then $\text{gKar}(\mathbf{C})$ is too, so we get induced a multiplication making $K_0(\text{gKar}(\mathbf{C}))$ into a $\mathbb{Z}[q, q^{-1}]$ -algebra.

In [BWW23], it is shown that the $\mathbb{Z}[q, q^{-1}]$ -algebra $K_0(\text{gKar}(\mathbf{NB}_t))$ is isomorphic to a certain $\mathbb{Z}[q, q^{-1}]$ -form U_t^i for the split ι -quantum group of rank one. We refer to [BWW23, Sec. 2] for the

full definition of this, just noting for now that $\mathbb{Q}(q) \otimes_{\mathbb{Z}[q, q^{-1}]} U_t^i$ is the polynomial algebra $\mathbb{Q}(q)[b]$, and the $\mathbb{Z}[q, q^{-1}]$ -form U_t^i is free as a $\mathbb{Z}[q, q^{-1}]$ -module with basis $b^{(n)}$ ($n \geq 0$) defined from

$$b^{(n)} := \frac{b^{\sigma_t(n)}}{[n]_q!} \prod_{\substack{i=0 \\ i \equiv t \pmod{2}}}^{n-1} (b^2 - [i]_q^2) \quad \text{where} \quad \sigma_t(n) := \begin{cases} 0 & \text{if } n \text{ is even} \\ -1 & \text{if } n \text{ is odd and } t = 0 \\ 1 & \text{if } n \text{ is odd and } t = 1. \end{cases} \quad (2.19)$$

The basis $b^{(n)}$ ($n \geq 0$) is the ι -canonical basis of U_t^i in the general sense of [BW18a, BW18b] associated to the parameter t . These elements are also known as ι -divided powers. Instead of the closed formula (2.19), which was worked out originally in [BW18c], $b^{(n)}$ can be also defined recursively: we have that $b^{(0)} := 1$ and

$$b \cdot b^{(n)} = \begin{cases} [n+1]_q b^{(n+1)} + [n]_q b^{(n-1)} & \text{if } n \equiv t \pmod{2} \\ [n+1]_q b^{(n+1)} & \text{if } n \not\equiv t \pmod{2}. \end{cases} \quad (2.20)$$

Let $\text{Par}(r \times c)$ be the set of partitions whose Young diagram fits into the $r \times c$ rectangle, and $\text{Par}_{\neq t}(r \times c)$ be the subset consisting of those partitions whose non-zero parts are $\neq t \pmod{2}$. Another notable formula [BWW23, Cor. 2.13] gives that

$$b^n = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} [n-2i]_q! \left(\sum_{\lambda \in \text{Par}_{\neq t}(i \times (n-2i))} [\lambda_1+1]_q^2 \cdots [\lambda_i+1]_q^2 \right) b^{(n-2i)} \quad (2.21)$$

for any $n \geq 0$.

Theorem 2.7. *There is an isomorphism of $\mathbb{Z}[q, q^{-1}]$ -algebras $K_0(\text{gKar}(\mathbf{NB}_t)) \xrightarrow{\sim} U_t^i$ taking the isomorphism class of*

$$B^{(n)} := q^{\binom{n}{2}} [B^{*n}, \mathbf{e}_n] \in \text{gKar}(\mathbf{NB}_t) \quad (2.22)$$

to the ι -canonical basis vector $b^{(n)}$.

Proof. This follows from [BWW23, Th. B], but a couple of comments are in order. One is that we have extended the ground ring from \mathbb{k} to $\Lambda^{[2]}$, but this causes no problem since $\Lambda^{[2]}$ is a connected positively graded algebra. More likely to cause confusion, in [BWW23], results were not explained in terms of graded Karoubi envelopes, rather, they were formulated in terms of finitely generated graded projective left modules over the path algebra NB of \mathbf{NB}_t . The graded category NB-pgmod of finitely generated projective graded left NB -modules is *contravariantly* equivalent to $\text{gKar}(\mathbf{NB}_t)$ via the Yoneda equivalence. Being contravariant, the canonical isomorphism

$$K_0(\text{gKar}(\mathbf{NB}_t)) \xrightarrow{\sim} K_0(\text{NB-pgmod})$$

induced by the Yoneda equivalence is *anti-linear*. This accounts for the fact that the grading shift in (2.22) is the negation of the grading shift in the definition of the corresponding indecomposable projective module $P(n)$ defined in [BWW23]. \square

From Theorem 2.7 and (2.21), we get that

$$B^{*n} \cong \bigoplus_{i=0}^{\lfloor \frac{n}{2} \rfloor} [n-2i]_q! \left(\sum_{\lambda \in \text{Par}_{\neq t}(i \times (n-2i))} [\lambda_1+1]_q^2 \cdots [\lambda_i+1]_q^2 \right) B^{(n-2i)} \quad (2.23)$$

for any $n \geq 0$. There is a unique top degree summand in this direct sum decomposition equal to $q^{-\binom{n}{2}} B^{(n)}$. This is the ‘‘image’’ of the idempotent \mathbf{e}_n .

2.4. Bubble slides. Given $f(x) = \sum_{r \geq 0} c_r x^r \in \mathbb{k}[x]$ and a dot in some string diagram s , we denote

$$\sum_{r \geq 0} c_r \times (\text{the morphism obtained from } s \text{ by labeling the dot by } r)$$

by attaching what we call a *pin* to the dot, labeling the node at the head of the pin by $f(x)$:

$$\begin{array}{c} \bullet \\ | \\ \boxed{f(x)} \end{array} := \sum_{r \geq 0} c_r \begin{array}{c} | \\ \bullet \end{array} r \in \text{End}_{\mathbf{NB}_t}(B). \quad (2.24)$$

More generally, $f(x)$ here could be a polynomial with coefficients in the algebra $\mathbb{k}((u^{-1}))$ of formal Laurent series in an indeterminate u^{-1} ; then the string s decorated with a pin labelled $f(x)$ defines a morphism in $\mathbf{NB}_t((u^{-1}))$. We think of this as being a generating function for a family of morphisms.

Now we can discuss “bubble slides”. We obviously have that

$$\boxed{e^{[2]}(u)} \Big| = \Big| \boxed{e^{[2]}(u)} \quad (2.25)$$

since \mathbf{NB}_t is a $\Lambda^{[2]}$ -linear monoidal category. The following is [BWW24, Th. 2.5(5)]:

$$\boxed{q(u)} \Big| = \begin{array}{c} | \\ \bullet \\ | \end{array} \boxed{\frac{u-x}{u+x}} \boxed{q(u)}. \quad (2.26)$$

Writing $\sqrt{q(u)}$ for the unique square root of $q(u)$ in $1 + u^{-1}\Gamma[[u^{-1}]]$, (2.26) is equivalent to

$$\boxed{\sqrt{q(u)}} \Big| = \begin{array}{c} | \\ \bullet \\ | \end{array} \boxed{\frac{u-x}{u+x}} \boxed{\sqrt{q(u)}}. \quad (2.27)$$

The following lemma describes the bubble slides for elementary and complete symmetric functions; the formulae are the same as the one for $\sqrt{q(u)}$.

Lemma 2.8. *We have that*

$$\boxed{e(u)} \Big| = \begin{array}{c} | \\ \bullet \\ | \end{array} \boxed{\frac{u-x}{u+x}} \boxed{e(u)}, \quad \boxed{h(u)} \Big| = \begin{array}{c} | \\ \bullet \\ | \end{array} \boxed{\frac{u-x}{u+x}} \boxed{h(u)}. \quad (2.28)$$

Proof. Recall that $e^{[2]}(u) = e(u)e(-u)$ and $q(u) = e(u)e(-u)^{-1}$. Hence, $e(u)^2 = e^{[2]}(u)q(u)$ and $e(u) = \sqrt{e^{[2]}(u)}\sqrt{q(u)}$ (again, for $e^{[2]}(u)$, we take the square root with constant term 1). The first identity in (2.28) is now clear from (2.25) and (2.27). To deduce the second one, replace u by $-u$ and then take inverses on both sides. \square

Corollary 2.9. *We have that* $\boxed{\mathfrak{e}_r} \Big| + \Big| \boxed{\mathfrak{e}_r} = 2 \sum_{s=0}^r \boxed{\mathfrak{e}_{r-s}} \Big|_s = 2 \sum_{s=0}^r (-1)^s \Big|_s \boxed{\mathfrak{e}_{r-s}}$.

Proof. For the first equality, equate coefficients of u^{-r} on both sides of the first relation from (2.28) using the identity $\frac{u-x}{u+x} = 2(1+xu^{-1})^{-1} - 1 = 2 \sum_{s \geq 0} (-1)^s x^s u^{-s} - 1$. The second equality follows from the first by applying R. \square

2.5. The cyclotomic nil-Brauer category. The endomorphism of B appearing in Corollary 2.9 will play an important role, so we introduce some special notation for it: let

$$\boxed{r} := \boxed{\mathfrak{e}_r} \Big| + \Big| \boxed{\mathfrak{e}_r}. \quad (2.29)$$

Definition 2.10. For $l \geq 0$, let \mathbf{I}_l be the four-sided ideal of \mathbf{NB}_t generated by \boxed{l} and $\boxed{\mathfrak{e}_r}$ ($r > l$). By “four-sided ideal” here, we mean that it is a *two-sided ideal* of the Λ -linear category \mathbf{NB}_t , i.e. a family of Λ -submodules $\mathbf{I}_l(B^{*n}, B^{*m}) \subseteq \text{Hom}_{\mathbf{NB}_t}(B^{*n}, B^{*m})$ for all $m, n \geq 0$ closed under vertical composition on top and bottom with any morphism, which is also a *two-sided tensor ideal*, i.e. it is closed under

horizontal composition on left and right by any morphism. The quotient $\mathbf{NB}_l/\mathbf{I}_l$ is again a Λ -linear strict graded monoidal category, which we call the *level l cyclotomic nil-Brauer category*.

Theorem 2.11. *The four-sided ideal \mathbf{I}_l is equal both to the right tensor ideal and to the left tensor ideal generated by \boxed{l} and $\textcircled{e_r}$ ($r > l$).*

Proof. Let \mathbf{J}_l be the right tensor ideal generated by \boxed{l} and $\textcircled{e_r}$ ($r > l$). In a series of claims below, we will show that \mathbf{J}_l is a two-sided tensor ideal. This is sufficient to complete the proof of the theorem. Indeed, since \mathbf{J}_l has the same generators as \mathbf{I}_l , it follows that $\mathbf{I}_l = \mathbf{J}_l$, so \mathbf{I}_l is the right tensor ideal with these generators. Applying \mathbf{R} , it also follows that \mathbf{I}_l is the left tensor ideal with these generators.

Claim 1: *We have that $\overbrace{\boxed{l}}^{\curvearrowright} \in \mathbf{J}_l(B \star B, \mathbb{1})$ and $\underbrace{\boxed{l}}_{\curvearrowleft} \in \mathbf{J}_l(\mathbb{1}, B \star B)$.* The first assertion follows because $\overbrace{\boxed{l}}^{\curvearrowright} = \overbrace{\boxed{l}}^{\curvearrowright}$, as is clear from the definition (2.29). The second one is similar (or one can apply \mathbf{T}).

Claim 2: *We have that $\boxed{r} \in \mathbf{J}_l(B, B)$ for all $r \geq l$.* This is proved by induction on r , the base case $r = l$ being immediate from the definition of \mathbf{J}_l . Now suppose $r > l$. By Corollary 2.9 we have that

$$\boxed{r} = 2 \sum_{s=0}^r \textcircled{e_{r-s}} \downarrow^s = 2 \textcircled{e_r} \downarrow + \sum_{s=1}^r \textcircled{e_{r-s}} \downarrow^{s-1} = 2 \textcircled{e_r} \downarrow + \sum_{s=0}^{r-1} \textcircled{e_{r-1-s}} \downarrow^s = 2 \textcircled{e_r} \downarrow + \boxed{r-1}. \quad (2.30)$$

This lies in \mathbf{J}_l by induction (and because $\textcircled{e_r} \in \mathbf{J}_l$).

Claim 3: *We have that $\textcircled{e_r} \in \mathbf{J}_l(B, B)$ for any $r > l$.* This follows from Claim 2 because (by (2.29))

$$\textcircled{e_r} = \boxed{r} - \textcircled{e_r}.$$

Claim 4: *We have that $\overbrace{\boxed{r}}^{\curvearrowright} \in \mathbf{J}_l(B \star B, B \star B)$ for all $r \geq l$.* By (2.29), we have that

$$\overbrace{\boxed{r}}^{\curvearrowright} = \overbrace{\boxed{r}}^{\curvearrowright} + \overbrace{\textcircled{e_r}}^{\curvearrowright} - \overbrace{\textcircled{e_r}}^{\curvearrowright}.$$

The latter two terms on the right-hand side are zero by (2.10), thus the claim follows.

Claim 5: *We have that $\overbrace{\boxed{l}}^{\times} \in \mathbf{J}_l(B \star B, B \star B)$.* By the second relation from (2.12), we have that

$$\overbrace{\boxed{l}}^{\times} = \overbrace{\boxed{l}}^{\curvearrowright} - \overbrace{\boxed{l}}^{\curvearrowleft} + \overbrace{\boxed{l}}^{\curvearrowright}.$$

The first and third terms on the right hand side lie in $\mathbf{J}_l(B \star B, B \star B)$ by Claims 1 and 4. It remains to show that the second term on the right hand side belongs to $\mathbf{J}_l(B \star B, B \star B)$ too. We have

$$\overbrace{\boxed{l}}^{\curvearrowleft} = \overbrace{\boxed{l+1}}^{\curvearrowleft} - 2 \textcircled{e_{l+1}}$$

by (2.30). Both of the terms on the right hand side here lie in $\mathbf{J}_l(B \star B, B \star B)$ by Claims 4 and 3.

interpreting $f^{(-1)}\eta_l$ as 0. We find it helpful to visualize this as follows:

$$\begin{array}{ccc}
 & f^{(0)}\eta_l & \\
 & [1]_q \left(\begin{array}{c} \nearrow \\ \downarrow \end{array} \right) q^{1-l} [l]_q & \uparrow \\
 & f^{(1)}\eta_l & \\
 & [2]_q \left(\begin{array}{c} \nearrow \\ \downarrow \end{array} \right) q^{3-l} [\ell-1]_q & \\
 & \vdots & \\
 & [\ell-1]_q \left(\begin{array}{c} \nearrow \\ \downarrow \end{array} \right) q^{l-3} [2]_q & \\
 & f^{(l-1)}\eta_l & \\
 & [\ell]_q \left(\begin{array}{c} \nearrow \\ \downarrow \end{array} \right) q^{l-1} [1]_q & \\
 & f^{(l)}\eta_l & \\
 f \downarrow & & \uparrow \rho(f)
 \end{array} \tag{2.32}$$

Recall finally that there is a unique symmetric bilinear form $(\cdot, \cdot)_l : \mathbb{V}(l) \times \mathbb{V}(l)$ such that $(\eta_l, \eta_l)_l = 1$ and $(uv_1, v_2)_l = (v_1, \rho(u)v_2)_l$ for all $v_1, v_2 \in \mathbb{V}(l)$ and $u \in \mathbb{U}$. The standard basis is orthogonal with

$$(f^{(n)}\eta_l, f^{(n)}\eta_l)_l = q^{-n(l-n)} \begin{bmatrix} l \\ n \end{bmatrix}_q \in 1 + \mathbb{N}[q^{-1}], \tag{2.33}$$

which is the Poincaré polynomial for the cohomology of the Grassmannian $\text{Gr}_{n,l}$.

The ι -quantum group \mathbb{U}'_l is naturally a $\mathbb{Z}[q, q^{-1}]$ -subalgebra of \mathbb{U} , with $b \in \mathbb{U}'_l$ being the element $f + \rho(f) \in \mathbb{U}$. Of course, the \mathbb{U} -module $\mathbb{V}(l)$ can also be viewed as a \mathbb{U}'_l -module by restriction. By (2.31), we have that $b\eta_l = f\eta_l$ and

$$b \cdot f^{(n)}\eta_l = [n+1]_q f^{(n+1)}\eta_l + q^{2n-l-1} [l-n+1]_q f^{(n-1)}\eta_l \tag{2.34}$$

for $n \geq 1$. The ι -canonical basis of \mathbb{U}'_l descends to the ι -canonical basis $b^{(n)}\eta_l$ ($0 \leq n \leq l$) of $\mathbb{V}(l)$, and $b^{(n)}\eta_l = 0$ for $n > l$. These statements all follow from [BW18a, BW18b].

Remark 2.13. Let I_l be the kernel of the \mathbb{U}'_l -module homomorphism $\mathbb{U}'_l \rightarrow \mathbb{V}(l), u \mapsto u\eta_l$. From the preceding discussion, it follows that I_l is generated by the ι -canonical basis elements $b^{(n)}$ for $n > l$. In fact, by (2.20), this ideal is generated already just by $b^{(l+1)}$. The quotient \mathbb{U}'_l/I_l is isomorphic to $\mathbb{V}(l)$ as a \mathbb{U}'_l -module, but it has the additional structure of a $\mathbb{Z}[q, q^{-1}]$ -algebra.

The ι -canonical basis of $\mathbb{V}(l)$ has an alternative characterization; in fact, this is the way the ι -canonical basis gets introduced in the first place in [BW18a, BW18b]. To explain this, we need the ι -bar involution $\mathbb{V}(l) \rightarrow \mathbb{V}(l), v \mapsto \bar{v}$. This is the unique anti-linear map fixing η_l such that $\overline{b\bar{v}} = b\bar{v}$ for all $v \in \mathbb{V}(l)$. This is *not* the usual bar involution on $\mathbb{V}(l)$ as it is definitely not true that $\overline{f\bar{v}} = f\bar{v}$. Then, for $0 \leq n \leq l$, $b^{(n)}\eta_l \in \mathbb{V}(l)$ is the unique bar-invariant vector such that

$$b^{(n)}\eta_l = f^{(n)}\eta_l + (a q^{-1} \mathbb{Z}[q^{-1}]\text{-linear combination of the vectors } f^{(i)}\eta_l \text{ for } i < n). \tag{2.35}$$

There is also an explicit formula expressing the ι -canonical basis in terms of the standard bases:

Theorem 2.14 ([BW18c, (2.16),(2.17),(3.8),(3.9)]). *We have that*

$$b^{(n)}\eta_l = \begin{cases} \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} q^{-i(l+2i-n-1)} \begin{bmatrix} (l+2i-n)/2 \\ i \end{bmatrix}_{q^2} f^{(n-2i)}\eta_l & \text{if } 0 \leq n \leq l \text{ and } n \equiv t \pmod{2}, \\ \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} q^{-i(l+2i-n)} \begin{bmatrix} (l+2i-n-1)/2 \\ i \end{bmatrix}_{q^2} f^{(n-2i)}\eta_l & \text{if } 0 \leq n \leq l \text{ and } n \not\equiv t \pmod{2} \\ 0 & \text{if } n > l. \end{cases}$$

Proof. This follows from the four formulae cited from [BW18c], but it is some work to convert these into the more concise form recorded here. Alternatively, one can give an independent proof by using (2.31) to verify that our reformulation satisfies the recurrence relation (2.20). \square

Corollary 2.15. *For $0 \leq n \leq \lfloor \frac{l}{2} \rfloor$, we have that $(\eta_l, b^{(2n)}\eta_l)_l = q^{-n(l+t-1)} \begin{bmatrix} (l-t)/2 \\ n \end{bmatrix}_{q^2}$.*

Here is one more observation about the bilinear form on $V(l)$.

Lemma 2.16. *The top degree term of the Laurent polynomial $(b^n\eta_l, b^n\eta_l)_l$ is $q^{2\binom{n}{2}}$ for $0 \leq n \leq l$, and it is $q^{(2n-l)(l-1)}$ for $n \geq l$.*

Proof. Since $(b^n\eta_l, b^n\eta_l)_l = (\eta_l, b^{2n}\eta_l)_l$, we need to compute the η_l -coefficient of

$$b^{2n}\eta_l = (f + \rho(f))^{2n}\eta_l.$$

Such coefficients arise from the monomials in the expansion of $(f + \rho(f))^{2n}$ that have f appearing n times and $\rho(f)$ appearing n times. The coefficient of η_l arising when each of these monomials is applied to η_l is easily computed by contemplating (2.32). The Laurent polynomials on the edges of this diagram are monic of degrees $0, 1, \dots, l-1$ from top to bottom. It follows that the biggest degree arises from a unique monomial, namely, $f^n \rho(f)^n$ if $n \leq l$ or $f^l (\rho(f)f)^{n-l} \rho(f)^l$ if $n \geq l$. Since the Laurent polynomials are monic, this produces the top degree term in the statement of the lemma. \square

2.7. The special cases $l = 0$ and $l = 1$. The definition of \mathbf{cNB}_l makes sense when $l = 0$ and $l = 1$, but these are degenerate cases and we treat them separately here. In the case $l = 0$, \mathbf{cNB}_0 is the trivial graded monoidal category with one object $\mathbb{1}$ and $\text{End}_{\mathbf{cNB}_0}(\mathbb{1}) = \mathbb{k}$. Its graded Karoubi envelope is monoidally equivalent to $\mathbf{gVec}_{\text{fd}}$, with $K_0(\mathbf{gKar}(\mathbf{cNB}_0)) \cong V(0)$ so that $[\mathbb{1}]$ is identified with η_0 . There is nothing more to be said about this case.

If $l = 1$ then \mathbf{cNB}_l is slightly more interesting, but still easy to understand directly. It can be presented as a strict graded \mathbb{k} -linear monoidal category with one generating object B and three generating morphisms

$$\begin{array}{ccc} \textcircled{e_1} : \mathbb{1} \rightarrow \mathbb{1}, & \cap : B \star B \rightarrow \mathbb{1}, & \cup : \mathbb{1} \rightarrow B \star B, \\ \text{(degree 2)} & \text{(degree 0)} & \text{(degree 0)} \end{array}$$

subject to the following relations:

$$\bigcirc = \text{id}_{\mathbb{1}}, \quad \left| \textcircled{e_1} = -\textcircled{e_1} \right|, \quad \left| \cup = \cup \right|, \quad \left| \cap = \cap \right|, \quad \left| \left| = \bigcup \right. \right|. \quad (2.36)$$

This follows quite easily from the defining relations above; the dot is redundant as a generator since $\uparrow = \left| \textcircled{e_1} \right|$ in \mathbf{NB}_1 by Corollary 2.9, and the crossing can be seen to be 0 using the second relation from (2.8).

Theorem 2.17. *View the morphism space $\mathrm{Hom}_{\mathbf{cNB}_1}(B^{*n}, B^{*m})$ as a right $\mathbb{k}[x]$ -module so that x acts by horizontal composition on the right by $\textcircled{1}$. Then this morphism space is $\{0\}$ unless $m \equiv n \pmod{2}$, in which case it is free of rank one as a right $\mathbb{k}[x]$ -module with basis given by the string diagram obtained by drawing $\lfloor m/2 \rfloor$ side-by-side caps at the top and $\lfloor n/2 \rfloor$ side-by-side cups at the bottom, plus a vertical string on the right hand side if m and n are odd.*

Proof. It is an exercise using the relations to see that $\mathrm{Hom}_{\mathbf{cNB}_1}(B^{*n}, B^{*m})$ is $\{0\}$ if $m \not\equiv n \pmod{2}$, and that it is spanned as a $\mathbb{Z}[x, x^{-1}]$ -module by the morphism just described otherwise. To prove the freeness, we let \mathbf{C} be the \mathbb{k} -linear graded category with objects \mathbb{N} and $\mathrm{Hom}_{\mathbf{C}}(n, m) := \mathbb{k}[x]$ if $m \equiv n \pmod{2}$ or $\{0\}$ otherwise, with the obvious composition law defined by multiplication of polynomials. We make this into a \mathbb{k} -linear graded monoidal category by defining $m \star n := m + n$, and defining the tensor product of morphisms $f(x) : m \rightarrow m', g(x) : n \rightarrow n'$ by setting $f(x) \star g(x) := f((-1)^n x)g(x) : m+n \rightarrow m'+n'$. It is trivial to check from the defining relations that there is a \mathbb{k} -linear graded monoidal functor $\mathbf{cNB}_1 \rightarrow \mathbf{C}$ taking $B \mapsto 1$, $\textcircled{1} \mapsto (x : 0 \rightarrow 0)$, and the cap and the cup to the morphisms $2 \rightarrow 0$ and $0 \rightarrow 2$ represented by $1 \in \mathbb{k}[x]$. The desired freeness is now obvious. \square

From this, it follows that $B^{*n} \cong \mathbb{1}$ if n is even, $B^{*n} \cong B$ if n is odd, and $\mathrm{End}_{\mathbf{cNB}_1}(\mathbb{1}) \cong \mathrm{End}_{\mathbf{cNB}_1}(B) \cong \mathbb{k}[x]$. Hence, there is a $\mathbb{Z}[q, q^{-1}]$ -module isomorphism

$$K_0(\mathrm{gKar}(\mathbf{cNB}_1)) \xrightarrow{\sim} \mathbf{V}(1), \quad [\mathbb{1}] \mapsto \eta_1, \quad [B] \mapsto b\eta_1. \quad (2.37)$$

Since $B \star B \cong \mathbb{1}$, we have that $[B]^2 = [\mathbb{1}]$, so as an algebra $K_0(\mathrm{gKar}(\mathbf{cNB}_1)) \cong \mathbb{Z}[q, q^{-1}][b]/(b^2 - 1)$.

3. REMINDERS ABOUT SINGULAR SOERGEL BIMODULES

In this expository section, we review some fundamental results about singular Soergel bimodules and the graphical calculus for them following [Wil11, ESW17]. Our general setup is essentially the same as in [Wil11, §3.1] with additional assumptions imposed so that we can appeal to the diagrammatic calculus of [ESW17]; see also [EMTW20, Ch. 24]. Later, we will appeal to these results only for the reflection realization in finite type D, for which it is well known that all of the assumptions hold.

3.1. Coxeter group realization. Let (W, S) be a Coxeter system with length function $\ell : W \rightarrow \mathbb{N}$ and Bruhat order \leq . We assume simple reflections are parametrized by another set¹ N (“nodes of the Coxeter diagram”) via a given bijection $N \rightarrow S, i \mapsto s_i$. In fact, excepting one use in a bullet point below, we no longer use the set S , to avoid conflict with the q -Schur algebra S to be defined later.

We fix a realization in the sense of [EW16, Def. 3.1]. So we are given a finite-dimensional vector space \mathfrak{h} over the ground field \mathbb{k} ($\mathrm{char} \mathbb{k} \neq 2$) together with subsets $\{\alpha_i \mid i \in N\} \subset \mathfrak{h}^*$ and $\{\alpha_i^\vee \mid i \in N\} \subset \mathfrak{h}$ such that

- $\langle \alpha_i, \alpha_i^\vee \rangle = 2$ for all $i \in N$;
- the function $S \rightarrow GL(\mathfrak{h}^*)$ taking s_i to the linear map $\lambda \mapsto \lambda - \langle \lambda, \alpha_i^\vee \rangle \alpha_i$ extends to a representation of W .

We assume in addition that the realization is

- *balanced* in the sense of [EW16, Def. 3.6];
- *finitary Demazure surjective* as in [EMTW20, Sec. 24.3.6] (called *generalized Demazure surjective* in [EKLP23, Def. 3.5]), meaning that the longest Demazure operator $\partial_I : R \rightarrow R^I$ is surjective for each $I \subseteq_f N$ (the notation here will be explained shortly);

¹This is the second author’s pickiness. It is perfectly reasonable to index everything by the set S but that leads to expressions like α_{s_i} not α_i to denote the simple root associated to $s_i \in S$.

- *faithful* and *reflection faithful* as in [Wil11, Sec. 4.1]. This means for any $w \in W$ that the fixed point space $\{v \in \mathfrak{h}^* \mid w(v) = v\}$ is \mathfrak{h}^* if and only if $w = 1$, and it is of codimension 1 if and only if w is conjugate to a simple reflection.

Using further language yet to be introduced, these assumptions are required to ensure

- the Demazure operators $\partial_i (i \in N)$ satisfy the braid relations, and the sets Φ_I^+ of “positive roots” introduced below are of the claimed size;
- the upgraded Chevalley theorem as formulated in [EMTW20, Th. 24.36, Th. 24.40] (and proved again in [EKLP23, Sec. 4]) holds, i.e., we have available the appropriate squares of Frobenius extensions (see Subsections 3.2 and 3.3);
- the hypotheses of [Wil11] are satisfied (see Subsection 3.4 below where the important results deduced from these hypotheses discussed further).

For further justification of the assumptions, see [EMTW20, Sec. 24.3.6] and [EKLP23, Sec. 3].

Let $R := \mathbb{k}[\mathfrak{h}]$ be the symmetric algebra on the dual space \mathfrak{h}^* , viewed as a graded algebra with \mathfrak{h}^* in degree 2. The Coxeter group acts naturally on R by graded algebra automorphisms. For $i \in N$, the Demazure operator $\partial_i : R \rightarrow R$ is the degree -2 linear map defined by

$$\partial_i(f) := \frac{f - s_i(f)}{\alpha_i}.$$

The endomorphisms $\partial_i (i \in N)$ generate a copy of the *nil-Coxeter algebra* associated to the Coxeter group W . They satisfy the same braid relations as the simple reflections in the Coxeter group, but the quadratic relation $s_i^2 = 1$ is replaced by $\partial_i^2 = 0$.

For $I \subseteq N$, let W_I be the parabolic subgroup $\langle s_i \mid i \in I \rangle$ of W , and R^I be the subalgebra of R consisting of its fixed points. We say that I is *finitary* if W_I is a finite group, and use the notation $I \subseteq_f N$ to indicate that I is such a subset. The letters I, J, K will be reserved for finitary subsets. For $I \subseteq_f N$, let w_I be the longest element of W_I , and ∂_I be the product of Demazure operators corresponding to a reduced expression for w_I . Let

$$\tilde{\pi}_I := \sum_{w \in W_I} q^{-2\ell(w)}, \quad \pi_I := \sum_{w \in W_I} q^{\ell(w_I) - 2\ell(w)}. \quad (3.1)$$

The first of these is the usual Poincaré polynomial of W_I , and the second is its bar-invariant renormalization, which is a Laurent polynomial of degree $\ell(w_I)$. Let Φ_I be the finite set $\{w(\alpha_i) \mid w \in W_I, i \in I\}$, and

$$\Phi_I^+ := \{w(\alpha_i) \mid w \in W_I, i \in I \text{ such that } \ell(ws_i) = \ell(w) + 1\}. \quad (3.2)$$

The assumption that the realization is balanced implies² that $|\Phi_I^+| = \ell(w_I)$. Finally, let

$$\mu_I := \prod_{\alpha \in \Phi_I^+} \alpha \in R. \quad (3.3)$$

This polynomial is W_I -anti-invariant (meaning $s_i \mu_I = -\mu_I$ for $i \in I$) and has degree $2\ell(w_I)$.

3.2. Chains of Frobenius extensions. Suppose that we are given $J \subseteq I \subseteq_f N$. We have that $W_J \subseteq W_I$, hence, R^I is a subalgebra of R^J . We denote the set of minimal length W_I/W_J -coset representatives by $(W_I/W_J)_{\min}$. Let

$$\eta_I^J := \prod_{\substack{J \subseteq K \subseteq I \\ |K| \equiv |I| \pmod{2}}} \mu_K \bigg/ \prod_{\substack{J \subseteq K \subseteq I \\ |K| \not\equiv |I| \pmod{2}}} \mu_K. \quad (3.4)$$

²This is proven for dihedral groups in [Eli16, Appendix]. The general case is not in the literature to our knowledge. The result will be obvious in our applications, since we make use of realizations associated to typical root data.

The inclusion-exclusion principle implies that this is a polynomial in R (rather than a rational function). In fact, since each μ_K is W_J -anti-invariant, η_I^J lies in R^J . For example:

- writing Ii for $I \cup \{i\}$ for $i \in N - I$ and assuming this set is finitary, we have that $\eta_{Ii}^I = \frac{\mu_{Ii}}{\mu_I} \in R^I$;
- writing Iij for $I \cup \{i, j\}$ for distinct $i, j \in N - I$ and assuming it is finitary, we have that $\eta_{Iij}^I = \frac{\mu_{Iij}}{\mu_i \mu_j} \in R^I$;
- writing $Iijk$ for $I \cup \{i, j, k\}$ for distinct $i, j, k \in N - I$ and assuming it is finitary, we have that $\eta_{Iijk}^I = \frac{\mu_{Iij} \mu_{Iik} \mu_{Ijk} \mu_I}{\mu_{ijk} \mu_i \mu_j \mu_k} \in R^I$.

Continuing with $J \subseteq I \subseteq_f N$, a classical result of Demazure [Dem73] (see also [EKLP23, Th. 4.3]) shows that R^J is a symmetric graded Frobenius extension of R^I . The (unique up to a scalar) Frobenius trace is the (R^I, R^I) -bimodule homomorphism of degree $2(\ell(w_J) - \ell(w_I))$

$$\mathrm{tr}_I^J : R^J \rightarrow R^I, \quad f \mapsto \hat{\partial}_I^J(f), \quad (3.5)$$

where $\hat{\partial}_I^J$ denotes the element of the nil-Coxeter algebra defined by composing Demazure operators in the same order as a reduced expression for the longest element³ $w_I w_J^{-1}$ of $(W_I/W_J)_{\min}$. The comultiplication is the (R^J, R^J) -bimodule homomorphism of degree $2(\ell(w_I) - \ell(w_J))$

$$\Delta_I^J : R^J \rightarrow R^J \otimes_{R^I} R^J, \quad 1 \mapsto \sum_{b \in \mathbb{B}_I^J} b \otimes b^\vee, \quad (3.6)$$

where \mathbb{B}_I^J is a homogeneous basis for R^J as a free R^I -module and b^\vee is the dual basis element defined by $\mathrm{tr}_I^J(ab^\vee) = \delta_{a,b}$ ($a \in \mathbb{B}_I^J$).

The following observation about chains of Frobenius extensions appears in [ESW17, Sec. 2.2].

Lemma 3.1. *Let $K \subseteq J \subseteq I \subseteq_f N$. Then $\mathrm{tr}_I^K = \mathrm{tr}_I^J \circ \mathrm{tr}_J^K$. Moreover, $\{ab \mid a \in \mathbb{B}_J^K, b \in \mathbb{B}_I^J\}$ is a basis for R^K as a free graded R^I -module, with dual basis $\{a^\vee b^\vee \mid a \in \mathbb{B}_J^K, b \in \mathbb{B}_I^J\}$.*

Proof. The equality of Demazure operators $\hat{\partial}_I^K = \hat{\partial}_I^J \circ \hat{\partial}_J^K$ implies that $\mathrm{tr}_I^K = \mathrm{tr}_I^J \circ \mathrm{tr}_J^K$. Using the fact that tr_J^K is R^J -linear, we compute

$$\mathrm{tr}_I^K((ab) \cdot (a')^\vee (b')^\vee) = \mathrm{tr}_I^J \circ \mathrm{tr}_J^K(ab(a')^\vee (b')^\vee) = \mathrm{tr}_I^J(b(b')^\vee \mathrm{tr}_J^K(a(a')^\vee)) = \delta_{a,a'} \delta_{b,b'}.$$

Since \mathbb{B}_J^K spans R^K over R^J and \mathbb{B}_I^J spans R^J over R^I , it follows that $\{ab \mid a \in \mathbb{B}_J^K, b \in \mathbb{B}_I^J\}$ spans R^K over R^I . Linear independence is a consequence of the existence of a dual basis. \square

Using Lemma 3.1, the following useful identities are established in [ESW17, (2.7)]: for $K \subseteq J \subseteq I \subseteq_f N$ and $f \in R^K$, we have that

$$\sum_{b \in \mathbb{B}_I^K} b \otimes \mathrm{tr}_J^K(fb^\vee) = \sum_{a \in \mathbb{B}_I^J} af \otimes a^\vee \in R^K \otimes_{R^I} R^J, \quad (3.7)$$

$$\sum_{b \in \mathbb{B}_I^K} \mathrm{tr}_J^K(bf) \otimes b^\vee = \sum_{a \in \mathbb{B}_I^J} a \otimes fa^\vee \in R^J \otimes_{R^I} R^K. \quad (3.8)$$

In particular, taking $K = J$, this gives that

$$\sum_{b \in \mathbb{B}_I^J} bf \otimes b^\vee = \sum_{b \in \mathbb{B}_I^J} b \otimes fb^\vee \in R^J \otimes_{R^I} R^J \quad (3.9)$$

for all $f \in R^J$. Also [ESW17, (2.8)] gives

$$\sum_{b \in \mathbb{B}_I^K} b \mathrm{tr}_J^K(b^\vee) = \mu_I / \mu_J \in R^I. \quad (3.10)$$

³Although $w_J^{-1} = w_J$, we write $w_I w_J^{-1}$ to remind the reader that $\ell(w_I w_J^{-1}) = \ell(w_I) - \ell(w_J)$.

Taking $K = J$, this implies that $\sum_{b \in \mathbb{B}_I^J} bb^\vee = \mu_I/\mu_J$. Another special case recovers the counit axiom

$$\sum_{b \in \mathbb{B}_I^J} b \operatorname{tr}_I^J(b^\vee) = 1. \quad (3.11)$$

3.3. Diagrammatics for singular Bott-Samelson bimodules. Suppose that $J \subseteq I \subseteq_f N$, so that $R^I \subseteq R^J$. Working with the usual graded categories $R^I\text{-gmod}$ and $R^J\text{-gmod}$ of graded left modules, we have the restriction and induction functors

$$\operatorname{Res}_I^J : R^J\text{-gmod} \rightarrow R^I\text{-gmod}, \quad \operatorname{Ind}_I^J : R^I\text{-gmod} \rightarrow R^J\text{-gmod},$$

which are defined by tensoring on the left with R^J viewed as an (R^I, R^J) -bimodule or as an (R^J, R^I) -bimodule, respectively. They form an adjoint pair $(\operatorname{Ind}_I^J, \operatorname{Res}_I^J)$ via the canonical (degree 0) adjunction between tensor and hom. Since $R^I \subseteq R^J$ is a graded Frobenius extension, there is also an adjunction the other way around, after a suitable grading shift. To make the grading shifts on the various units and counits of adjunction more balanced, we incorporate a grading shift into *restriction*⁴, working henceforth with the functors $q^{\ell(w_I) - \ell(w_J)} \operatorname{Res}_I^J$ and Ind_I^J . We have that

$$q^{\ell(w_I) - \ell(w_J)} \operatorname{Res}_I^J = B_{I,J} \otimes_{R^I} -, \quad \operatorname{Ind}_I^J = B_{J,I} \otimes_{R^I} -, \quad (3.12)$$

where $B_{I,J}$ denotes the graded (R^I, R^J) -bimodule $q^{\ell(w_I) - \ell(w_J)} R^J$, and $B_{J,I}$ denotes the graded (R^J, R^I) -bimodule R^I . The two adjunctions give natural degree-preserving isomorphisms

$$\operatorname{Hom}_{R^J} (B_{J,I} \otimes_{R^I} M, M') \cong q^{\ell(w_J) - \ell(w_I)} \operatorname{Hom}_{R^I} (M, B_{I,J} \otimes_{R^I} M'), \quad (3.13)$$

$$\operatorname{Hom}_{R^I} (B_{I,J} \otimes_{R^J} M', M) \cong q^{\ell(w_I) - \ell(w_J)} \operatorname{Hom}_{R^J} (M', B_{J,I} \otimes_{R^I} M) \quad (3.14)$$

for any graded left R^I -module M and R^J -module M' .

More generally, for $I, J \subseteq_f N$ and $K \subseteq I \cap J$, we define the graded (R^I, R^J) -bimodule

$$B_{I,J}^K := q^{\ell(w_I) - \ell(w_K)} R^K. \quad (3.15)$$

This is naturally isomorphic to $B_{I,K} \otimes_{R^K} B_{K,J}$. We denote $B_{I,J}^{I \cap J}$ simply by $B_{I,J}$. When $I \subseteq J$ or $I \supseteq J$, this notation is consistent with the notation for induction/restriction bimodules that appeared in (3.12).

Remark 3.2. The general results explained in the next subsection (specifically, Corollary 3.8 in the light of Theorems 3.10 to 3.12) imply that $B_{I,J}$ is an indecomposable bimodule for any $I, J \subseteq_f N$. Moreover, for $K \subseteq I \cap J$, we have that $B_{I,J}^K \cong \frac{\pi_{I \cap J}}{\pi_K} B_{I,J}$ as a graded (R^I, R^J) -bimodule.

Definition 3.3 ([Wil11]). The category $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im}$ of *singular Bott-Samelson bimodules* is the full⁵ sub-bicategory of $\mathfrak{g}\mathfrak{B}\mathfrak{im}$ with objects R^I for $I \subseteq_f N$ and 1-morphisms given by the bimodules

$$B_{I_0, I_1}^{J_1} \otimes_{R^{I_1}} B_{I_1, I_2}^{J_2} \otimes_{R^{I_2}} \cdots \otimes_{R^{I_{n-1}}} B_{I_{n-1}, I_n}^{J_n}, \quad (3.16)$$

for all finitary $I_0 \supseteq J_1 \subseteq I_1 \supseteq \cdots \subseteq I_{n-1} \supseteq J_n \subseteq I_n$. We will denote the object R^I of $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im}$ simply by I , so that the object set of $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im}$ is identified with the set of finitary subsets of N . Then, we define the graded bicategory of *singular Soergel bimodules* to be the graded Karoubi envelope $\mathfrak{g}\mathfrak{Kar}(\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im})$. We identify this with a sub-bicategory of $\mathfrak{g}\mathfrak{B}\mathfrak{im}$ in the natural way, so that singular Soergel bimodules are summands of finite direct sums of grading shifts of singular Bott-Samelson bimodules.

⁴This convention is consistent with the choice of duality made in (3.45) below; see also Remark 3.9.

⁵We mean that it has the same 2-morphism spaces.

Now we introduce the string calculus for (the strictification of) $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}$ following [ESW17]. By transitivity of induction and restriction, to generate $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}$, it suffices to consider the bimodules $B_{I,i}$ and $B_{i,I}$ for $I \subset N$ and $i \in N - I$ such that Ii is finitary. We use the string diagrams $I \uparrow i$ and $i \downarrow I$ to denote the identity endomorphisms of these bimodules. For extra clarity, the strings can also be colored by the color i but this is not essential since the string color is determined by the labels of the adjacent 2-cells. We use the following rightward cap and rightward cup to denote the bimodule homomorphisms arising from the natural adjunction between induction and restriction:

$$\begin{aligned} \begin{array}{c} \text{---} i \text{---} \\ \downarrow \\ \text{---} I \text{---} \end{array} & : B_{I,i} \otimes_{R^i} B_{i,I} \rightarrow R^I, & f \otimes g & \mapsto fg, \\ \begin{array}{c} \text{---} I \text{---} \\ \uparrow \\ \text{---} i \text{---} \end{array} & : R^I \rightarrow B_{i,I} \otimes_{R^I} B_{I,i}, & 1 & \mapsto 1 \otimes 1. \end{aligned}$$

The leftward cap and leftward cup denote the bimodule homomorphisms arising from the Frobenius adjunction, i.e., the trace and comultiplication maps (3.5) and (3.6):

$$\begin{aligned} \begin{array}{c} \text{---} i \text{---} \\ \downarrow \\ \text{---} I \text{---} \end{array} & : B_{i,I} \otimes_{R^I} B_{I,i} \rightarrow R^I, & f \otimes g & \mapsto \text{tr}_I^I(fg), \\ \begin{array}{c} \text{---} I \text{---} \\ \uparrow \\ \text{---} i \text{---} \end{array} & : R^I \rightarrow B_{I,i} \otimes_{R^I} B_{i,I}, & 1 & \mapsto \sum_{b \in \mathbb{B}_I^I} b \otimes b^\vee. \end{aligned}$$

The (clockwise) rightward cap and leftward cup are of positive degree $\ell(w_{Ii}) - \ell(w_I)$ and the (counterclockwise) rightward cup and leftward cap are of negative degree $\ell(w_I) - \ell(w_{Ii})$. The adjunctions give the zig-zag relations

$$I \begin{array}{c} \uparrow \\ \text{---} i \text{---} \\ \downarrow \\ \text{---} I \text{---} \end{array} = I \begin{array}{c} \uparrow \\ \text{---} i \text{---} \end{array} = I \begin{array}{c} \uparrow \\ \text{---} i \text{---} \\ \downarrow \\ \text{---} I \text{---} \end{array}, \quad i \begin{array}{c} \downarrow \\ \text{---} I \text{---} \\ \uparrow \\ \text{---} i \text{---} \end{array} = i \begin{array}{c} \downarrow \\ \text{---} I \text{---} \end{array} = i \begin{array}{c} \downarrow \\ \text{---} I \text{---} \\ \uparrow \\ \text{---} i \text{---} \end{array}. \quad (3.17)$$

It makes sense to include an element a of the algebra R^I into a 2-cell labelled I ; we draw this by putting a into a lightly shaded node in the region. We obviously have that

$$I \begin{array}{c} \uparrow \\ \text{---} (f) \text{---} \\ \downarrow \\ \text{---} i \text{---} \end{array} = I \begin{array}{c} \uparrow \\ \text{---} (f) \text{---} \end{array}, \quad i \begin{array}{c} \downarrow \\ \text{---} (f) \text{---} \\ \uparrow \\ \text{---} I \text{---} \end{array} = i \begin{array}{c} \downarrow \\ \text{---} (f) \text{---} \end{array} \quad (3.18)$$

for $f \in R^I \subset R^I$. In particular, elements $f \in R^N$ slide across strings of all colors, i.e., $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{i}\mathfrak{m}$ is an R^N -linear graded bicategory. Closed bubbles evaluate according to the relations

$$\begin{array}{c} \text{---} (f) \text{---} \\ \downarrow \\ \text{---} I \text{---} \end{array} = \boxed{\text{tr}_I^I(f)} \quad i, \quad \begin{array}{c} \text{---} i \text{---} \\ \downarrow \\ \text{---} I \text{---} \end{array} = \boxed{\eta_I^I} \quad I, \quad (3.19)$$

for $f \in R^I$. Also (3.11) can be formulated diagrammatically as

$$i \begin{array}{c} \downarrow \\ \text{---} I \text{---} \\ \uparrow \\ \text{---} i \text{---} \end{array} = \sum_{b \in \mathbb{B}_I^I} \begin{array}{c} \text{---} (b) \text{---} \\ \downarrow \\ \text{---} I \text{---} \\ \uparrow \\ \text{---} i \text{---} \\ \downarrow \\ \text{---} (b^\vee) \text{---} \end{array}. \quad (3.20)$$

There are also upward and downward crossings of strings of different colors i and j , which are the obvious degree 0 bimodule isomorphisms arising from transitivity of induction and restriction:

$$\begin{aligned} \begin{array}{c} \text{---} i \text{---} \\ \swarrow \quad \searrow \\ \text{---} I_j \text{---} \\ \nwarrow \quad \nearrow \\ \text{---} I_i \text{---} \end{array} & : B_{I,I_j} \otimes_{R^{I_j}} B_{I_j,I_i} \rightarrow B_{I,i} \otimes_{R^I} B_{I,i,j}, & f \otimes 1 & \mapsto f \otimes 1, \\ \begin{array}{c} \text{---} I_j \text{---} \\ \swarrow \quad \searrow \\ \text{---} I_i \text{---} \\ \nwarrow \quad \nearrow \\ \text{---} I \text{---} \end{array} & : B_{I_j,I_i} \otimes_{R^I} B_{I,i} \rightarrow B_{I_j,I_j} \otimes_{R^{I_j}} B_{I,i}, & 1 \otimes f & \mapsto 1 \otimes f. \end{aligned}$$

This notation implicitly assumes that i, j are different elements of $N - I$ such that Iij is finitary. The following *easy* Reidemeister II relations are clear:

$$\begin{array}{c} \begin{array}{c} \nearrow I \\ \searrow Ij \\ \nearrow Iij \\ \searrow I \\ \nearrow I \\ \searrow I \\ \nearrow Iij \\ \searrow I \end{array} = \begin{array}{c} \uparrow \\ \uparrow \end{array}, \quad \begin{array}{c} \begin{array}{c} \nearrow Iij \\ \searrow I \\ \nearrow I \\ \searrow Ij \\ \nearrow I \\ \searrow Iij \\ \nearrow I \\ \searrow I \end{array} = \begin{array}{c} \downarrow \\ \downarrow \end{array}. \end{array} \quad (3.21)$$

There are sideways crossings which are defined by rotating the upward or downward ones:

$$\begin{array}{c} \begin{array}{c} \nearrow I \\ \searrow Ij \\ \nearrow Iij \\ \searrow I \end{array} := \begin{array}{c} \nearrow I \\ \searrow Ij \\ \nearrow Iij \\ \searrow I \end{array} = \begin{array}{c} \nearrow I \\ \searrow Ij \\ \nearrow Iij \\ \searrow I \end{array} : B_{Ij,Iij} \otimes_{R^I} B_{Iij,I} \rightarrow B_{Ij,I} \otimes_{R^I} B_{I,Ii}, \quad 1 \otimes 1 \mapsto 1 \otimes 1, \\ \begin{array}{c} \nearrow Iij \\ \searrow I \\ \nearrow I \\ \searrow Ij \\ \nearrow I \\ \searrow Iij \\ \nearrow I \\ \searrow I \end{array} := \begin{array}{c} \nearrow Iij \\ \searrow I \\ \nearrow I \\ \searrow Ij \\ \nearrow I \\ \searrow Iij \\ \nearrow I \\ \searrow I \end{array} = \begin{array}{c} \nearrow Iij \\ \searrow I \\ \nearrow I \\ \searrow Ij \\ \nearrow I \\ \searrow Iij \\ \nearrow I \\ \searrow I \end{array} : B_{Ii,I} \otimes_{R^I} B_{I,Ij} \rightarrow B_{Ii,Iij} \otimes_{R^{Iij}} B_{Iij,Ij}, \quad f \otimes 1 \mapsto \sum_{b \in \mathbb{B}_{Iij}^I} \text{tr}_{Ii}^I(fb) \otimes b^\vee. \end{array}$$

The equalities here reflect the cyclicity of this bicategory. Both sideways crossings have degree $\ell(w_{Iij}) + \ell(w_I) - \ell(w_{Ii}) - \ell(w_{Ij})$, which is half the degree of the element η_{Iij}^I . The mnemonic to remember this degree is: “big plus small minus middle minus middle.” This degree is always non-negative, and it is 0 if and only if i and j live in different connected components of Iij . The *hard* Reidemeister II relations are

$$\begin{array}{c} \begin{array}{c} \nearrow I \\ \searrow Ij \\ \nearrow Iij \\ \searrow I \end{array} = \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} I \\ Iij \\ Ij \end{array}, \quad \begin{array}{c} \begin{array}{c} \nearrow Iij \\ \searrow I \\ \nearrow I \\ \searrow Ij \\ \nearrow I \\ \searrow Iij \\ \nearrow I \\ \searrow I \end{array} = \sum_{b \in \mathbb{B}_{Iij}^I} \begin{array}{c} I \\ Iij \\ Ij \end{array} \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} Iij \\ Ij \\ I \end{array} \end{array} \quad (3.22)$$

for $f \in R^I$. The second of these can be rewritten in several equivalent ways using the following identity, which follows from (3.7) to (3.9):

$$\sum_{b \in \mathbb{B}_{Iij}^I} \text{tr}_{Ii}^I(fb) \otimes b^\vee = \sum_{b \in \mathbb{B}_{Iij}^I} \text{tr}_{Ii}^I(b) \otimes \text{tr}_{Ij}^I(fb^\vee) = \sum_{b \in \mathbb{B}_{Iij}^I} \text{tr}_{Ii}^I(bf) \otimes \text{tr}_{Ij}^I(b^\vee) = \sum_{b \in \mathbb{B}_{Iij}^I} b \otimes \text{tr}_{Ij}^I(fb^\vee) \quad (3.23)$$

in $R^{Ii} \otimes_{R^{Iij}} R^{Ij}$. There is a special case of (3.22) we wish to call out explicitly: when i and j are in distinct connected components of Iij , we have the *distant* Reidemeister II relations

$$\begin{array}{c} \begin{array}{c} \nearrow I \\ \searrow Ij \\ \nearrow Iij \\ \searrow I \end{array} = \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} I \\ I \\ Ij \end{array}, \quad \begin{array}{c} \begin{array}{c} \nearrow Iij \\ \searrow I \\ \nearrow I \\ \searrow Ij \\ \nearrow I \\ \searrow Iij \\ \nearrow I \\ \searrow I \end{array} = \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} Iij \\ Iij \\ Ij \end{array}. \end{array} \quad (3.24)$$

Thus, in this case, the sideways crossings are mutually inverse isomorphisms.

The final general relations from [ESW17] are the Reidemeister III relations, *easy* and *hard*:

$$\begin{array}{c} \begin{array}{c} \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \end{array} = \begin{array}{c} \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \end{array}, \quad \begin{array}{c} \begin{array}{c} \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \end{array} = \begin{array}{c} \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \\ \nearrow Ijk \\ \searrow I \end{array} \end{array} \quad (3.25)$$

Note also that if $\{i, j, k\}$ is not contained in one connected component of $Iijk$, then $\eta_{Iijk}^I = 1$, leading to the *distant* Reidemeister III relation. There are several more variations on Reidemeister III, which may be obtained from the ones above by rotating using the cyclic structure.

Another way to obtain further relations is to apply the graded 2-functor

$$\mathbb{R} : \mathfrak{B} \circ \mathfrak{B} \text{im} \rightarrow \mathfrak{B} \circ \mathfrak{B} \text{im}^{\text{rev}} \quad (3.26)$$

defined as follows: it fixes objects, takes the graded (R^I, R^J) -bimodule M to $q^{\ell(w_J) - \ell(w_I)} M$ viewed as an (R^J, R^I) -bimodule, and takes a bimodule homomorphism to the same underlying function, which is automatically also a homomorphism with respect to the new bimodule structure. In particular, it maps the bimodule $B_{I,K}^J$ to $B_{K,I}^J$. In terms of string diagrams, \mathbb{R} reflects diagrams in a vertical axis and reverses the orientation on all strings.

Finally, we record a useful consequence of the relations described so far. This was originally discovered as [EKL24, Theorem 6.7]. For $i \in I \subseteq_f N$, we write $\hat{I}i$ as a shorthand for $I - \{i\}$.

Lemma 3.4 (Switchback relation). *Suppose we are given I and distinct $j, k \in N - I$ with $Ijk \subseteq_f N$, and also $i \in Ik$ such that $\partial_{Ijk}^{Ij} = \partial_{Iijk}^{Iik} \circ \partial_{Ik}^I$. Then the following relation holds:*

$$\hat{I}ijk \begin{array}{c} \uparrow Ijk \\ \text{---} Iik \text{---} Ik \text{---} I \text{---} Ij \\ \downarrow Ijk \end{array} = \hat{I}ijk \begin{array}{c} \uparrow \\ Ijk \\ \downarrow \end{array} Ij. \quad (3.27)$$

Proof. By (3.22) and easy relations, we have that

$$\begin{aligned} \hat{I}ijk \begin{array}{c} \uparrow Ijk \\ \text{---} Iik \text{---} Ik \text{---} I \text{---} Ij \\ \downarrow Ijk \end{array} &= \sum_{b \in \mathbb{B}_{Ijk}^{Ij}} \hat{I}ijk \begin{array}{c} \uparrow \\ \text{---} Iik \text{---} \boxed{\text{tr}_{Ik}^I(b)} \text{---} Ijk \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ b^\vee \\ \downarrow \end{array} Ij = \sum_{b \in \mathbb{B}_{Ijk}^{Ij}} \hat{I}ijk \begin{array}{c} \uparrow \\ \text{---} Iik \text{---} \boxed{\text{tr}_{Ik}^J(b)} \text{---} Ijk \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ Ijk \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ b^\vee \\ \downarrow \end{array} Ij \\ &= \sum_{b \in \mathbb{B}_{Ijk}^{Ij}} \hat{I}ijk \begin{array}{c} \uparrow \\ \text{---} \boxed{\text{tr}_{Iijk}^{Iik} \text{tr}_{Ik}^I(b)} \text{---} Ijk \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ Ijk \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ b^\vee \\ \downarrow \end{array} Ij = \sum_{b \in \mathbb{B}_{Ijk}^{Ij}} \hat{I}ijk \begin{array}{c} \uparrow \\ \text{---} \boxed{\text{tr}_{Ijk}^{Ij}(b)} \text{---} Ijk \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ Ijk \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ b^\vee \\ \downarrow \end{array} Ij \\ &= \sum_{b \in \mathbb{B}_{Ijk}^{Ij}} \hat{I}ijk \begin{array}{c} \uparrow \\ \text{---} \boxed{\text{tr}_{Ijk}^{Ij}(b)} \text{---} Ijk \\ \downarrow \end{array} \begin{array}{c} \uparrow \\ Ijk \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ b^\vee \\ \downarrow \end{array} Ij = \hat{I}ijk \begin{array}{c} \uparrow \\ Ijk \\ \downarrow \end{array} Ij. \end{aligned}$$

The fourth equality is where we used the hypothesis $\partial_{Ijk}^{Ij} = \partial_{Iijk}^{Iik} \circ \partial_{Ik}^I$. The last equality uses a bubble slide (3.18), followed by the counit axiom (3.11). \square

The hypothesis $\partial_{Ijk}^{Ij} = \partial_{Iijk}^{Iik} \circ \partial_{Ik}^I$ appearing in Lemma 3.4 is an instance of the *switchback relation* introduced in [EK23]. A complete list of all situations in which it holds can be found in [EK23, Sec. 6].

Remark 3.5. The relations recorded in this subsection are not intended to be a complete presentation of $\mathfrak{B} \circ \mathfrak{B} \text{im}$, and indeed no such presentation exists currently in the literature.

3.4. The Soergel-Williamson categorification theorem. Next we give an account of the main results of [Wil11]. Note that our q is denoted v in [Wil11]. We have made a few other expository changes. Our setup is biased towards writing translation functors on the left, and consequently many of the statements below have been obtained from the ones in [Wil11] by twisting with the anti-involution ρ on the q -Schur algebra from (3.33) or the symmetry \mathbb{R} on singular Bott-Samelson bimodules from (3.26).

Let H be the Hecke algebra associated to the Coxeter group W over the ground ring $\mathbb{Z}[q, q^{-1}]$. This is the free $\mathbb{Z}[q, q^{-1}]$ -module with basis $\{h_w \mid w \in W\}$ viewed as an $\mathbb{Z}[q, q^{-1}]$ -algebra with multiplication satisfying

$$h_i h_w = \begin{cases} h_{s_i w} & \text{if } \ell(s_i w) > \ell(w) \\ h_{s_i w} - (q - q^{-1})h_w & \text{if } \ell(s_i w) < \ell(w) \end{cases} \quad (3.28)$$

for each $w \in W$ and $i \in I$, where h_i denotes h_{s_i} for short. In particular, $(h_i + q)(h_i - q^{-1}) = 1$ and $h_i^{-1} = h_i + (q - q^{-1})$. The *bar involution* is the anti-linear algebra involution $H \rightarrow H, h \mapsto \bar{h}$ defined so that $\bar{\bar{h}} := h_{w^{-1}}$ for each $w \in W$. Also useful is the linear algebra anti-involution $\rho : H \rightarrow H, h_w \mapsto h_{w^{-1}}$. It commutes with the bar involution, so $\omega : H \rightarrow H, h \mapsto \rho(\bar{h})$ is an anti-linear algebra anti-involution fixing each basis vector h_w ($w \in W$). We denote the *Kazhdan-Lusztig basis* of H (renormalized as in [Soe97]) by b_w ($w \in W$). So b_w is the unique bar-invariant element of $h_w + \sum_{w' < w} q\mathbb{Z}[q]h_{w'}$. We have that $\rho(b_w) = b_{w^{-1}}$.

For $I \subseteq_f N$, let H_I be the subalgebra of H with basis h_w ($w \in W_I$). The longest Kazhdan-Lusztig basis element of H_I is

$$b_{w_I} = \sum_{w \in W_I} q^{\ell(w_I) - \ell(w)} h_w. \quad (3.29)$$

Note that $h_i b_{w_I} = b_{w_I} h_i = q^{-1} b_{w_I}$ for all $i \in I$. It follows that

$$b_{w_I} b_{w_I} = \pi_I b_{w_I} \quad \text{where} \quad \pi_I := \sum_{w \in W_I} q^{\ell(w_I) - 2\ell(w)}. \quad (3.30)$$

Here, π_I is the renormalized Poincaré polynomial from (3.1).

Given $I, J \subseteq_f N$, we are going to consider the set $W_I \backslash W / W_J$ of (W_I, W_J) -double cosets in W . We let $(W_I \backslash W / W_J)_{\min}$ denote the set of minimal length coset representatives. For $d \in (W_I \backslash W / W_J)_{\min}$ we often use the shorthand IdJ to succinctly encode the triple of I, d and J , hence, the (W_I, W_J) -double coset $W_I d W_J$. On the other hand, given just the set $W_I d W_J$, one can recover d as the minimal length element, but I and J themselves are not uniquely determined. For $d \in (W_I \backslash W / W_J)_{\min}$, let

$$\text{LR}(IdJ) := \{i \in I \mid s_i d = d s_j \text{ for some } j \in J\}. \quad (3.31)$$

This set is the *left redundancy*⁶ of the double coset $W_I d W_J$. We have that $W_I \cap d W_J d^{-1} = W_{\text{LR}(IdJ)}$ (Kilmoyer's theorem); this is the stabilizer of $d W_J$ under the natural left action of W_I on W / W_J . Elements of $W_I d W_J$ can be written uniquely as udv for $u \in (W_I / W_{\text{LR}(IdJ)})_{\min}$, $v \in W_J$, and we have that $\ell(udv) = \ell(u) + \ell(d) + \ell(v)$ (Howlett's theorem). The unique element of maximal length in the double coset $W_I d W_J$ is $w_I w_{\text{LR}(IdJ)}^{-1} d w_J$, of length

$$\ell(IdJ) := \ell(w_I) - \ell(w_{\text{LR}(IdJ)}) + \ell(d) + \ell(w_J). \quad (3.32)$$

Definition 3.6 ([Wil11, Sec. 2.3]). The (generalized) *q-Schur algebra*⁷ is

$$\mathbb{S} := \bigoplus_{I, J \subseteq_f N} (b_{w_I} H) \cap (H b_{w_J}),$$

which is a direct sum of free $\mathbb{Z}[q, q^{-1}]$ -submodules of H . We view it as a $\mathbb{Z}[q, q^{-1}]$ -algebra with multiplication $\circ : \mathbb{S} \times \mathbb{S} \rightarrow \mathbb{S}$ defined by

$$x \circ y := \begin{cases} \frac{1}{\pi_J} xy & \text{if } J = J' \\ 0 & \text{otherwise,} \end{cases}$$

⁶Analogously, there is also the *right redundancy* $\text{RR}(IdJ) := \{j \in J \mid d s_j = s_i d \text{ for some } i \in I\}$.

⁷In [Wil11], this is called the *Hecke algebroid*, and others refer to it as the *Schur algebroid*.

$I, J, J', K \subseteq N$, $x \in (b_{w_I}H) \cap (Hb_{w_J})$ and $y \in (b_{w_{J'}}H) \cap (Hb_{w_K})$; here, we have used (3.30) to see that xy is divisible by π_J in the first case. (This algebra could also be introduced by starting from the $\mathbb{Z}[q, q^{-1}]$ -linear category \mathbf{S} with objects $I \subseteq_f N$ and morphisms $\text{Hom}(J, I) := (b_{w_I}H) \cap (Hb_{w_J})$, with the operation \circ just defined being its composition law. Then \mathbf{S} is the path algebra of \mathbf{S} . We will only use \mathbf{S} not \mathbf{S} subsequently.)

We denote the element b_{w_I} of the summand $(b_{w_I}H) \cap (Hb_{w_I})$ of \mathbf{S} by 1_I (by itself, b_{w_I} is ambiguous as an element of \mathbf{S}). We then have that $1_I \circ 1_J = \delta_{I,J}1_I$. So the elements 1_I ($I \subseteq_f N$) are mutually orthogonal idempotents summing to the identity in \mathbf{S} , and $1_I \circ \mathbf{S} \circ 1_J = (b_{w_I}H) \cap (Hb_{w_J})$. The Hecke algebra \mathbf{H} itself is naturally identified with the idempotent truncation $1_\emptyset \circ \mathbf{S} \circ 1_\emptyset$ (on which \circ coincides with the usual multiplication). The *bar involution* $\mathbf{S} \rightarrow \mathbf{S}$, $h \mapsto \bar{h}$ is the anti-linear algebra involution defined on the summand $1_I \circ \mathbf{S} \circ 1_J$ by the restriction of the bar involution from \mathbf{H} . There is also a linear algebra anti-involution

$$\rho : \mathbf{S} \rightarrow \mathbf{S} \quad (3.33)$$

defined by the maps $1_I \circ \mathbf{S} \circ 1_J \rightarrow 1_J \circ \mathbf{S} \circ 1_I$ obtained by the restrictions of $\rho : \mathbf{H} \rightarrow \mathbf{H}$ for all $I, J \subseteq_f N$. Again, ρ commutes with the bar involution, so the composition $\omega : \mathbf{S} \rightarrow \mathbf{S}$ of ρ and the bar involution (in either order) is an anti-linear algebra anti-involution of \mathbf{S} .

For our purposes, there are two important bases of \mathbf{S} , the *standard basis* $\{h_{IdJ}\}$ and the *Kazhdan-Lusztig basis* $\{b_{IdJ}\}$, both indexed by the symbols IdJ for $I, J \subseteq_f N$ and $d \in (W_I \backslash W / W_J)_{\min}$. They are defined by

$$h_{IdJ} := \sum_{w \in W_I d W_J} q^{\ell(IdJ) - \ell(w)} h_w, \quad b_{IdJ} := b_{w_I w_{\text{LR}(IdJ)}^{-1} d w_J}, \quad (3.34)$$

both viewed as elements of the summand $1_I \circ \mathbf{S} \circ 1_J$. We have that $b_{II} = h_{II} = 1_I$. Also

$$\rho(h_{IdJ}) = h_{Jd^{-1}I}, \quad \rho(b_{IdJ}) = b_{Jd^{-1}I}. \quad (3.35)$$

In general, the Kazhdan-Lusztig basis element b_{IdJ} can be characterized as the unique bar-invariant element of

$$h_{IdJ} + \sum_{\substack{d' \in (W_I \backslash W / W_J)_{\min} \\ d' < d}} q\mathbb{Z}[q]h_{Id'J}.$$

Since 1 is minimal, we have that $b_{II} = h_{II}$. We denote this instead by $b_{I,I}$; this parallels the notation for the bimodules $B_{I,J}$ in (3.12) and their generalizations introduced just after (3.15). These special elements generate \mathbf{S} as an algebra. In fact, \mathbf{S} is already generated by $b_{I,I}$ and $b_{i,i}$ for $I \subset N$ and $i \in N - I$ with Ii finitary. We also let

$$b_{I,J}^K := b_{I,K} \circ b_{K,J} \quad (3.36)$$

for $I, J \subseteq_f N$ and $K \subseteq I \cap J$; this parallels (3.15).

Lemma 3.7 ([Wil11, Prop. 2.8]). *The following hold for any $I, J, K \subseteq_f N$ with $I \subseteq J$ or $I \supseteq J$ and $d \in (W_J \backslash W / W_K)_{\min}$:*

(1) *If $I \subseteq J$ then*

$$b_{I,J} \circ h_{JdK} = \sum_{d' \in W_J d W_K \cap (W_I \backslash W / W_K)_{\min}} q^{\ell(JdK) - \ell(Id'K)} h_{Id'K}.$$

(2) *If $I \supseteq J$ then*

$$b_{I,J} \circ h_{JdK} = q^{\ell(d') - \ell(d)} \frac{\pi_{\text{LR}}(Id'K)}{\pi_{\text{LR}}(JdK)} h_{Id'K}$$

where d' is the minimal length element of $W_I d W_K$.

Corollary 3.8. *For $I, J \subseteq_f N$ and $K \subseteq I \cap J$, we have that $b_{I,J}^K = \frac{\pi_{I \cap J}}{\pi_K} b_{I,J}$.*

The final basic notion is a bilinear form $(-, -) : \mathbb{S} \times \mathbb{S} \rightarrow \mathbb{Z}[q, q^{-1}]$. Let $\text{tr} : \mathbb{H} \rightarrow \mathbb{Z}[q, q^{-1}]$ be the usual symmetric Frobenius trace on the Hecke algebra, i.e., the $\mathbb{Z}[q, q^{-1}]$ -linear map with $\text{tr}(h_w) = \delta_{w,1}$ for all $w \in W$. Then, $(-, -)$ is defined so that different summands $1_I \circ \mathbb{S} \circ 1_J$ are orthogonal to each other, and

$$(x, y) := \frac{1}{\pi_I} \text{tr}(\rho(x)y) \quad (3.37)$$

for $x, y \in 1_I \circ \mathbb{S} \circ 1_J$. From the definition of the form, we have that $(zx, y) = (x, \rho(z)y)$ for any $x, y, z \in \mathbb{S}$. The standard basis of \mathbb{S} is an orthogonal basis with respect to this form, with

$$(h_{IdJ}, h_{IdJ}) := q^{\ell(IdJ) - \ell(d)} \frac{\pi_J}{\pi_{\text{LR}(IdJ)}}. \quad (3.38)$$

Hence, the form is symmetric. This all follows from [Wil11, Lem. 2.13], noting that the form $\langle -, - \rangle$ there is related to ours by $\langle x, y \rangle = (\rho(x), \rho(y))$. We record also an equivalent way of writing (3.38) which is convenient when applying Theorem 3.10 below:

$$q^{\ell(w_I)} \overline{(h_{IdJ}, h_{IdJ})} = \frac{\tilde{\pi}_J}{\tilde{\pi}_{\text{LR}(IdJ)}} \in 1 + \mathbb{N}[q^{-2}]. \quad (3.39)$$

Now we come to the results from [Wil11] relating the q -Schur algebra to singular Soergel bimodules. Suppose that $I, J \subseteq N$. As well as singular Soergel bimodules, it is important to consider the graded (R^I, R^J) -bimodule $R_d^{\text{LR}(IdJ)}$ for $d \in (W_I \setminus W/W_J)_{\min}$. This denotes $R^{\text{LR}(IdJ)}$ viewed as a graded (R^I, R^J) -bimodule, with action defined by $a \cdot v \cdot b := avd(b)$. By [Wil11, Lem. 4.2(1)], we have that

$$R_d^{\text{LR}(IdJ)} \cong \begin{cases} q^{\ell(w_{\text{LR}(IdJ)}) - \ell(w_I)} \frac{\pi_I}{\pi_{\text{LR}(IdJ)}} R^I & \text{as a graded left } R^I\text{-module} \\ q^{\ell(w_{\text{LR}(IdJ)}) - \ell(w_J)} \frac{\pi_J}{\pi_{\text{LR}(IdJ)}} R^J & \text{as a graded right } R^J\text{-module.} \end{cases} \quad (3.40)$$

Also [Wil11, Cor. 4.13] gives that

$$\text{Hom}_{R^I - R^J}(R_d^{\text{LR}(IdJ)}, R_{d'}^{\text{LR}(Id'J)}) \cong \begin{cases} R^{\text{LR}(IdJ)} & \text{if } d = d' \\ \{0\} & \text{otherwise,} \end{cases} \quad (3.41)$$

for any $d, d' \in (W_I \setminus W/W_J)_{\min}$. The *costandard* and *standard bimodules* are

$$\nabla_{IdJ} := q^{\ell(IdJ) - \ell(w_J)} R_d^{\text{LR}(IdJ)}, \quad (3.42)$$

$$\Delta_{IdJ} := q^{-2\ell(d)} \nabla_{IdJ}, \quad (3.43)$$

respectively. In the special case $d = 1$, we have that $B_{I,J} := \nabla_{11J} = \Delta_{11J}$; these are the same as the bimodules introduced just after (3.15).

The 2-functor \mathbb{R} defined in (3.26) makes sense on all graded (R^I, R^J) -bimodules, not just on Bott-Samelsons. It reverses tensor products, mapping a graded (R^I, R^J) -bimodule M to $q^{\ell(w_J) - \ell(w_I)} M$ viewed as a graded (R^J, R^I) -bimodule. It is clear from (3.42) and (3.43) that

$$\mathbb{R}(\Delta_{IdJ}) = \Delta_{Jd^{-1}I}, \quad \mathbb{R}(\nabla_{IdJ}) = \nabla_{Jd^{-1}I}. \quad (3.44)$$

In particular, we have that $\mathbb{R}(B_{I,J}) = B_{J,I}$. There is also a duality functor \mathbb{D} defined on a graded (R^I, R^J) -bimodule M by

$$\mathbb{D}(M) := q^{2\ell(w_I) - 2\ell(w_J)} \text{Hom}_{-R^J}(M, R^J), \quad (3.45)$$

viewed as a graded (R^I, R^J) -bimodule in the natural way. Arguing as in the proof of [Wil11, Prop. 6.17], remembering the modified grading shifts in (3.42) and (3.43), we have that

$$\mathbb{D}(\nabla_{IdJ}) \cong \Delta_{IdJ}, \quad \mathbb{D}(\Delta_{IdJ}) \cong \nabla_{IdJ}. \quad (3.46)$$

In particular, this shows that $\mathbb{D}(B_{I,J}) \cong B_{I,J}$. Another important point is that

$$\mathbb{D}(M \otimes_{R^J} M') \cong \mathbb{D}(M) \otimes_{R^J} \mathbb{D}(M'), \quad (3.47)$$

for graded (R^I, R^J) - and (R^J, R^K) -bimodules M and M' such that M is a singular Soergel bimodule. It suffices to prove this in the special case that M is the self-dual bimodule B_{IJ} with $I \subseteq J$ or $I \supseteq J$. When $I \supseteq J$, the result follows using that restriction commutes with duality (tautologically), then one deduces the result when $I \subseteq J$ using the adjunctions (3.13) and (3.14); cf. [Wil11, Prop. 6.15]. This argument also shows that all of the singular Bott-Samelson bimodules (3.16) are self-dual.

Remark 3.9. We have changed the grading shift in the definition of duality compared to [Wil11], which forces several other changes. Our choice of grading shift on duality is one of two sensible ways to correct a minor error in Williamson’s original setup. This is also explained in [Wil23], where it is described as the “more comprehensive fix.”

Let \leq be a total order on $(W_I \backslash W / W_J)_{\min}$ refining the Bruhat order \leq . The next definitions are independent of this choice (up to isomorphism) thanks to the hin-und-her lemmas in [Wil11]. For $d \in (W_I \backslash W / W_J)_{\min}$ and an (R^I, R^J) -bimodule M , we define the subquotients $\Gamma_{\leq d} M / \Gamma_{< d} M, \Gamma_{\geq d} M / \Gamma_{> d} M$ of M in terms of their supports, as explained in [Wil11, Sec. 4.5].

- Following [Wil11, Def. 6.1], a graded (R^I, R^J) -bimodule M has a ∇ -flag if there exist (necessarily unique) Laurent polynomials $(M : \nabla_{IdJ})_q \in \mathbb{N}[q, q^{-1}]$ such that $\Gamma_{\leq d} M / \Gamma_{< d} M \cong (M : \nabla_{IdJ})_q \nabla_{IdJ}$ for all $d \in (W_I \backslash W / W_J)_{\min}$. Assuming M has a ∇ -flag, we define its ∇ -character to be

$$\text{ch}_{\nabla}(M) := \sum_{d \in (W_I \backslash W / W_J)_{\min}} (M : \nabla_{IdJ})_q \overline{h_{IdJ}} \in 1_I \circ \mathbb{S} \circ 1_J. \quad (3.48)$$

Note also that our ch_{∇} is the one in [Wil23] post-composed with the bar involution. This is a harmless change which seems more natural to us since it leads to (3.50), the idea being that ∇ -flags are dual to the Δ -flags which are introduced next.

- Following [Wil11, Def. 6.12], a graded (R^I, R^J) -bimodule M has a Δ -flag there exist $(M : \Delta_{IdJ})_q \in \mathbb{N}[q, q^{-1}]$ such that $\Gamma_{\geq d} M / \Gamma_{> d} M \cong (M : \Delta_{IdJ})_q \Delta_{IdJ}$ for all $d \in (W_I \backslash W / W_J)_{\min}$. Assuming M has a Δ -flag, its Δ -character is

$$\text{ch}_{\Delta}(M) := \sum_{d \in (W_I \backslash W / W_J)_{\min}} (M : \Delta_{IdJ})_q h_{IdJ} \in 1_I \circ \mathbb{S} \circ 1_J. \quad (3.49)$$

This is the same as in [Wil23].

The duality \mathbb{D} interchanges the two sorts of flag. Moreover, by (3.46), we have that

$$\text{ch}_{\Delta}(\mathbb{D}(M)) = \overline{\text{ch}_{\nabla}(M)} \quad (3.50)$$

on bimodules with ∇ -flags.

Now we can state the main results. These identify the q -Schur algebra \mathbb{S} with $K_0(\text{gKar}(\mathfrak{B} \subseteq \mathfrak{B}\text{im}))$, the split Grothendieck ring consisting of degree 0 isomorphism classes of singular Soergel bimodules.

Theorem 3.10 (Homomorphism formula [Wil11, Th. 7.9]). *Let M, M' be graded (R^I, R^J) -bimodules such that either M is a singular Soergel bimodule and M' has a ∇ -flag, or M has a Δ -flag and M' is a singular Soergel bimodule. There is an isomorphism*

$$\text{Hom}_{R^I - R^J}(M, M') \cong q^{\ell(w_I)} \overline{(\text{ch}_{\Delta}(M), \text{ch}_{\nabla}(M'))} R^J$$

of graded R^J -modules.

Theorem 3.11 (Classification of indecomposables [Will1, Th. 7.10]). *For each $d \in (W_I \backslash W / W_J)_{\min}$, there is a unique (up to isomorphism) self-dual indecomposable graded (R^I, R^J) -bimodule B_{IdJ} such that $B_{IdJ} / \Gamma_{<d} B_{IdJ} \cong \nabla_{IdJ}$ (equivalently, $B_{IdJ} / \Gamma_{>d} B_{IdJ} \cong \Delta_{IdJ}$). These bimodules for all I, J and d give a full set of pairwise inequivalent indecomposable 1-morphisms in $\mathfrak{gKar}(\mathfrak{B} \circlearrowleft \mathfrak{Bim})$ (up to grading shifts).*

Theorem 3.12 (Categorification theorem [Will1, Th. 7.12]). *If M is a singular Soergel bimodule then it has both a Δ -flag and a ∇ -flag, and*

$$\mathrm{ch}(M) := \mathrm{ch}_{\Delta}(M) = \mathrm{ch}_{\nabla}(M). \quad (3.51)$$

The map ch induces a $\mathbb{Z}[q, q^{-1}]$ -algebra isomorphism $\mathrm{ch} : K_0(\mathfrak{gKar}(\mathfrak{B} \circlearrowleft \mathfrak{Bim})) \xrightarrow{\sim} \mathcal{S}$ taking the isomorphism class of the graded (R^I, R^I) -bimodule R^I to 1_I for each $I \subseteq N$. It intertwines the anti-linear involution of $K_0(\mathfrak{gKar}(\mathfrak{B} \circlearrowleft \mathfrak{Bim}))$ arising from the duality \mathbb{D} with the bar involution on \mathcal{S} , and it intertwines the linear anti-involution arising from the symmetry \mathcal{R} with ρ .

4. EXTENDED SINGULAR SOERGEL BIMODULES IN TYPE D_l

Now we specialize to the case of interest in this article: singular Soergel bimodules for the Weyl group W of type D_l . We will actually work with a variant which we call *extended* singular Soergel bimodules, the purpose of which we explain in Subsection 4.2. We also introduce an extended version of the q -Schur algebra, and compute explicitly the Kazhdan-Lusztig basis of the part of it that relates to nil-Brauer. Finally, we use the extended diagrammatic calculus to derive some difficult relations.

4.1. Realization of the root system of type D_l . Fix $l \geq 2$ and consider the root system of type D_l . We label the nodes of the Dynkin diagram by the set $N := \{\pm 1, 2, \dots, l-1\}$ as in the introduction. We adopt the usual reflection realization of the corresponding Coxeter group W , which satisfies all of the hypotheses assumed in Subsection 3.1. So we let \mathfrak{h} be the vector space with basis x_1, \dots, x_l , and identify \mathfrak{h}^* with \mathfrak{h} via the non-degenerate symmetric bilinear form defined by declaring that this basis is orthonormal. The root system Φ is $\{\pm x_j \pm x_i \mid 1 \leq i, j \leq l, i \neq j\}$, and we have simply that $\alpha^\vee = \alpha$ for each $\alpha \in \Phi$. We make the following choice for simple roots:

$$\begin{aligned} \alpha_1 &:= x_2 - x_1 \\ \alpha_2 &:= x_3 - x_2 \quad \alpha_3 := x_4 - x_3 \quad \cdots \quad \alpha_{l-1} := x_l - x_{l-1} \\ \alpha_{-1} &:= x_2 + x_1 \end{aligned}$$

The corresponding set of positive roots is $\{x_j \pm x_i \mid 1 \leq i < j \leq l\}$. For $i \in N$, the simple reflection s_i is the reflection in the hyperplane orthogonal to α_i . Except when $i = -1$, s_i permutes x_i and x_{i+1} fixing all other x_j , while s_{-1} permutes x_1 and $-x_2$ fixing all other x_j .

A useful additional feature is the existence of the *graph automorphism* $\gamma : N \rightarrow N$. This switches 1 and -1 and fixes all other elements. Let $s_0 : \mathfrak{h} \rightarrow \mathfrak{h}$ be the reflection with $s_0(x_1) = -x_1$, fixing all other x_j . This corresponds to γ in that $s_0(\alpha_i) = \alpha_{\gamma(i)}$ for each $i \in N$. The reflections s_0, s_1, \dots, s_{l-1} generate the Weyl group of type B_l , with the Weyl group W of type D_l being the subgroup generated by $s_{-1} = s_0 s_1 s_0, s_1, \dots, s_{l-1}$. All s_i extend to algebra automorphisms of $R = \mathbb{k}[x_1, \dots, x_l]$ (defined in the previous section to be the symmetric algebra of \mathfrak{h}^*). We prefer to denote the automorphism of R defined by s_0 by

$$\gamma : R \rightarrow R \quad f \mapsto s_0(f). \quad (4.1)$$

We reserve the unusual letter O for the special subset $\{2, \dots, l-1\} \subset N$, since it appears very often. Remember that we write simply Ii for $I \cup \{i\}$ (assuming $i \notin I$) and $I\hat{i}$ for $I - \{i\}$ (assuming $i \in I$). Mainly because they look nicer in diagrams, we will use the following shorthands:

$$I+ := I1, \quad I- := I(-1), \quad I\pm := I1(-1) \quad (4.2)$$

for $I \subseteq O$. So we are using the symbol $+$ to indicate inclusion of 1, the symbol $-$ to indicate inclusion of -1 , and \pm to indicate the inclusion of both. The graph automorphism γ interchanges $+$ and $-$. Recall also for $I \subseteq N$ that R^I denotes the invariant subalgebra R^{W_I} of R . The following table gives some useful facts about some of these algebras and the corresponding parabolic subgroups of W , for $n \in O$:

I	Type of W_I	$\ell(w_I)$	π_I
$O\pm$	D_l	$2\binom{l}{2}$	$[2]_q^{l-1}[l-1]_{q^2}[l]_q$
$O+$ or $O-$	A_{l-1}	$\binom{l}{2}$	$[l]_q!$
O	A_{l-2}	$\binom{l-1}{2}$	$[l-1]_q!$
$O\hat{n}\pm$	$A_{l-n-1} \times D_n$	$\binom{l-n}{2} + 2\binom{n}{2}$	$[l-n]_q![2]_q^{n-1}[n-1]_{q^2}[n]_q$
$O\hat{n}+$ or $O\hat{n}-$	$A_{l-n-1} \times A_{n-1}$	$\binom{l-n}{2} + \binom{n}{2}$	$[l-n]_q![n]_q!$
$O\hat{n}$	$A_{l-n-1} \times A_{n-2}$	$\binom{l-n}{2} + \binom{n-1}{2}$	$[l-n]_q![n-1]_q!$

(4.3)

Also, here are explicit descriptions of some of the algebras of invariants:

- R^N is free with generators given by the elementary symmetric polynomials $e_r(x_1^2, \dots, x_l^2)$ for $r = 1, \dots, l-1$ together with $e_l(x_1, \dots, x_l) = x_1 \cdots x_l$.
- R^{O+} is the algebra A of symmetric polynomials which appeared already in the introduction and Remark 2.2. It is freely generated by $e_r(x_1, \dots, x_l)$ for $r = 1, \dots, l$.
- $R^O = \mathbb{k}[x_1, x_2, \dots, x_l]^{S_1 \times S_{l-1}}$, which is generated by x_1 and $e_r(x_2, \dots, x_l)$ for $r = 1, \dots, l-1$. This was seen in the introduction when the (A, A) -bimodule B was defined.

4.2. Motivational interlude. We will be interested primarily in double cosets $W_I \backslash W / W_J$ where $I, J \in \{O+, O-\}$, and in the corresponding categories of singular Bott-Samelson bimodules, that is, we wish to study in depth the sub-bicategory of $\mathfrak{B} \mathfrak{S} \mathfrak{B} \text{im}$ with two objects $\{R^{O+}, R^{O-}\}$, and the corresponding piece of \mathfrak{S} . Our goal is to relate these bimodules to the cyclotomic nilBrauer category \mathfrak{cNB}_l , but there is some awkward book-keeping involved in this since we are comparing a monoidal category with a bicategory with two objects.

The approach we found to be most convenient is to transform any R^{O-} -module into an R^{O+} -module using the graph automorphism. In this way, we are able to replace the aforementioned bicategory-with-two-objects simply with a monoidal category of (R^{O+}, R^{O+}) -bimodules. See Subsection 4.5 for details. The Grothendieck group of this formal construction is a piece of the extended q -Schur algebra described in the next section. The general approach taken also fits well with the ι -Schur-Weyl duality from [BW18b, Ch. 5], although we will not discuss this further here.

An alternate approach would be to instead replace \mathfrak{NB}_l with a 2-category with two objects. This is analogous to the use of the two-colored Temperley-Lieb algebra in [Eli17]. There are advantages to this approach, e.g., one can eliminate the pesky sign in the first relations from (2.9) and (2.12) by rescaling the dot based on the ambient colors. However, the disadvantage of having to keep track of two versions of each bimodule seemed to us to be more burdensome. For example, in the two-colored approach, one has to work with two generating bimodules $B_{O-,O} \otimes_{R^O} B_{O,O+}$ and $B_{O+,O} \otimes_{R^O} B_{O,O-}$. In the extended category, due to the added twists, they both yield the same bimodule over (R^{O+}, R^{O+}) .

4.3. The extended q -Schur algebra. Let \mathfrak{S} be the q -Schur algebra of type D_l as in Definition 3.6. The graph automorphism defines a $\mathbb{Z}[q, q^{-1}]$ -algebra involution $\gamma : \mathfrak{H} \rightarrow \mathfrak{H}, h_i \mapsto h_{\gamma(i)}$ of the Hecke algebra, which extends to an automorphism $\gamma : \mathfrak{S} \rightarrow \mathfrak{S}$ taking 1_I to $1_{\gamma(I)}$. The *extended q -Schur algebra* is the $\mathbb{Z}[q, q^{-1}]$ -module $\mathfrak{S} \oplus s_0 \mathfrak{S}$, where $s_0 \mathfrak{S}$ is a copy of \mathfrak{S} with elements denoted $s_0 x$ ($x \in \mathfrak{S}$), viewed as a $\mathbb{Z}[q, q^{-1}]$ -algebra with the extended multiplication \circ defined by

$$(x + s_0 y) \circ (u + s_0 v) := (x \circ u + \gamma(y) \circ v) + s_0 (y \circ u + \gamma(x) \circ v) \quad (4.4)$$

for $x, y, u, v \in S$. From the standard and Kazhdan-Lusztig bases for S , we obtain standard and Kazhdan-Lusztig bases also for $S \oplus s_0 S$. These consist of the elements $\{h_{IdJ}, s_0 h_{IdJ}\}$ or the elements $\{b_{IdJ}, s_0 b_{IdJ}\}$ for $I, J \subseteq N$ and $d \in (W_I \backslash W / W_J)_{\min}$. The Kazhdan-Lusztig basis is invariant under the bar involution on $S \oplus s_0 S$, which is defined simply by

$$\overline{x + s_0 y} := \bar{x} + s_0 \bar{y}. \quad (4.5)$$

The linear anti-involution ρ from (3.33) extends to $S \oplus s_0 S$ by setting

$$\rho(x + s_0 y) := \rho(x) + s_0 \gamma(\rho(y)). \quad (4.6)$$

Finally, we extend the symmetric bilinear form on S from (3.37) to $S \oplus s_0 S$ by

$$(x + s_0 y, u + s_0 v) := (x, u) + (y, v). \quad (4.7)$$

Remark 4.1. The Weyl group W , its Hecke algebra H and the q -Schur algebra S here are of type D_l . The Weyl group of type B_l is $W \sqcup s_0 W$, and $H \oplus s_0 H$ with twisted multiplication defined like in (4.4) is the Hecke algebra of type B_l at unequal parameters, with the short root parameter being 1. Similarly, our extended q -Schur algebra $S \oplus s_0 S$ can be related to the $(q, 1)$ -Schur algebra of type B_l . We will not use this observation here so leave the formulation of a more precise statement to the reader; the construction in [Bao19] is also somewhat relevant.

4.4. The subalgebra V and its Kazhdan-Lusztig basis. In the remainder of the paper, the focus will be on the subalgebra

$$V := 1_{O_+} \circ (S \oplus s_0 S) \circ 1_{O_+} = (1_{O_+} \circ S \circ 1_{O_+}) \oplus s_0 (1_{O_-} \circ S \circ 1_{O_+}) \quad (4.8)$$

of $S \oplus s_0 S$, which has the idempotent 1_{O_+} as its identity element. We are now going to describe its Kazhdan-Lusztig basis explicitly. In fact, we will show that it is combinatorially the same as the ι -canonical basis of $V(l)$ seen already in Theorem 2.14. We denote the special element $s_0 b_{O_-, O_+} \in V$ simply by b (it will turn out to be “the same” as the b in Section 2):

$$b := s_0 b_{O_-, O_+} \stackrel{(3.36)}{=} s_0 b_{O_-, O} \circ b_{O, O_+}. \quad (4.9)$$

Since $\gamma(b_{O_-, O_+}) = b_{O_+, O_-}$, we have that

$$b^{\circ n} = \begin{cases} 1_{O_+} & \text{if } n = 0 \\ (b_{O_+, O_-} \circ b_{O_-, O_+})^{\circ \frac{n}{2}} & \text{if } n \text{ is even} \\ s_0 b_{O_-, O_+} \circ (b_{O_+, O_-} \circ b_{O_-, O_+})^{\circ \frac{n-1}{2}} & \text{if } n \text{ is odd.} \end{cases} \quad (4.10)$$

By Corollary 4.6 below, the elements $b^{\circ n}$ ($0 \leq n \leq l$) give a rational basis for V , i.e., they are a basis for the $\mathbb{Q}(q)$ -vector space $\mathbb{Q}(q) \otimes_{\mathbb{Z}[q, q^{-1}]} V$.

We will start now to draw elements of $W \sqcup s_0 W$ by permutation-type diagrams with symmetry about the middle axis, labeling the boundary in order from left to right by $-x_l, \dots, -x_2, -x_1; x_1, x_2, \dots, x_l$. For example, the following are the pictures for s_0, s_1 and s_2 when $l = 3$:

$$s_0 = \begin{array}{c} \begin{array}{ccccccc} -x_3 & -x_2 & -x_1 & x_1 & x_2 & x_3 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \\ -x_3 & -x_2 & -x_1 & x_1 & x_2 & x_3 & \end{array} \\ \text{,} \end{array} \quad s_1 = \begin{array}{c} \begin{array}{ccccccc} -x_3 & -x_2 & -x_1 & x_1 & x_2 & x_3 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \\ -x_3 & -x_2 & -x_1 & x_1 & x_2 & x_3 & \end{array} \\ \text{,} \end{array} \quad s_2 = \begin{array}{c} \begin{array}{ccccccc} -x_3 & -x_2 & -x_1 & x_1 & x_2 & x_3 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \\ -x_3 & -x_2 & -x_1 & x_1 & x_2 & x_3 & \end{array} \\ \text{.} \end{array}$$

Lemma 4.4. *The set $(W_{O-} \backslash W/W_{O+})_{\min}$ equals $\{s_0 d_n \mid 1 \leq n \leq l \text{ with } n \text{ odd}\}$. For these values of n , we have that*

$$\text{LR}((O-)s_0 d_n(O+)) = \begin{cases} O & \text{if } n = 1 \\ O- & \text{if } n = l \\ O\hat{n}- & \text{otherwise.} \end{cases}$$

Also

$$\begin{aligned} W_{O-} s_0 d_n W_{O+} &= W_{O-} s_0 d_n W_{O+} \sqcup W_{O-} d_{n+1} W_{O+} && \text{for odd } 1 \leq n \leq l-1, \\ W_{O-} s_0 d_l W_{O+} &= W_{O-} s_0 d_l W_{O+} && \text{when } l \text{ is odd.} \end{aligned}$$

Proof. Similar arguments to the previous two lemmas. \square

Let $t := \frac{(-1)^{l-1} + 1}{2}$, which is 0 or 1 according to whether l is even or odd. Recall the U_t^l -module $V(l)$ from Subsection 2.6 with its monomial basis $f^{(n)} \eta_l$ ($0 \leq n \leq l$) and t -canonical basis $b^{(n)} \eta_l$ ($0 \leq n \leq l$), and the $\mathbb{Z}[q, q^{-1}]$ -algebra V from (4.8), with its standard and Kazhdan-Lusztig bases. Both spaces have also been equipped with a bar involution and a symmetric bilinear form.

Theorem 4.5. *There is a unique isomorphism of free $\mathbb{Z}[q, q^{-1}]$ -modules*

$$\begin{aligned} \phi : V(l) &\xrightarrow{\sim} V, \\ \overline{f^{(n)} \eta_l} &\mapsto \begin{cases} h_{(O+)d_n(O+)} & \text{for even } 0 \leq n \leq l \\ s_0 h_{(O-)s_0 d_n(O+)} & \text{for odd } 1 \leq n \leq l, \end{cases} \end{aligned} \quad (4.11)$$

$$b^{(n)} \eta_l \mapsto \begin{cases} b_{(O+)d_n(O+)} & \text{for even } 0 \leq n \leq l \\ s_0 b_{(O-)s_0 d_n(O+)} & \text{for odd } 1 \leq n \leq l. \end{cases} \quad (4.12)$$

In fact, viewing $V(l)$ as a $\mathbb{Z}[q, q^{-1}]$ -algebra by identifying it with the quotient U_t^l/\mathbb{I}_l as explained in Remark 2.13, ϕ is an algebra isomorphism. Also, $\phi(bv) = b \circ \phi(v)$, $\phi(\bar{v}) = \overline{\phi(v)}$, and $(v, w) = q^{\binom{l}{2}} (\overline{\phi(v)}, \overline{\phi(w)})$ for all $v, w \in V(l)$.

Proof. We define ϕ by (4.11). First we claim that $\phi(bv) = b\phi(v)$ for any $v \in V(l)$. The claim implies that ϕ is an algebra isomorphism since $\mathbb{Q}(q) \otimes_{\mathbb{Z}[q, q^{-1}]} U_t^l$ is generated by b . To prove it, we may assume that $v = f^{(n)} \eta_l$ for some $0 \leq n \leq l$. We have that $b_{O-, O+} = b_{O-, O} \circ b_{O, O+}$ and $b_{O+, O-} = b_{O+, O} \circ b_{O, O-}$. If n is even then, in view also of (2.34), the proof reduces to checking that

$$b_{O-, O} \circ b_{O, O+} \circ h_{(O+)d_n(O+)} = [n+1]_q h_{(O-)s_0 d_{n+1}(O+)} + q^{l-2n+1} [l-n+1]_q h_{(O-)s_0 d_{n-1}(O+)},$$

omitting the first term if $n = l$ and the last term if $n = 0$. If n is odd, we need to show instead that

$$b_{O+, O} \circ b_{O, O-} \circ h_{(O-)s_0 d_n(O+)} = [n+1]_q h_{(O+)d_{n+1}(O+)} + q^{l-2n+1} [l-n+1]_q h_{(O+)d_{n-1}(O+)},$$

omitting the first term if $n = l$. This follows in both cases by a computation using Lemma 3.7, (3.32), (4.3), and the information about double cosets in Lemmas 4.2 to 4.4. For example, if n is even and $0 < n < l$, then Lemma 3.7(1) implies that

$$b_{O, O+} \circ h_{(O+)d_n(O+)} = q^{l-n} h_{O d_n(O+)} + h_{O s_0 d_{n+1}(O+)},$$

there being two terms in the summation ($d' = d_n$ and $d' = d_{n+1}$). Then we use Lemma 3.7(2) to get

$$b_{O-, O} \circ b_{O, O+} \circ h_{(O+)d_n(O+)} = q^{1-n} [l-n+1]_q q^{l-n} h_{(O-)s_0 d_{n-1}(O+)} + [n+1]_q h_{(O-)s_0 d_{n+1}(O+)},$$

as required. The other cases follow by similar calculations.

Next we check that $\phi(\bar{v}) = \overline{\phi(v)}$. Since $V(l)$ is generated as a U_l -module by the highest weight vector η_l , and the elements denoted b on both sides of the picture are bar-invariant, the proof of this reduces using the claim established in the previous paragraph just to checking that $\phi(\overline{\eta_l}) = \overline{\phi(\eta_l)}$. This is clear since η_l and $\phi(\eta_l) = 1_{O_+}$ are both bar-invariant.

Now we can show that (4.12) holds. Applying the bar involution to (2.35) shows that $b^{(n)}\eta_l$ is bar-invariant and equals $\overline{f^{(n)}\eta_l}$ (a $q\mathbb{Z}[q]$ -linear combination of other $\overline{f^{(i)}\eta_l}$). We deduce from (4.11) and (4.12) that $\phi(b^{(n)}\eta_l)$ is bar-invariant and equals $h_{(O_+)d_n(O_+)} + (*)$ if n is even or $s_0h_{(O_-)s_0d_n(O_+)} + (*)$ if n is odd, where $(*)$ is a $q\mathbb{Z}[q]$ -linear combination of other standard basis elements. It follows that $\phi(b^{(n)}\eta_l)$ equals $b_{(O_+)d_n(O_+)}$ if n is even or $s_0b_{(O_-)s_0d_n(O_+)}$ if n is odd, since these properties characterize this Kazhdan-Lusztig basis.

Finally, we check that $(v, w) = q^{\binom{l}{2}}(\overline{\phi(v)}, \overline{\phi(w)})$. We may assume that $v = f^{(m)}\eta_l$ and $w = f^{(n)}\eta_l$ for $0 \leq m, n \leq l$. By the orthogonality of the respective standard bases, both sides are 0 if $m \neq n$, so assume that $m = n$. Using (2.33), we are reduced to showing that

$$q^{-n(l-n)} \begin{bmatrix} l \\ n \end{bmatrix}_q = \begin{cases} q^{\binom{l}{2}} \overline{(h_{(O_+)d_n(O_+)}, h_{(O_+)d_n(O_+)})} & \text{if } n \text{ is even} \\ q^{\binom{l}{2}} \overline{(h_{(O_-)s_0d_n(O_+)}, h_{(O_-)s_0d_n(O_+)})} & \text{if } n \text{ is odd.} \end{cases}$$

This follows from (3.39) and (4.3), using also the descriptions of stabilizers in Lemmas 4.3 and 4.4. \square

The following corollaries follow from the theorem using (2.21), Corollary 2.15, and Lemma 2.16, respectively.

Corollary 4.6. *For any $n \geq 0$, we have in $S \oplus s_0S$ that*

$$b^{\circ n} = \begin{cases} \sum_{\substack{0 \leq i \leq \frac{n}{2} \\ n-2i \leq l}} [n-2i]_q! \left(\sum_{\lambda \in \text{Par}_{\neq i}(i \times (n-2i))} [\lambda_1 + 1]_q^2 \cdots [\lambda_i + 1]_q^2 \right) b_{(O_+)d_{n-2i}(O_+)} & \text{if } n \text{ is even} \\ \sum_{\substack{0 \leq i \leq \frac{n-1}{2} \\ n-2i \leq l}} [n-2i]_q! \left(\sum_{\lambda \in \text{Par}_{\neq i}(i \times (n-2i))} [\lambda_1 + 1]_q^2 \cdots [\lambda_i + 1]_q^2 \right) s_0 b_{(O_-)s_0d_{n-2i}(O_+)} & \text{if } n \text{ is odd.} \end{cases}$$

Corollary 4.7. *For $0 \leq n \leq \lfloor \frac{l}{2} \rfloor$, we have that $q^{\binom{l}{2}} \overline{(1_{O_+}, b_{(O_+)d_{2n}(O_+)})} = q^{-n(l+n-1)} \left[\begin{smallmatrix} l-t \\ n \end{smallmatrix} \right]_{q^2}$.*

Corollary 4.8. *The top degree term of the Laurent polynomial $q^{\binom{l}{2}} \overline{(b^{\circ n}, b^{\circ n})}$ is $q^{2\binom{n}{2}}$ when $0 \leq n \leq l$, and it is $q^{(2n-l)(l-1)}$ when $n > l$.*

4.5. Extended singular Soergel bimodules. Let $\mathfrak{B} \ni \mathfrak{B} \text{im}$ be the graded bicategory of singular Bott-Samelson bimodules from Definition 3.3 for the realization of D_l fixed in Subsection 4.1. We remind again that $\gamma : N \rightarrow N$ denotes the graph automorphism.

Definition 4.9. For $I \subseteq N$, we write s_0R^I for the graded $(R^{\gamma(I)}, R^I)$ -bimodule that is a copy $\{s_0f \mid f \in R^I\}$ of the graded vector space R^I (here s_0 is a formal symbol). It is equipped with the natural right action of R^I , and a twisted left action of $R^{\gamma(I)}$ defined by $x \cdot (s_0f) := s_0\gamma(x)f$. Similarly, there is a graded $(R^I, R^{\gamma(I)})$ -bimodule $R^I s_0 = \{f s_0 \mid f \in R^I\}$ with the natural left action and the twisted right action.

The functor $s_0R^I \otimes_{R^I} -$ (resp., $- \otimes_{R^I} R^I s_0$) amounts to twisting the left (resp., right) action of R^I into an action of $R^{\gamma(I)}$ by pull-back along the isomorphism $\gamma : R^{\gamma(I)} \xrightarrow{\sim} R^I$. We refer to s_0R^I and $R^I s_0$ as *twisting bimodules*. There is an obvious isomorphism of graded $(R^I, R^{\gamma(I)})$ -bimodules $s_0R^{\gamma(I)} \xrightarrow{\sim} R^I s_0$

which sends $s_0 f \mapsto \gamma(f)s_0$. Because of this isomorphism we avoid using $R^I s_0$ entirely in the definitions below, having chosen to prefer $s_0 R^{\gamma(I)}$.

Definition 4.10. Let $e\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im}$, the graded bicategory of *extended singular Bott-Samelson bimodules* (of type D_I), be the full sub-bicategory of $\mathfrak{g}\mathfrak{B}\mathfrak{im}$ with the same objects R^I ($I \subseteq N$) as $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im}$, but with 1-morphisms given by tensor products of the bimodules $B_{I,K}^J$ for all $I \supseteq K \subseteq J$ as before plus the new twisting bimodules $s_0 R^I$ for $I \subseteq N$. The graded bicategory of *extended singular Soergel bimodules* is its graded Karoubi envelope $\mathfrak{g}\mathfrak{Kar}(e\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im})$, which we identify with a full sub-bicategory of $\mathfrak{g}\mathfrak{B}\mathfrak{im}$ in the natural way.

In string diagrams, we use a *dashed string* $\gamma(I) \mid I$ separating 2-cells labelled I to the right and $\gamma(I)$ to the left to denote the identity endomorphism of the twisting bimodule $s_0 R^I$. In contrast, our previous diagrams were built from *undashed strings*. We have the obvious bubble slides:

$$\gamma(I) \mid \textcircled{f} I = \gamma(I) \mid \textcircled{\gamma(f)} I \quad (4.13)$$

for $f \in R^I$ (so $\gamma(f) \in R^{\gamma(I)}$). Left-tensoring with the twisting bimodule $s_0 R^I$ is obviously an equivalence of categories with quasi-inverse given by left-tensoring with $s_0 R^{\gamma(I)}$. One of the resulting adjunctions defines degree 0 graded bimodule isomorphisms $s_0 R^{\gamma(I)} \otimes_{R^{\gamma(I)}} s_0 R^I \xrightarrow{\sim} R^I$ and $R^I \xrightarrow{\sim} s_0 R^I \otimes_{R^I} s_0 R^{\gamma(I)}$ represented in string diagrams by the dashed caps and cups

$$\begin{aligned} \textcircled{\gamma(I)} I &: s_0 R^{\gamma(I)} \otimes_{R^{\gamma(I)}} s_0 R^I \rightarrow R^I, & f \otimes g &\mapsto \gamma(f)g, \\ I \textcircled{\gamma(I)} &: R^{\gamma(I)} \rightarrow s_0 R^I \otimes_{R^I} s_0 R^{\gamma(I)}, & 1 &\mapsto 1 \otimes 1. \end{aligned}$$

These satisfy the zig-zag identities, in addition to being mutually inverse isomorphisms:

$$I \textcircled{\gamma(I)} \textcircled{\gamma(I)} I = I \textcircled{\gamma(I)} I \textcircled{\gamma(I)} I = I \textcircled{\gamma(I)} I \textcircled{\gamma(I)} I, \quad \textcircled{\gamma(I)} I = \gamma(I) \mid I \mid \gamma(I), \quad \textcircled{\gamma(I)} I = \text{id}_{R^I}. \quad (4.14)$$

Using these relations, one can argue that any two diagrams built entirely from dashed strings and their cups and caps, and having the same boundary (thus representing morphisms between the same tensor products of twisting bimodules), are actually equal. (Formally, this augmented string calculus is encoding a categorical action of the cyclic group C_2 .)

More interestingly there are *mixed crossings*, that is, crossings of dashed strings with undashed strings. These represent the following graded bimodule homomorphisms of degree 0:

$$\begin{aligned} \begin{array}{c} I \\ \swarrow \gamma(I) \\ \searrow \gamma(Ii) \\ Ii \end{array} &: B_{\gamma(I),\gamma(Ii)} \otimes_{R^{\gamma(Ii)}} s_0 R^{Ii} \rightarrow s_0 R^I \otimes_{R^I} B_{I,Ii}, & f \otimes 1 &\mapsto 1 \otimes \gamma(f), \\ \begin{array}{c} \swarrow \gamma(Ii) \\ \searrow \gamma(I) \\ I \end{array} &: s_0 R^I \otimes_{R^I} B_{I,Ii} \rightarrow B_{\gamma(I),\gamma(Ii)} \otimes_{R^{\gamma(Ii)}} s_0 R^{Ii}, & 1 \otimes f &\mapsto \gamma(f) \otimes 1, \\ \begin{array}{c} Ii \\ \swarrow \gamma(Ii) \\ \searrow \gamma(I) \\ I \end{array} &: B_{\gamma(Ii),\gamma(I)} \otimes_{R^{\gamma(I)}} s_0 R^I \rightarrow s_0 R^{Ii} \otimes_{R^{Ii}} B_{Ii,I}, & f \otimes 1 &\mapsto 1 \otimes \gamma(f), \\ \begin{array}{c} \swarrow \gamma(I) \\ \searrow \gamma(Ii) \\ Ii \end{array} &: s_0 R^{Ii} \otimes_{R^{Ii}} B_{Ii,I} \rightarrow B_{\gamma(Ii),\gamma(I)} \otimes_{R^{\gamma(I)}} s_0 R^I, & 1 \otimes f &\mapsto \gamma(f) \otimes 1. \end{aligned}$$

Although we are not labelling strings with their colors, in all four of the above diagrams, the undashed strings change color from i to $\gamma(i)$ at the point that they cross the dashed string.

The mixed crossings are mutual inverses, that is, they satisfy Reidemeister II, in the obvious ways:

$$\begin{array}{c} \begin{array}{c} \gamma(I) \\ \nearrow \\ I \\ \searrow \\ \gamma(I) \end{array} Ii = \begin{array}{c} \uparrow \\ \gamma(I) \end{array} \begin{array}{c} | \\ \gamma(I) \end{array} \begin{array}{c} | \\ Ii \end{array}, \end{array} \quad \begin{array}{c} \begin{array}{c} I \\ \nearrow \\ I \\ \searrow \\ I \end{array} Ii = \begin{array}{c} | \\ \gamma(I) \end{array} \begin{array}{c} | \\ Ii \end{array} \begin{array}{c} \uparrow \\ \end{array}, \end{array} \quad (4.15)$$

$$\begin{array}{c} \begin{array}{c} \gamma(I) \\ \searrow \\ Ii \\ \nearrow \\ \gamma(I) \end{array} I = \begin{array}{c} \downarrow \\ \gamma(I) \end{array} \begin{array}{c} | \\ \gamma(I) \end{array} \begin{array}{c} | \\ I \end{array}, \end{array} \quad \begin{array}{c} \begin{array}{c} Ii \\ \nearrow \\ \gamma(I) \\ \searrow \\ Ii \end{array} I = \begin{array}{c} | \\ \gamma(I) \end{array} \begin{array}{c} | \\ Ii \end{array} \begin{array}{c} \downarrow \\ \end{array}. \end{array} \quad (4.16)$$

There are also Reidemeister III relations for diagrams involving dashed strings. These are obvious and we will not spell them out further. Consequently, when at least one dashed string is involved, we can even draw triple intersections without any ambiguity.

The upshot of this discussion is that dashed strings are easy to manipulate graphically—they slide across all other features, applying the graph automorphism to bubbles in the process, and they can be cut open and restitched in any reasonable way.

The graded 2-functor \mathbb{R} from (3.26) extends to a graded 2-functor

$$\mathbb{R} : \mathfrak{eB} \mathfrak{S} \mathfrak{Bim} \rightarrow \mathfrak{eB} \mathfrak{S} \mathfrak{Bim}^{\text{rev}} \quad (4.17)$$

This fixes objects, takes the graded (R^I, R^J) -bimodule M to $q^{\ell(w_J) - \ell(w_I)} M$ viewed as an (R^J, R^I) -bimodule, and takes the twisting bimodule $s_0 R^I$ to $s_0 R^{\gamma(I)}$. To define \mathbb{R} on a bimodule homomorphism θ , the same underlying function defines a bimodule homomorphism between the bimodules obtained instead by replacing each $s_0 R^I$ with $R^I s_0$; one then needs to conjugate with the appropriate isomorphisms $s_0 R^{\gamma(I)} \xrightarrow{\sim} R^I s_0$ to obtain from this the bimodule homomorphism $\mathbb{R}(\theta)$. In terms of string diagrams, the functor \mathbb{R} reflects diagrams in a vertical axis and reverses the orientation on all undashed strings as before; dotted bubbles of the form \textcircled{f} in 2-cells are unchanged. (The duality \mathbb{D} from (3.45) obviously extends too, but we will not use this below.)

The Soergel-Williamson Theorems 3.10 to 3.12 are easily adapted to the extended setting. Although we do not believe that extended singular Soergel bimodules are yet in the literature, one can find a study of extended (ordinary) Soergel bimodules in [Eli, Ch.3, Appendix], and it is straightforward to adapt the proofs to our setting. We just make a few comments.

Any graded bimodule that is a 1-morphism in $\mathfrak{eB} \mathfrak{S} \mathfrak{Bim}$ is a tensor product of singular Bott-Samelson bimodules and some number of twisting bimodules. We say that the bimodule is *untwisted* or *twisted* according to whether the number of twisting bimodules in the tensor product is even or odd. These two types of bimodule can easily be distinguished since left and right multiplication by $e_i(x_1, \dots, x_l) = x_1 \cdots x_l \in R^N$ are equal on untwisted bimodules, and they differ by a sign on twisted ones. In particular, no non-zero untwisted bimodule is isomorphic to a twisted one.

Mixed crossings will give an isomorphism between any object in $\mathfrak{eB} \mathfrak{S} \mathfrak{Bim}$ and a bimodule of the form $M \otimes X$, where M is a tensor product of twisting bimodules and X is a singular Bott-Samelson bimodule. Dashed cups and caps give additional isomorphisms, letting one assume that M is either the identity bimodule (if untwisted) or a single twisting bimodule (if twisted). Finally, we claim that tensoring with id_M gives an isomorphism

$$\text{Hom}_{\mathfrak{eB} \mathfrak{S} \mathfrak{Bim}}(X, X') \cong \text{Hom}_{\mathfrak{eB} \mathfrak{S} \mathfrak{Bim}}(M \otimes X, M \otimes X'). \quad (4.18)$$

A priori, the right-hand side also contains diagrams with dashed strings appearing willy-nilly, but the relations allow one to slide all dashed strings to the left and then pull them straight.

The following is easily deduced from this discussion:

- The classification of indecomposable untwisted bimodules in $e\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im}$ is exactly the same as in $\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im}$, as described by Theorem 3.11. The classification of indecomposable twisted bimodules is almost the same, except one needs to twist once so that, up to isomorphism and grading shift, the indecomposable twisted (R^I, R^J) -bimodules in $\mathfrak{g}\mathfrak{K}\mathfrak{ar}(e\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im})$ are the bimodules $s_0R^I \otimes_{R^I} B_{IdJ}$ for $I, J \subseteq N$ and $d \in (W_I \backslash W / W_J)_{\min}$.
- There are no non-zero homomorphisms between a twisted and an untwisted bimodule. Given two untwisted bimodules, the homomorphism formula is the same as before. Given twisted bimodules, there is an analogous formula, but one needs to allow Δ - and ∇ -flags which involve the twisted standard and costandard bimodules $s_0R^I \otimes_{R^I} \Delta_{IdJ}$ and $s_0R^I \otimes_{R^I} \nabla_{IdJ}$, leading to a slightly modified definitions of the character maps ch_Δ and ch_∇ . (Alternatively, one can just tensor one more time with a twisting bimodule to reduce to the usual homomorphism formula in the untwisted setting.)
- Finally, the categorification theorem is easily adapted. The modified character map ch takes the indecomposable untwisted bimodule B_{IdJ} to $b_{IdJ} \in \mathfrak{S}$ as before, and it takes the twisted bimodule $s_0R^I \otimes_{R^I} B_{IdJ}$ to s_0b_{IdJ} . One recovers the extended q -Schur algebra:

$$K_0(\mathfrak{g}\mathfrak{K}\mathfrak{ar}(e\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im})) \cong \mathfrak{S} \oplus s_0\mathfrak{S}, \quad (4.19)$$

4.6. Diagrammatics for extended singular Bott-Samelson bimodules. We now start to use the diagrammatic calculus explained in Subsection 3.3 more seriously for making calculations in the graded bicategory $e\mathfrak{B}\mathfrak{S}\mathfrak{B}\mathfrak{im}$ from Definition 4.10. Traditionally in this subject an unlabelled 2-cell in a string diagram is used to denote the 2-cell labelled by the empty set, but from now on:

An unlabelled 2-cell in a string diagram denotes the 2-cell labelled by O .

Thus O is our “ground state”. Then we decorate the 2-cell with $+$, $-$ or \pm to denote the 2-cell labelled by $O+$, $O-$ or $O\pm$, and we decorate it with \hat{n} , $\hat{n}+$, $\hat{n}-$ or $\hat{n}\pm$ to denote the 2-cell labelled by $O\hat{n}$, $O\hat{n}+$, $O\hat{n}-$ or $O\hat{n}\pm$ for $n \in O$, using the shorthands (4.2). In fact, moving forwards, we will rarely need 2-cells with labels of the form $I-$ for $I \subseteq O$; it will be enough to consider string diagrams with 2-cells labelled I , $I+$ or $I\pm$ for $I \subseteq O$. Typically I will be O or $O\hat{n}$.

Now we introduce an important notational shorthand to the string calculus for extended singular Soergel bimodules. Suppose that $I \subseteq O$, so that $\gamma(I) = I$. We will use the *special undotted strings*

$$\begin{array}{c} \uparrow \\ I \\ \uparrow \end{array} I_+ := I \uparrow I_+, \quad \begin{array}{c} \uparrow \\ I_+ \\ \uparrow \end{array} I_\pm := I_+ \uparrow I_\pm, \quad \begin{array}{c} I_\pm \\ \downarrow \\ I_+ \end{array} := I_\pm \downarrow I_+, \quad \begin{array}{c} I_+ \\ \downarrow \\ I \end{array} := I_+ \downarrow I \quad (4.20)$$

to denote the identity endomorphisms of the untwisted bimodules B_{I,I_+} , B_{I_+,I_\pm} , B_{I_\pm,I_+} and $B_{I_+,I}$. We use the *special dotted strings*

$$\begin{array}{c} \uparrow \\ I \\ \uparrow \end{array} I_+ := I \uparrow I_+, \quad \begin{array}{c} \uparrow \\ I_+ \\ \uparrow \end{array} I_\pm := I_+ \uparrow I_\pm, \quad \begin{array}{c} I_\pm \\ \downarrow \\ I_+ \end{array} := I_\pm \downarrow I_+, \quad \begin{array}{c} I_+ \\ \downarrow \\ I \end{array} := I_+ \downarrow I \quad (4.21)$$

to denote the identity endomorphisms of the twisted bimodules $s_0R^I \otimes_{R^I} B_{I,I_+}$, $B_{I_+,I_\pm} \otimes_{R^I} s_0R^{I\pm}$, $s_0R^{I\pm} \otimes_{R^I} B_{I_\pm,I_+}$ and $s_0R^{I+} \otimes_{R^I} B_{I_+,I}$. It is helpful to remember that the dashed line, when it appears in these diagrams, passes through the 2-cells whose label is invariant under γ . We reiterate that these special dotted strings are not identity morphisms of new objects in the category, they are merely a shorthand for the identity morphisms of existing objects. In this shorthand, the equality (4.22) is tautological, and (4.25) is a consequence of (4.13). Actually, the special undotted string is an ordinary

string labelled either $+$ or $-$, but we find the addition of color to it makes the diagrams in which it arises later on more readable.

Notice that $I+\downarrow I\uparrow I+ = I+\downarrow I\uparrow I\uparrow I+ = I+\downarrow I\uparrow I+$, indeed, these represent the identity endomorphism of the twisted (R^{I+}, R^{I+}) -bimodule $B_{I+,I} \otimes_{R^I} s_0 R^I \otimes_{R^I} B_{I,I+}$. Most important is the case $I = O$ here, when

$$+\downarrow\uparrow+ = +\downarrow\uparrow+ = +\downarrow\uparrow+ \quad (4.22)$$

is the identity endomorphism of the twisted (R^{O+}, R^{O+}) -bimodule

$$B := B_{O+,O} \otimes_{R^O} s_0 R^O \otimes_{R^O} B_{O,O+}. \quad (4.23)$$

This is isomorphic via an obvious contraction to the bimodule B defined in the introduction. We will use the present definition of B from now on. At the level of characters, B corresponds to the element $b \in s_0 S$ from (4.9):

$$\text{ch}(B) = b_{O+,O} \circ (s_0 b_{O,O+}) = s_0 b_{O-,O} \circ b_{O,O+} = s_0 b_{O-,O+} = b. \quad (4.24)$$

Note also that

$$+\downarrow\uparrow+ = +\downarrow\uparrow+ . \quad (4.25)$$

The symmetry R from (4.17) acts on string diagrams as before, preserving the type (undotted or dotted) of special strings. For example:

$$R \left(+\downarrow\uparrow+ \right) = +\downarrow\uparrow+ , \quad R \left(+\downarrow\uparrow+ \right) = +\downarrow\uparrow+ . \quad (4.26)$$

There are several more diagrammatic shorthands to be introduced involving the two new types of colored strings. These should all seem unsurprising by this point. Continuing with any $I \subseteq O$, there are colored caps and cups satisfying zig-zag identities:

$$\begin{array}{cccccccc} \text{I}^+ \downarrow \text{I} & \text{I} \uparrow \text{I}^+ & \text{I} \downarrow \text{I}^+ & \text{I}^+ \uparrow \text{I} & \text{I}^\pm \downarrow \text{I}^+ & \text{I}^+ \uparrow \text{I}^\pm & \text{I}^+ \downarrow \text{I}^\pm & \text{I}^\pm \uparrow \text{I}^+ \\ \text{I}^+ \downarrow \text{I} & \text{I} \uparrow \text{I}^+ & \text{I} \downarrow \text{I}^+ & \text{I}^+ \uparrow \text{I} & \text{I}^\pm \downarrow \text{I}^+ & \text{I}^+ \uparrow \text{I}^\pm & \text{I}^+ \downarrow \text{I}^\pm & \text{I}^\pm \uparrow \text{I}^+ \end{array}$$

For example, the last two are defined by

$$\text{I}^+ \downarrow \text{I}^\pm := \text{I}^+ \downarrow \text{I}^\pm \text{I}^\pm , \quad \text{I}^\pm \uparrow \text{I}^+ := \text{I}^\pm \uparrow \text{I}^\pm \text{I}^+ ,$$

and they satisfy the zig-zag identities by combining the ones for ordinary and dashed strings seen before. The other cases are similar.

Now we discuss crossings. Undotted special strings are ordinary strings (with added color for clarity), so we have already defined crossings between them and other ordinary strings. There are also crossings of dotted special strings with others strings which we call *special crossings*. For $i \in I \subseteq O$, the upward special crossings are defined as follows:

$$\begin{array}{cc} \begin{array}{c} \text{I}^+ \text{I}^+ \\ \text{I} \text{I}^+ \\ \text{I}^+ \text{I}^+ \end{array} := \begin{array}{c} \text{I} \text{I}^+ \\ \text{I} \text{I}^+ \\ \text{I}^+ \text{I}^+ \end{array} , & \begin{array}{c} \text{I}^+ \text{I}^+ \\ \text{I} \text{I}^+ \\ \text{I}^+ \text{I}^+ \end{array} := \begin{array}{c} \text{I} \text{I}^+ \\ \text{I} \text{I}^+ \\ \text{I}^+ \text{I}^+ \end{array} , \\ \begin{array}{c} \text{I}^+ \text{I}^\pm \\ \text{I} \text{I}^\pm \\ \text{I}^+ \text{I}^\pm \end{array} := \begin{array}{c} \text{I} \text{I}^\pm \\ \text{I} \text{I}^\pm \\ \text{I}^+ \text{I}^\pm \end{array} , & \begin{array}{c} \text{I}^+ \text{I}^\pm \\ \text{I} \text{I}^\pm \\ \text{I}^+ \text{I}^\pm \end{array} := \begin{array}{c} \text{I} \text{I}^\pm \\ \text{I} \text{I}^\pm \\ \text{I}^+ \text{I}^\pm \end{array} , \end{array}$$

referring to the discussion after (4.16) for interpretation of the triple crossings. One defines downward and sideways special crossings by rotating the ones above, noting the cyclicity of the string calculus.

The special crossings satisfy the easy Reidemeister II relations, as follows directly from (3.21) and the definitions, using the simple behavior of dashed strings. Other relations involving the new colored strings can be worked out in just the same way. Here are a few more examples, mainly to draw attention to the importance of the type of special strings (i.e., undotted vs. dotted):

$$\begin{array}{c} \uparrow \\ \textcircled{f} \end{array} + = \textcircled{f} \uparrow +, \quad \begin{array}{c} \uparrow \\ \textcircled{f} \end{array} + = \textcircled{\gamma(f)} \uparrow + \quad \text{for } f \in R^{O+}, \quad (4.27)$$

$$\begin{array}{c} \textcircled{f} \\ \downarrow \end{array} + = \textcircled{\text{tr}_{O+}^O(f)} +, \quad \begin{array}{c} \textcircled{f} \\ \downarrow \end{array} + = \textcircled{\text{tr}_{O+}^O(\gamma(f))} + = \textcircled{\gamma(\text{tr}_{O-}^O(f))} + \quad \text{for } f \in R^O, \quad (4.28)$$

$$\begin{array}{c} \textcircled{+} \\ \downarrow \end{array} = \textcircled{\eta_{O+}^O}, \quad \begin{array}{c} \textcircled{+} \\ \downarrow \end{array} = \textcircled{\gamma(\eta_{O+}^O)} = \textcircled{\eta_{O-}^O}. \quad (4.29)$$

This calculus will be used extensively in the rest of the paper.

Remark 4.11. Since thinking in a foreign language is hard, we sometimes find it easier to replace the special strings with ordinary and dashed strings according to (4.20) and (4.21), slide the resulting dashed strings to the edges, then work with the ordinary (i.e., non-extended) string calculus after that. A roughly equivalent technique is to use the following mutually inverse isomorphisms to relate $B^{\otimes 2}$ with the untwisted singular Soergel bimodule $B_{O+,O-} \otimes_{R^{O-}} B_{O-,O+}$:

$$\begin{array}{c} \uparrow \\ \textcircled{+} \\ \downarrow \end{array} : B^{\otimes 2} \xrightarrow{\sim} B_{O+,O-} \otimes_{R^{O-}} B_{O-,O+}, \quad \begin{array}{c} \downarrow \\ \textcircled{+} \\ \uparrow \end{array} : B_{O+,O-} \otimes_{R^{O-}} B_{O-,O+} \xrightarrow{\sim} B^{\otimes 2}. \quad (4.30)$$

There are similar isomorphisms removing extra twisting bimodules in tensor products $B^{\otimes n}$ for all larger values of n , e.g., $B^{\otimes 3} \cong s_0 R_{O-} \otimes_{R^{O-}} B_{O-,O+} \otimes_{R^{O+}} B_{O+,O-} \otimes_{R^{O-}} B_{O-,O+}$. Using these, relations involving extended string calculus become equivalent to relations involving ordinary string calculus. For an example, see Remark 4.17.

4.7. Calculations. Many of our interesting calculations require some additional understanding of the underlying Frobenius extensions. As a first example, we look again at the first equality in (4.29). The circle $\textcircled{+}$ is, by definition, equal to multiplication by the product-coproduct element for the Frobenius extension $R^{O+} \hookrightarrow R^O$. For practice, let us re-check that this product-coproduct element is η_{O+}^O , as we already known by the general theory. The longest element of $(W_{O+}/W_O)_{\min}$ is $s_{l-1} \cdots s_2 s_1$, so the trace of the Frobenius extension $R^{O+} \hookrightarrow R^O$ is $\partial_{l-1} \cdots \partial_2 \partial_1$. For $i = 1, \dots, l-1$, $\partial_i(f) = \frac{f - s_i(f)}{x_{i+1} - x_i}$ by our choice of simple roots in Subsection 4.1. The reader can verify (a standard exercise in Schubert calculus) that an explicit pair of dual bases for R^O as a free R^{O+} -module are given by $((-1)^r x_1^r)_{0 \leq r \leq l-1}$ and $(e_{l-1-r}(x_2, \dots, x_l))_{0 \leq r \leq l-1}$. Consequently $\textcircled{+}$ is multiplication by

$$\sum_{r=0}^{l-1} (-1)^r x_1^r e_{l-1-r}(x_2, \dots, x_l) = (x_2 - x_1) \cdots (x_l - x_1) = \eta_{O+}^O.$$

Now recall that $A = R^{O+} = \mathbb{k}[x_1, \dots, x_l]^{S^l}$. The elementary symmetric polynomial $e_r(x_1, \dots, x_l)$ is the image of the elementary symmetric function $e_r \in \Lambda$ under the the natural evaluation map $\text{ev}_l : \Lambda \rightarrow A$ from (2.5). We also let $h_r(x_1, \dots, x_l)$ and $q_r(x_1, \dots, x_l)$ be the evaluations of the r th complete symmetric function and the q -function q_r from (2.2), respectively.

Lemma 4.12. *For $r \geq 0$, we have that $\partial_{l-1} \cdots \partial_2 \partial_1((-x_1)^r) = (-1)^{r-l+1} h_{r-l+1}(x_1, \dots, x_l)$ (interpreted as 0 in case $r < l-1$).*

Proof. Given that $x_1^r = h_r(x_1)$, this follows from the well-known fact that

$$\partial_i h_k(x_1, \dots, x_i) = -h_{k-1}(x_1, \dots, x_{i+1}).$$

□

Lemma 4.13. *We have that*

$$\mathrm{tr}_{O_+}^O(x_1^n \eta_{O_-}^O) = \begin{cases} \frac{(-1)^{l-1}+1}{2} \mathrm{id}_A & \text{if } n = 0 \\ -(-1)^l \frac{1}{2} q_n(x_1, \dots, x_l) \mathrm{id}_A & \text{if } n > 0. \end{cases}$$

(The number $\frac{(-1)^{l-1}+1}{2}$ is the element of $\{0, 1\}$ which is equal to $l \pmod 2$. Later, this will match the parameter t in the nilBrauer category \mathbf{NB}_t .)

Proof. We use the identity $e_r(x_2, \dots, x_l) + x_1 e_{r-1}(x_2, \dots, x_l) = e_r(x_1, \dots, x_l)$, which is valid for all $r \geq 0$ if one interprets $e_{-1}(x_2, \dots, x_l)$ as 0, to see that

$$\begin{aligned} \mathrm{tr}_{O_+}^O(x_1^n \eta_{O_-}^O) &= \partial_{l-1} \cdots \partial_1 (x_1^n (x_2 + x_1) \cdots (x_l + x_1)) \\ &= \sum_{r=0}^{l-1} \partial_{l-1} \cdots \partial_1 (x_1^{n+l-1-r} e_r(x_2, \dots, x_l)) \\ &= \frac{1}{2} \sum_{r=0}^{l-1} \partial_{l-1} \cdots \partial_1 (x_1^{n+l-1-r} e_r(x_2, \dots, x_l)) + \frac{1}{2} \sum_{r=1}^l \partial_{l-1} \cdots \partial_1 (x_1^{n+l-r} e_{r-1}(x_2, \dots, x_l)) \\ &= \frac{1}{2} \sum_{r=0}^{l-1} \partial_{l-1} \cdots \partial_1 (x_1^{n+l-1-r} (e_r(x_2, \dots, x_l) + x_1 e_{r-1}(x_2, \dots, x_l))) + \\ &\qquad\qquad\qquad \frac{1}{2} \partial_{l-1} \cdots \partial_1 (x_1^n e_{l-1}(x_2, \dots, x_l)) \\ &= \frac{1}{2} \sum_{r=0}^{l-1} \partial_{l-1} \cdots \partial_1 (x_1^{n+l-1-r} e_r(x_1, \dots, x_l)) + \frac{1}{2} \partial_{l-1} \cdots \partial_1 (x_1^n e_{l-1}(x_2, \dots, x_l)) \\ &= \frac{1}{2} \sum_{r=0}^{l-1} e_r(x_1, \dots, x_l) \partial_{l-1} \cdots \partial_1 (x_1^{n+l-1-r}) + \frac{1}{2} \partial_{l-1} \cdots \partial_1 (x_1^n e_{l-1}(x_2, \dots, x_l)). \end{aligned}$$

Now we use Lemma 4.12 to simplify further. When $n = 0$, using also the easy identity

$$\partial_{l-1} \cdots \partial_1 (x_2 x_3 \cdots x_l) = 1, \tag{4.31}$$

it reduces to $\frac{(-1)^{l-1}+1}{2}$. When $n > 0$, we obviously have that $x_1^n e_{l-1}(x_2, \dots, x_l) = x_1^{n-1} e_l(x_1, \dots, x_l)$, and the expression simplifies to

$$\begin{aligned} \frac{1}{2} \sum_{r=0}^l e_r(x_1, \dots, x_l) \partial_{l-1} \cdots \partial_1 (x_1^{n+l-1-r}) &= (-1)^{l-1} \frac{1}{2} \sum_{r=0}^{\min(l,n)} e_r(x_1, \dots, x_l) h_{n-r}(x_1, \dots, x_l) \\ &= (-1)^{l-1} \frac{1}{2} \sum_{r=0}^n e_r(x_1, \dots, x_l) h_{n-r}(x_1, \dots, x_l) \\ &= -(-1)^l \frac{1}{2} q_n(x_1, \dots, x_l), \end{aligned}$$

as required. □

Corollary 4.14. *We have that*

$$\left(\begin{array}{c} \oplus \\ \otimes \\ \boxplus \end{array} \right) x_1^n + = \begin{cases} \frac{(-1)^{l-1}+1}{2} \mathrm{id}_A & \text{if } n = 0 \\ -(-1)^l \frac{1}{2} q_n(x_1, \dots, x_l) \mathrm{id}_A & \text{if } n > 0. \end{cases}$$

Lemma 4.15. *Assuming $l \geq 3$, we have that*

$$\begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \end{array} = \begin{array}{c} \uparrow \\ + \\ \downarrow \end{array}. \quad (4.32)$$

Proof. This is an instance of Lemma 3.4. One needs to check the switchback relation $\partial_{O_+}^O = \partial_O^{O_2^+} \circ \partial_{O_2^+}^O$, which follows because both sides equal $\partial_{l-1} \cdots \partial_1$. \square

Lemma 4.16. *Assuming $l \geq 3$, the following relations hold:*

$$\begin{array}{c} \downarrow \\ \uparrow \\ + \\ \downarrow \\ \uparrow \\ + \end{array} = \frac{1}{2} \left(\begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \\ \alpha_{-1} \end{array} + \begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \\ \alpha_{-1} \end{array} \right), \quad (4.33)$$

$$\begin{array}{c} \uparrow + \\ \downarrow + \end{array} + \begin{array}{c} \downarrow + \\ \uparrow + \end{array} = \frac{1}{2} \left(\begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \\ \alpha_1 \end{array} + \begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \\ \alpha_1 \end{array} \right). \quad (4.34)$$

There are analogous relations in the case $l = 2$: one just has to omit 2-colored circles and all labels $\hat{2}$.

Proof. Suppose first that $l \geq 3$. Using that $\eta_{O_2^\pm}^{O_2^\pm} = 1$ and taking the basis $\mathbb{B}_{O_2^\pm}^{O_2^\pm}$ to be $1, \alpha_{-1}$ with dual basis $\frac{1}{2}\alpha_{-1}, \frac{1}{2}$, we have that

$$\begin{array}{c} \downarrow \\ \uparrow \\ + \\ \downarrow \\ \uparrow \\ + \end{array} \stackrel{(4.32)}{=} \begin{array}{c} \downarrow \\ \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \end{array} + \begin{array}{c} \downarrow \\ \uparrow + \\ \hat{2}+ \quad \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \end{array} \stackrel{(3.21)}{=} \begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \\ \alpha_{-1} \end{array} + \begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \\ \alpha_{-1} \end{array} \stackrel{(3.24)}{=} \begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \\ \alpha_{-1} \end{array} + \begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \\ \alpha_{-1} \end{array} \stackrel{(3.20)}{=} \frac{1}{2} \left(\begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \\ \alpha_{-1} \end{array} + \begin{array}{c} \uparrow + \\ \hat{2} \quad \hat{2}+ \quad \hat{2} \\ \downarrow + \\ \alpha_{-1} \end{array} \right).$$

Now we slide the α_{-1} bubbles across the special undotted strings using (4.27). To deduce the second identity, recall the equality (4.22). By expanding the special undotted and dotted strings in terms of ordinary and dashed strings, then using the easy relations satisfied by dashed strings, the identity just

displayed can be written in an equivalent form as

$$\begin{array}{c} + \\ \downarrow \\ + \\ \uparrow \\ + \\ \downarrow \\ + \\ \uparrow \end{array} = \frac{1}{2} \left(\begin{array}{c} \uparrow + \\ \hat{2} \hat{2} + \hat{2} \\ \textcircled{\alpha_{-1}} \\ \hat{2} + \hat{2} \pm \hat{2} + \\ \hat{2} \hat{2} + \hat{2} \\ \downarrow + \end{array} + \begin{array}{c} \uparrow + \\ \hat{2} \hat{2} + \hat{2} \\ \textcircled{\alpha_{-1}} \\ \hat{2} + \hat{2} \pm \hat{2} + \\ \hat{2} \hat{2} + \hat{2} \\ \downarrow + \end{array} \right). \quad (4.35)$$

Now we rotate counterclockwise through 90° using the cyclic structure, then slide the α_{-1} bubbles across the special dotted strings, which flips them to α_1 bubbles.

This argument is easily modified to treat the case $l = 2$, when there is no color 2. One can skip the first line altogether since (3.24) can be applied directly (without needing to remove any simple reflections). \square

Remark 4.17. Pre- and post-composing (4.33) with the isomorphisms from Remark 4.11, we get the following equivalent relation in the ordinary string calculus:

$$\begin{array}{c} + \\ \downarrow \\ + \\ \uparrow \\ - \\ \downarrow \\ + \\ \uparrow \end{array} = \frac{1}{2} \left(\begin{array}{c} - \\ \hat{2} \hat{2} - \hat{2} \\ \textcircled{\alpha_1} \\ \hat{2} + \hat{2} \pm \hat{2} + \\ \hat{2} \hat{2} - \hat{2} \\ \downarrow - \end{array} + \begin{array}{c} - \\ \hat{2} \hat{2} - \hat{2} \\ \textcircled{\alpha_1} \\ \hat{2} + \hat{2} \pm \hat{2} + \\ \hat{2} \hat{2} - \hat{2} \\ \downarrow - \end{array} \right).$$

This illustrates the alternate approach mentioned in Subsection 4.2.

Now we consider the n th tensor power $B^{\otimes n}$ of the (A, A) -bimodule B (the tensor product being over $A = R^{O^+}$). Its character is the element $b^{\otimes n} \in \mathbf{V}$ from (4.10). We are going to investigate certain endomorphisms which will be shown later to be primitive homogeneous idempotents.

Lemma 4.18. *If $0 \leq n \leq l$ then the endomorphism algebra of $B^{\otimes n}$ is 1-dimensional in degree $-2\binom{n}{2}$ and 0 in all smaller degrees. If $n > l$ then this endomorphism algebra is 0 in all degrees $\leq -2\binom{n}{2}$.*

Proof. If n is even, this follows from the homomorphism formula (Theorem 3.10) and Corollary 4.8. If n is odd, it follows from the modified homomorphism formula discussed in Subsection 4.5 (or the original one can be applied after left-tensoring with s_0A). When $n > l$ the degree bound from the homomorphism formula is $(2n - l)(l - 1)$, so it remains to observe that $(2n - l)(l - 1) < 2\binom{n}{2}$. This may be seen by replacing n by $l + k$ for $k > 0$, then checking that

$$2\binom{l+k}{2} - (2(l+k) - l)(l-1) = (l+k)(l+k-1) - (l+2k)(l-1) = k(k+1) > 0.$$

\square

For $1 \leq n \leq l$, we define an endomorphism $\mathbf{u}_n \in \text{End}_{A-A}(B^{\otimes n})$ by setting

$$\mathbf{u}_n := \begin{cases} \begin{array}{c} \text{Diagram 1} \\ \text{if } n = 1 \end{array} & \text{if } n = 1 \\ \begin{array}{c} \text{Diagram 2} \\ \text{if } n \text{ is even} \end{array} & \text{if } n \text{ is even} \\ \begin{array}{c} \text{Diagram 3} \\ \text{if } n \neq 1 \text{ is odd,} \end{array} & \text{if } n \neq 1 \text{ is odd,} \end{cases} \quad (4.36)$$

where there are a total of $n \downarrow \uparrow$ -pairs of strings at the top and bottom, $(n-1)$ cups at the top, and $(n-1)$ caps at the bottom. We have that $\mathbf{u}_n = \check{\mathbf{u}}_n \circ \hat{\mathbf{u}}_n$ where $\check{\mathbf{u}}_n$ denotes the top half of the above diagram and $\hat{\mathbf{u}}_n$ is its bottom half.

Lemma 4.19. For $1 \leq n \leq l$, both of the endomorphisms $\check{\mathbf{u}}_n$ and $\hat{\mathbf{u}}_n$ are of degree $\binom{l}{2} - n(l-1)$. This equals $-\binom{n}{2}$ if $n = l-1$ or $n = l$.

Proof. It is clear that $\check{\mathbf{u}}_n$ and $\hat{\mathbf{u}}_n$ are of the same degree. To compute the degree of $\check{\mathbf{u}}_n$, we use (4.3) to see:

- Each of the $(n-1)$ cups is of degree $\ell(w_{O_+}) - \ell(w_{O_{\pm}}) = \ell(w_{O_-}) - \ell(w_{O_{\pm}}) = -\binom{l}{2}$.
- Each of the n crossings is of degree $\ell(w_{O_{\pm}}) + \ell(w_O) - \ell(w_{O_+}) - \ell(w_{O_-}) = \binom{l-1}{2}$.

This gives the total degree $-(n-1)\binom{l}{2} + n\binom{l-1}{2} = \binom{l}{2} - n(l-1)$. □

Let $\mathbf{v}_0 := \text{id}_A$, $\mathbf{v}_1 := \downarrow \uparrow$, and $\mathbf{v}_l := \mathbf{u}_l$. Then for $2 \leq n \leq l-1$, we add an n -colored circle to the middle of the diagram defining \mathbf{u}_n to obtain

$$\mathbf{v}_n := \begin{cases} \begin{array}{c} \text{Diagram 4} \\ \text{if } n \text{ is even} \end{array} & \text{if } n \text{ is even} \\ \begin{array}{c} \text{Diagram 5} \\ \text{if } n \text{ is odd.} \end{array} & \text{if } n \text{ is odd.} \end{cases} \quad (4.37)$$

We reiterate that there are a total of $n \downarrow \uparrow$ -pairs and $(n-1)$ cups and caps at the top and bottom of these diagrams, so \mathbf{v}_n is an endomorphism of $B^{\otimes n}$. We have that $\mathbf{v}_n = \check{\mathbf{v}}_n \circ \hat{\mathbf{v}}_n$ where $\check{\mathbf{v}}_n$ denotes the top half of the above diagram and $\hat{\mathbf{v}}_n$ is its bottom half.

Lemma 4.20. *For $0 \leq n \leq l$, both of the endomorphisms $\check{\mathbf{v}}_n$ and $\hat{\mathbf{v}}_n$ are of degree $-\binom{n}{2}$.*

Proof. It is clear that $\check{\mathbf{v}}_n$ and $\hat{\mathbf{v}}_n$ are of the same degree. They have degree 0 if $n = 0$ or $n = 1$, and the result follows from Lemma 4.19 when $n = l$. So now we assume that $2 \leq n \leq l-1$, and proceed to compute the degree of $\check{\mathbf{v}}_n$. Using (4.3) again, we see:

- Each of the $n-1$ special cups is of degree $\ell(w_{O\hat{n}+}) - \ell(w_{O\hat{n}\pm}) = \ell(w_{O\hat{n}-}) - \ell(w_{O\hat{n}\pm}) = -\binom{n}{2}$.
- The n crossings of special strings are of degree $\ell(w_{O\hat{n}\pm}) + \ell(w_{O\hat{n}}) - \ell(w_{O\hat{n}+}) - \ell(w_{O\hat{n}-}) = \binom{n-1}{2}$;
- The ordinary cap of color n is of degree $\ell(w_{O\hat{n}}) - \ell(w_O) = -(n-1)(l-n)$ if n is odd or $\ell(w_{O\hat{n}+}) - \ell(w_{O+}) = \ell(w_{O\hat{n}-}) - \ell(w_{O-}) = -n(l-n)$ if n is even.
- Each of the crossings involving the ordinary string and a special string has degree $\ell(w_{O+}) + \ell(w_{O\hat{n}}) - \ell(w_{O\hat{n}+}) - \ell(w_O) = \ell(w_{O-}) + \ell(w_{O\hat{n}}) - \ell(w_{O\hat{n}-}) - \ell(w_O) = l-n$, and there are $(n-1)$ such if n is odd or n such if n is even.

The total degree coming from the last two points is 0. We are left with $-(n-1)\binom{n}{2} + n\binom{n}{2} = -\binom{n}{2}$. \square

We need two more families of endomorphisms. For $n \geq 1$, let

$$\mathbf{w}_n := \begin{cases} \begin{array}{c} \downarrow + \quad \curvearrowright + \quad \cdots \quad \curvearrowright + \quad \curvearrowright + \quad \uparrow + \\ \downarrow + \quad \curvearrowright + \quad \cdots \quad \curvearrowright + \quad \curvearrowright + \quad \uparrow + \end{array} & \text{if } n \text{ is even} \\ \begin{array}{c} \downarrow + \quad \curvearrowright + \quad \cdots \quad \curvearrowright + \quad \curvearrowright + \quad \uparrow + \\ \downarrow + \quad \curvearrowright + \quad \cdots \quad \curvearrowright + \quad \curvearrowright + \quad \uparrow + \end{array} & \text{if } n \text{ is odd,} \end{cases} \quad (4.38)$$

where there are $n-1$ caps in total. For $n \geq 0$, let

$$\mathbf{r}_n := \begin{cases} \begin{array}{c} \downarrow \boxed{x_1^{n-1}} \uparrow + \cdots + \downarrow \boxed{x_1^2} \uparrow + \downarrow \boxed{x_1} \uparrow + \downarrow \uparrow + \\ \downarrow \boxed{x_1^{n-1}} \uparrow + \cdots + \downarrow \boxed{x_1^2} \uparrow + \downarrow \boxed{x_1} \uparrow + \downarrow \uparrow + \end{array} & \text{if } n \text{ is even} \\ \begin{array}{c} \downarrow \boxed{x_1^{n-1}} \uparrow + \cdots + \downarrow \boxed{x_1^2} \uparrow + \downarrow \boxed{x_1} \uparrow + \downarrow \uparrow + \\ \downarrow \boxed{x_1^{n-1}} \uparrow + \cdots + \downarrow \boxed{x_1^2} \uparrow + \downarrow \boxed{x_1} \uparrow + \downarrow \uparrow + \end{array} & \text{if } n \text{ is odd.} \end{cases} \quad (4.39)$$

In particular, $\mathbf{r}_0 = \text{id}_A$. Note that the endomorphism \mathbf{r}_n is of degree $2\binom{n}{2}$.

Lemma 4.21. *Suppose that $n = l$ or $n = l-1$. Then*

$$\sum_{b \in \mathbb{B}_{O_{\pm}}^{O_{\pm}}} \text{tr}_{O_{+}}^O (x_1^{n-1} \text{tr}_{O_{-}}^O (\cdots (x_1^3 \text{tr}_{O_{-}}^O (x_1^2 \text{tr}_{O_{+}}^O (x_1 \text{tr}_{O_{-}}^O (b))) \cdots))) \otimes b^{\vee} = (-1)^{\binom{n}{2}} 1 \otimes 1$$

if n is even, and

$$\sum_{b \in \mathbb{B}_{O_{\pm}}^{O_{\pm}}} \text{tr}_{O_{-}}^O (x_1^{n-1} \text{tr}_{O_{+}}^O (\cdots (x_1^3 \text{tr}_{O_{-}}^O (x_1^2 \text{tr}_{O_{+}}^O (x_1 \text{tr}_{O_{-}}^O (b))) \cdots))) \otimes b^{\vee} = (-1)^{\binom{n}{2}} 1 \otimes 1$$

if n is odd.

Proof. Let $\partial_{\pm 1}^{(i)} := \partial_i \cdots \partial_2 \partial_{\pm 1}$, interpreted as $\partial_{\pm 1}$ if $i = 1$ and as 1 if $i = 0$. A reduced expression for the longest element of $(W_{O_{\pm}}/W_{O_{+}})_{\min}$ is $s_{(-1)^{l-1}}(s_2 s_{(-1)^{l-2}}) \cdots (s_{l-2} \cdots s_2 s_1)(s_{l-1} \cdots s_2 s_{-1})$. Hence, recalling (3.5), $\text{tr}_{O_{\pm}}^{O_{+}} = \partial_{(-1)^{l-1}}^{(1)} \partial_{(-1)^{l-2}}^{(2)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}$. Also $\text{tr}_{O_{+}}^O = \partial_1^{(l-1)}$ and $\text{tr}_{O_{-}}^O = \partial_{-1}^{(l-1)}$.

By degree considerations, all of the terms in the summation to be computed are 0 except for the one in which b is of maximal degree. We choose this top degree b so that the corresponding b^\vee equals 1. Thus, b is an element of R^{O^+} such that

$$\partial_{(-1)^{l-1}}^{(1)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b) = 1. \quad (4.40)$$

Now the result we are trying to prove is reduced to checking that

$$\partial_{(-1)^n}^{(l-1)} \left(x_1^{n-1} \partial_{(-1)^{n-1}}^{(l-1)} \left(\cdots \left(x_1^2 \partial_1^{(l-1)} \left(x_1 \partial_{-1}^{(l-1)}(b) \right) \right) \cdots \right) \right) = (-1)^{\binom{n}{2}}$$

for $n = l - 1$ or $n = l$. In view of (4.40), this follows from the $n = l - 1$ or $n = l$ cases of the following more general statement: for $1 \leq n \leq l$ we have that

$$\partial_{(-1)^n}^{(l-1)} \left(x_1^{n-1} \partial_{(-1)^{n-1}}^{(l-1)} \left(\cdots \left(x_1^2 \partial_1^{(l-1)} \left(x_1 \partial_{-1}^{(l-1)}(b) \right) \right) \cdots \right) \right) = (-1)^{\binom{n}{2}} \partial_{(-1)^n}^{(l-n)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b). \quad (4.41)$$

Now we proceed to prove (4.41) by induction on $n = 1, \dots, l$.

The base case $n = 1$ is vacuous—it is just asserting that $\partial_{-1}^{(l-1)}(b) = \partial_{-1}^{(l-1)}(b)$. For the induction step, we assume that (4.41) is true for some $1 \leq n \leq l - 1$, and must show that

$$\partial_{(-1)^{n+1}}^{(l-1)} \left(x_1^n \cdot (-1)^{\binom{n}{2}} \partial_{(-1)^n}^{(l-n)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b) \right) = (-1)^{\binom{n+1}{2}} \partial_{(-1)^{n+1}}^{(l-n-1)} \partial_{(-1)^n}^{(l-n)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b).$$

Equivalently,

$$\partial_{l-1} \cdots \partial_2 \partial_{(-1)^{n+1}} \left((-x_1)^n \cdot \partial_{(-1)^n}^{(l-n)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b) \right) = \partial_{(-1)^{n+1}}^{(l-n-1)} \partial_{(-1)^n}^{(l-n)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b). \quad (4.42)$$

To prove this, we will need the identity

$$\partial_i \partial_{(-1)^m}^{(l-m)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b) = 0, \quad (4.43)$$

for $2 \leq i \leq l - 1$ and $1 \leq m \leq l - 1$. To prove (4.43), we observe that

$$s_i(s_{l-m} \cdots s_2 s_{(-1)^m}) \cdots (s_{l-2} \cdots s_2 s_1)(s_{l-1} \cdots s_2 s_{-1})$$

is either not a reduced expression or it is braid-equivalent to ws_j for some $j \in \{1, 2, \dots, l - 1\}$. This can be seen by drawing the permutation diagram, or one can argue algebraically by induction on m . In the former case, (4.43) follows as the unreduced product of Demazure's is 0. In the latter case, (4.43) follows because $\partial_j(b) = 0$.

To prove (4.42), we use the product rule $\partial_i(fg) = \partial_i(f)g + s_i(f)\partial_i(g)$ to apply $\partial_{(-1)^{n+1}}, \partial_2, \dots, \partial_{l-1}$ in order to the product $(-x_1)^n \cdot \partial_{(-1)^n}^{(l-n)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b)$, starting with $\partial_{(-1)^{n+1}}$. In this way, we expand the left hand side of (4.42) as a sum of 2^{l-1} terms. As soon as one of the Demazure operators acts on the term $(-x_1)^n$ on the left, all remaining Demazure operators must also act on this term, else we get 0 by (4.43). Thus, the only potentially non-zero terms are

$$\partial_{l-1} \cdots \partial_{l-m} \left((-x_{l-m})^n \right) \cdot \partial_{l-m-1} \cdots \partial_2 \partial_{(-1)^{n+1}} \partial_{(-1)^n}^{(l-n)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b).$$

for $0 \leq m \leq l$. If $m > n$ then $\partial_{l-1} \cdots \partial_{l-m} \left((-x_{l-m})^n \right) = 0$ by degree considerations. So we may assume that $m \leq n$. If $m < n = l - 1$ the term is 0 since

$$\partial_{l-m-1} \cdots \partial_2 \partial_{(-1)^l} \partial_{(-1)^{l-1}}^{(1)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b) \stackrel{(4.40)}{=} \partial_{l-m-1} \cdots \partial_2 \partial_{(-1)^l}(1) = 0.$$

If $m < n < l - 1$ the term is 0 since

$$\partial_{l-m-1} \cdots \partial_2 \partial_{(-1)^{n+1}} \partial_{(-1)^n}^{(l-n)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b) = \partial_{l-m-1} \cdots \partial_{l-n} \partial_{(-1)^{n+1}}^{(l-n-1)} \partial_{(-1)^n}^{(l-n)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b),$$

which is 0 by (4.43) (taking i there to be $l - n$). Thus we are left just with the $m = n$ term

$$\partial_{l-1} \cdots \partial_{l-n} ((-x_{l-n-1})^n) \cdot \partial_{(-1)^{n+1}}^{(l-n-1)} \partial_{(-1)^n}^{(l-n)} \cdots \partial_1^{(l-2)} \partial_{-1}^{(l-1)}(b),$$

which is equal to the right hand side of (4.42) by a variant of Lemma 4.12. \square

Corollary 4.22. *For $n = l - 1$ or $n = l$, we have that*

$$\hat{\mathbf{u}}_n \circ \mathbf{r}_n \circ \check{\mathbf{u}}_n = \begin{cases} + \uparrow \pm \downarrow + & \text{if } n \text{ is even} \\ + \uparrow \pm \downarrow + & \text{if } n \text{ is odd.} \end{cases}$$

Proof. This follows from Lemma 4.21 by a calculation which is similar to the proof of Lemma 3.4. We illustrate with an example, taking $n = 5$. We recommend thinking also about an example in which n is even, at which point the pattern is clear. We need to show that

This is a situation in which we find it easiest to expand the special undotted and dotted strings as ordinary and dashed strings using (4.20) and (4.21). Remembering signs which arise because $\gamma(x_1) = -x_1$, this reduces the problem to proving that

in the ordinary string calculus for $\mathfrak{B} \subseteq \mathfrak{Bim}$. Now we simplify the left hand side of this working from right to left, starting with an application of hard RII (3.22), to get

Then we use (3.18) and easy RII (3.21) to move the rightmost bubble inside the second bubble from the right, and evaluating the internal bubble using (3.19) to reduce to

Repeating in this way eventually yields

$$\sum_{b \in \mathbb{B}_{O_{\pm}^0}^+} - \left(\text{tr}_{O_-^0}^O (x_1^4 \text{tr}_{O_+^0}^O (x_1^3 \text{tr}_{O_-^0}^O (x_1^2 \text{tr}_{O_+^0}^O (x_1 \text{tr}_{O_-^0}^O (b)))))) \right) \uparrow_{\pm} \downarrow_{\pm} (b^{\vee}) + .$$

It remains to apply Lemma 4.21 to obtain the result. □

Corollary 4.23. *For $2 \leq n \leq l - 1$, we have that*

$$\hat{\mathbf{v}}_n \circ \mathbf{r}_n \circ \check{\mathbf{u}}_n = \begin{cases} \begin{array}{c} \hat{n}+ \quad \hat{n}\pm \quad \hat{n}+ \\ \uparrow \quad \downarrow \\ \pm \\ \downarrow \\ \pm \end{array} & \text{if } n \text{ is even} \\ \begin{array}{c} \hat{n}+ \quad \hat{n}\pm \quad \hat{n}+ \\ \uparrow \quad \downarrow \\ \pm \\ \downarrow \\ \pm \end{array} & \text{if } n \text{ is odd.} \end{cases}$$

Proof. Again we explain the calculation with an example, taking $n = 4$. By the definitions, we have that

$$\hat{\mathbf{v}}_4 \circ \mathbf{r}_4 \circ \check{\mathbf{u}}_4 = + \begin{array}{c} \hat{n}+ \quad \hat{n}\pm \quad \hat{n}+ \\ \uparrow \quad \downarrow \\ \hat{n} \quad \hat{n} \quad \hat{n} \\ \circ \quad \circ \quad \circ \\ \downarrow \quad \downarrow \quad \downarrow \\ \pm \end{array} + .$$

Now we use easy RII, easy RIII and (3.18) to slide the big rightward cap downward to obtain

$$\begin{array}{c} \hat{n}+ \quad \hat{n}\pm \quad \hat{n}+ \\ \uparrow \quad \downarrow \\ \hat{n} \quad \hat{n} \quad \hat{n} \\ \circ \quad \circ \quad \circ \\ \downarrow \quad \downarrow \quad \downarrow \\ \pm \end{array} + .$$

The circles can now be removed by applying Corollary 4.22—the presence of the \hat{n} means that we are effectively working inside a D_n subsystem, so the hypothesis of Corollary 4.22 is valid. □

Lemma 4.24. *For $2 \leq n \leq l - 1$, we have that*

$$\sum_{b \in \mathbb{B}_{O_+^0}^O} b^{\vee} \text{tr}_{O_+^0}^{O\hat{n}} \left(b (\eta_{O_-^0}^{O\hat{n}})^{\lfloor \frac{n-1}{2} \rfloor} (\eta_{O_+^0}^{O\hat{n}})^{\lfloor \frac{n-1}{2} \rfloor} \right) = 1.$$

Proof. A reduced expression for the longest element of $(W_{O_+}/W_{O\hat{n}})_{\min}$ is

$$(s_{l-1} \cdots s_2 s_1)(s_{l-n+1} \cdots s_{l-1}) \cdots (s_3 \cdots s_{n+1})(s_2 \cdots s_n).$$

The corresponding product of Demazure operators computes $\text{tr}_{O_+^0}^{O\hat{n}}$. We also have that $\eta_{O_+^0}^{O\hat{n}} = (x_{n+1} - x_1)(x_{n+2} - x_1) \cdots (x_l - x_1)$ and $\eta_{O_-^0}^{O\hat{n}} = (x_{n+1} + x_1)(x_{n+2} + x_1) \cdots (x_l + x_1)$.

By degree considerations, all terms in the sum vanish except for the one in which b is of maximal degree. For this, we take $b = (-x_1)^{l-1}$ and $b^\vee = 1$. So the expression we are trying to calculate is

$$(\partial_{l-1} \cdots \partial_1)(\partial_{l-n+1} \cdots \partial_{l-1}) \cdots (\partial_3 \cdots \partial_{n+1})(\partial_2 \cdots \partial_n) \left((-x_1)^{l-1} \prod_{i=n+1}^l \left[(x_i + x_1)^{\lfloor \frac{n-1}{2} \rfloor} (x_i - x_1)^{\lfloor \frac{n-1}{2} \rfloor} \right] \right).$$

First, note that $(-x_1)^{l-1}$ is invariant under s_i for $i = 2, \dots, l-1$, and therefore commutes past the product of Demazure operators $(\partial_{l-n+1} \cdots \partial_{l-1}) \cdots (\partial_3 \cdots \partial_{n+1})(\partial_2 \cdots \partial_n)$. Now we observe that $\partial_2 \cdots \partial_n \left((x_{n+1} + x_1)^{\lfloor \frac{n-1}{2} \rfloor} (x_{n+1} - x_1)^{\lfloor \frac{n-1}{2} \rfloor} \right) = 1$. All that matters for this is that the expression inside the brackets is a monic polynomial of degree $(n-1)$ in x_{n+1} with coefficients in $\mathbb{k}[x_1]$, and $\partial_2 \cdots \partial_n (x_{n+1}^{n-1}) = 1$. Similarly, $\partial_3 \cdots \partial_{n+2} \left((x_{n+2} + x_1)^{\lfloor \frac{n-1}{2} \rfloor} (x_{n+2} - x_1)^{\lfloor \frac{n-1}{2} \rfloor} \right) = 1$, and so on. In this way, we reduce until we are left with $\partial_{l-1} \cdots \partial_1 ((-x_1)^{l-1})$. This is 1 by Lemma 4.12. \square

Corollary 4.25. *For $2 \leq n \leq l-1$, we have that*

$$\mathbf{w}_n \circ \mathbf{v}_n \circ \mathbf{r}_n \circ \check{\mathbf{u}}_n = \begin{cases} \begin{array}{c} \text{red arc} \\ + \\ \text{green arc} \\ \pm \end{array} & \text{if } n \text{ is even} \\ \begin{array}{c} \text{red arc} \\ + \\ \text{green arc} \\ \pm \end{array} & \text{if } n \text{ is odd.} \end{cases}$$

Proof. We first write $\mathbf{v}_n = \check{\mathbf{v}}_n \circ \hat{\mathbf{v}}_n$ and use Corollary 4.23 to simplify the bottom $\hat{\mathbf{v}}_n \circ \mathbf{r}_n \circ \check{\mathbf{u}}_n$ part of $\mathbf{w}_n \circ \check{\mathbf{v}}_n \circ \hat{\mathbf{v}}_n \circ \mathbf{r}_n \circ \check{\mathbf{u}}_n$. For example, if $n = 6$, it shows that

$$\mathbf{w}_6 \circ \mathbf{v}_6 \circ \mathbf{r}_6 \circ \check{\mathbf{u}}_6 = \begin{array}{c} \text{diagram with red and green arcs} \\ + \end{array}$$

Now it seems simplest to rewrite the special strings as ordinary strings using (4.20) and (4.21), sliding the dashed string to the top, to reduce the proof to showing using the ordinary string calculus that

$$\begin{array}{c} \text{diagram with red and green arcs} \\ + \end{array} = \begin{array}{c} \text{red arc} \\ + \\ \text{green arc} \\ \pm \end{array}$$

To prove this, we use hard RII (3.22) to commute the five caps at the top past the big cap to obtain

$$\begin{array}{c} \text{diagram with red and green arcs and caps} \\ + \end{array}$$

5. MAIN RESULTS

Now we prove the main results, which explain the combinatorial coincidences observed in Theorem 4.5. Throughout, $l \geq 2$ is an integer with $l \equiv t \pmod{2}$. Nil-Brauer notation is as in Section 2, and all other notation is for the type D_l root system as in Section 4. In particular, $A = \mathbb{k}[x_1, \dots, x_l] = R^{O+}$.

5.1. Construction of the monoidal functor Θ . Recall the subalgebra $\Lambda^{[2]}$ of the algebra of symmetric functions defined just before Lemma 2.1. As discussed in Remark 2.2, the image of $\Lambda^{[2]}$ under the evaluation homomorphism $\text{ev}_l : \Lambda \rightarrow A$ is the subalgebra generated by $e_r(x_1^2, \dots, x_l^2)$ ($1 \leq r \leq l$), that is, the algebra of invariants for the Weyl group of type B_l . As in the introduction, we denote this subalgebra by C . The action of $\Lambda^{[2]}$ factors through this quotient (via ev_l) to make the cyclotomic nil-Brauer category \mathbf{cNB}_l from Definition 2.10 into a strict C -linear graded monoidal category.

The endomorphism category $\mathbf{End}_{e\mathfrak{B} \otimes \mathfrak{B}\text{im}}(A)$ is a graded C -linear monoidal category whose objects are graded (A, A) -bimodules, and whose tensor product is $- \otimes_A -$. This will be our main focus from now on: we will show it categorifies the $\mathbb{Z}[q, q^{-1}]$ -algebra \mathbf{V} appearing in Theorem 4.5. We remark that the graded Karoubi envelope of $\mathbf{End}_{e\mathfrak{B} \otimes \mathfrak{B}\text{im}}(A)$ is generated as a graded Karoubian category by the special bimodule B from (4.23). This follows from Corollary 4.6.

Theorem 5.1. *There is a strict C -linear graded monoidal functor $\Theta : \mathbf{cNB}_l \rightarrow \mathbf{End}_{e\mathfrak{B} \otimes \mathfrak{B}\text{im}}(A)$ mapping the generating object B of \mathbf{cNB}_l to the bimodule B , and defined on generating morphisms by*

$$\downarrow \mapsto + \downarrow \begin{array}{c} \textcircled{x_1} \\ \downarrow \end{array} \uparrow + = - \left(+ \downarrow \begin{array}{c} \textcircled{x_1} \\ \downarrow \end{array} \uparrow + \right), \quad (5.1)$$

$$\cup \mapsto \begin{array}{c} \text{---} \uparrow \text{---} \\ \text{---} \downarrow \text{---} \end{array} + = \begin{array}{c} \text{---} \downarrow \text{---} \\ \text{---} \uparrow \text{---} \end{array} +, \quad (5.2)$$

$$\cap \mapsto \begin{array}{c} \text{---} \downarrow \text{---} \\ \text{---} \uparrow \text{---} \end{array} + = \begin{array}{c} \text{---} \uparrow \text{---} \\ \text{---} \downarrow \text{---} \end{array} +, \quad (5.3)$$

$$\times \mapsto + \begin{array}{c} \text{---} \uparrow \text{---} \\ \text{---} \downarrow \text{---} \end{array} \begin{array}{c} \text{---} \uparrow \text{---} \\ \text{---} \downarrow \text{---} \end{array} + = + \begin{array}{c} \text{---} \uparrow \text{---} \\ \text{---} \downarrow \text{---} \end{array} \begin{array}{c} \text{---} \uparrow \text{---} \\ \text{---} \downarrow \text{---} \end{array} +. \quad (5.4)$$

In the case $l = 2$, the circle of color 2 and all labels $\hat{2}$ here should simply be omitted. Furthermore:

- (1) For any symmetric polynomial $f \in A$, $\Theta(\downarrow \textcircled{f}) = \downarrow \textcircled{f} +$, which is the endomorphism of A defined by multiplication by f .
- (2) The functor Θ intertwines the symmetry \mathfrak{R} of \mathbf{cNB}_l arising from (2.13) with the symmetry \mathfrak{R} of $\mathbf{End}_{e\mathfrak{B} \otimes \mathfrak{B}\text{im}}(A)$ arising from (4.17).


Proof. We first justify the equalities in the formulae for the images of the generators under Θ in the statement of the theorem. This follows in each case by expanding the definitions of the undotted and dotted special strings like in (4.22). For the dot, it is a special case of (4.25) since $\gamma(x_1) = -x_1$. The other three are obvious once the definitions have been expanded, using the easy properties of the dashed string; for the crossing, we already used a similar strategy in the proof of Lemma 4.16. These equalities show that $\mathfrak{R} \circ \Theta = \Theta \circ \mathfrak{R}$ on all four of the generating morphisms of \mathbf{cNB}_l . Once the well-definedness of Θ has been established, this is all that is needed to prove the property (2).

Next we check that the degrees are consistent. The dot has degree 2, which is the same as the degree of x_1 . For the cap, the clockwise cap in $e\mathfrak{B} \otimes \mathfrak{B}\text{im}$ is of degree $\ell(w_{O-}) - \ell(w_O) = \ell(w_{O+}) - \ell(w_O)$ and the counterclockwise cap is of degree $\ell(w_O) - \ell(w_{O+}) = \ell(w_O) - \ell(w_{O-})$, giving the required total

degree 0. The cup is similar. For the crossing, note from the definition that $\Theta \left(\begin{array}{c} \diagup \\ \diagdown \end{array} \right)$ is \mathbf{v}_2 from (4.37), which is of the desired degree -2 thanks to Lemma 4.20.


To prove the existence of Θ , it suffices to check the defining relations for \mathbf{NB}_t , that is, the eight relations (2.7) to (2.10) plus the cyclotomic relations from Definition 2.10. We are first going to check (2.7) to (2.10). This establishes the existence of a graded $\Lambda^{[2]}$ -linear monoidal functor $\widehat{\Theta} : \mathbf{NB}_t \rightarrow \mathbf{End}_{\mathfrak{eB} \subseteq \mathfrak{Bim}}(A)$, viewing $\mathbf{End}_{\mathfrak{eB} \subseteq \mathfrak{Bim}}(A)$ as a $\Lambda^{[2]}$ -linear monoidal category via the evaluation homomorphism $\Lambda^{[2]} \rightarrow C$. After that, we will discuss property (1), leaving the cyclotomic relation to the end.

- The zig-zag relations from (2.7) follow easily using (3.17) and the zig-zag relations for the dashed string.
- The bubble relation from (2.7) follows by Corollary 4.14, taking $n = 0$.
- The first relation from (2.9) follows by the Interchange Law. The sign appears due to (4.25) since $\gamma(x_1) = -x_1$.
- The first relation from (2.8) follows by cyclicity of the diagrammatic calculus for $\mathfrak{eB} \subseteq \mathfrak{Bim}$:



In view of the alternative forms for the cap and the crossing in (5.3) and (5.4), which was already established in the opening paragraph of the proof, this is what we want.

- The dot sliding relation from (2.9) follows because



The difficult step here is the equality (*). Subtracting (4.34) from (4.33), one obtains a result similar to the above except with $\frac{1}{2}(\alpha_{-1} - \alpha_1)$ in certain regions labeled $\hat{2}$. But $x_1 = \frac{1}{2}(\alpha_{-1} - \alpha_1)$. Since x_1 is invariant under s_2 (unlike $\alpha_{\pm 1}$), it slides across the string of color 2 to the desired region with label O .

- For the Reidemeister I relation from (2.8), we give an easy degree argument. The curl is of degree -2 , so it maps to a graded bimodule homomorphism $B \otimes_A B \rightarrow A$ of degree -2 . This homomorphism must be 0 because by adjunction we have that

$$\text{Hom}_{A-A}(B \otimes_A B, A) \cong \text{Hom}_{R^{O+} \cdot R^{O+}}(B_{O+, O-} \otimes_{R^{O-}} B_{O-, O+}, R^{O+}) \cong \text{End}_{R^{O-} \cdot R^{O+}}(B_{O-, O+}).$$

Since B_{O^-, O^+} is isomorphic to a shift of R^O as a graded (R^{O^-}, R^{O^+}) -bimodule, they have the same endomorphism ring. As a bimodule R^O is generated by 1 (a simple argument on linear generators) and is non-negatively graded, so its endomorphism ring is zero in negative degrees.

- For the Reidemeister II relation from (2.10), we must show that the degree -4 endomorphism

$$\Theta \left(\text{crossing} \right) = + \text{diagram} + \text{diagram} \quad (5.5)$$

of the graded (A, A) -bimodule $B \otimes_A B$ is 0. One way to see this is to use the homomorphism formula (Theorem 3.10) together with the $n = 2$ case of Corollary 4.8 to prove that the lowest degree of any non-zero endomorphism of this bimodule is -2 . We regard this as a hard degree argument since it is applying some heavy machinery. Here is a more elementary proof, which also prepares for the proof of Reidemeister III in the next point. Identifying B simply with $q^{l-1} s_0 R^O$ in the obvious way, $B \otimes_A B$ is generated by the vectors $x_1^r \otimes x_1^s$ for $r, s \geq 0$, so it suffices to show that the endomorphism (5.5) is 0 on $x_1^r \otimes x_1^s$ for all $r, s \geq 0$. This follows by easy degree considerations in the case $r = s = 0$. To deduce the vanishing in general, note that the image of $x_1^r \otimes x_1^s$ under (5.5) is the same as the image of $1 \otimes 1$ under the following:


$$\Theta \left((-1)^s \text{crossing}_{r,s} \right) = + \text{diagram} + \text{diagram} \quad (5.6)$$

We have checked that the relations (2.7) to (2.9) are satisfied in $\mathbf{End}_{e\mathfrak{B} \otimes \mathfrak{B} \text{im}}(A)$, so we already have in our hands a monoidal functor from the monoidal category defined in the same way as \mathbf{NB}_r but omitting the relations (2.10) to $\mathbf{End}_{e\mathfrak{B} \otimes \mathfrak{B} \text{im}}(A)$. From this and (2.15), we deduce that the endomorphism (5.6) is equal to

$$\Theta \left((-1)^s \text{crossing}_{r,s} \right) = + \text{diagram} + \text{diagram} \quad (5.7)$$

Finally, the endomorphism (5.7) maps $1 \otimes 1$ to 0 by the $r = s = 0$ case treated earlier.

- For the Reidemeister III relation from (2.8), we must show that the degree -6 endomorphism



of the graded (A, A) -bimodule $B \otimes_A B \otimes_A B$ is 0. Identifying each B with $q^{l-1} s_0 R^O$, this bimodule is generated by $x_1^r \otimes x_1^s \otimes x_1^t$ for $r, s, t \geq 0$, so it suffices to show that the above endomorphism sends each of these vectors to 0. This follows by same strategy as was just used to prove Reidemeister II, using (2.16) in place of (2.15).

At this point, we have in our hands a $\Lambda^{[2]}$ -linear monoidal functor $\hat{\Theta} : \mathbf{NB}_l \rightarrow \mathbf{End}_{\mathfrak{eB}\mathfrak{B}\mathfrak{im}}(A)$. Next, we show for any $f \in \Lambda$ that $\hat{\Theta}(f)$ is the endomorphism $\overline{\text{ev}_l(f)}$ of A defined by multiplication by the image of f under the evaluation homomorphism $\Lambda \twoheadrightarrow A$. This implies the statement (1). Since $\hat{\Theta}$ is $\Lambda^{[2]}$ -linear by construction, and Λ is generated by $\Lambda^{[2]}$ together with the symmetric functions q_n for all $n > 0$, it suffices to show that $\hat{\Theta}(q_n) = \overline{\text{ev}_l(q_n)}$. By (2.17), q_n corresponds (up to scalar) to a dotted bubble \bigcirc_n in the nilBrauer category, and Corollary 4.14 shows that $\hat{\Theta}$ sends the dotted bubble to q_n (up to the inverse scalar) in $\mathbf{End}_{\mathfrak{eB}\mathfrak{B}\mathfrak{im}}(A)$.

Now we complete the proof by checking the cyclotomic relations, so that $\hat{\Theta}$ descends to the cyclotomic quotient \mathbf{cNB}_l . Since $e_r(x_1, \dots, x_l) = 0$ for $r > l$, the relations involving bubbles are clear. To show that $\hat{\Theta}$ maps (2.29) to 0, it suffices to show that the left action of $e_l(x_1, \dots, x_l) = x_1 \cdots x_l$ on B differs from the right action by a sign. This holds because B is a twisted bimodule. \square

Remark 5.2. Some of the relation checks in the proof of Theorem 5.1 were performed directly using diagrammatic transformations. However, we gave indirect arguments for the Reidemeister I, II and III relations. In fact, it is not hard to give a direct proof of the Reidemeister I relation, using (3.19), (3.21) and (3.22). To prove the Reidemeister II relation diagrammatically, one needs additional relations not recorded in Subsection 3.3, namely, the idempotent decompositions which categorify the type A MOY relations. Using these, one can simplify the inside of the middle circle in (5.7). However, we have *not* been able to work out a purely diagrammatical proof of Reidemeister III, and expect that such an argument would require the use of idempotent decompositions specific to type D.

5.2. Characters of indecomposables. Restricting from the graded bicategory $\mathfrak{eB}\mathfrak{B}\mathfrak{im}$ just to the graded monoidal category $\mathbf{End}_{\mathfrak{eB}\mathfrak{B}\mathfrak{im}}(A)$, and recalling the definition (4.8), the Soergel-Williamson character map (modified slightly to fit the extended setup) gives an isomorphism of $\mathbb{Z}[q, q^{-1}]$ -algebras

$$\text{ch} : K_0(\mathbf{gKar}(\mathbf{End}_{\mathfrak{eB}\mathfrak{B}\mathfrak{im}}(A))) \xrightarrow{\sim} \mathbf{V}. \tag{5.8}$$

Hence, by Theorem 4.5, $K_0(\mathbf{gKar}(\mathbf{End}_{\mathfrak{eB}\mathfrak{B}\mathfrak{im}}(A)))$ is isomorphic as a module to the integral form $\mathbf{V}(l)$ for the $(l + 1)$ -dimensional $U_q \mathfrak{sl}_2$ -module, and as an algebra to the quotient U_l/I_l from Remark 2.13. By Theorem 3.11 and the classification of double cosets in Lemmas 4.3 and 4.4, we know that a full set of self-dual indecomposable objects in $\mathbf{gKar}(\mathbf{End}_{\mathfrak{eB}\mathfrak{B}\mathfrak{im}}(A))$ are given by the (A, A) -bimodules

$B_{(O+)d_n(O+)}$ for $0 \leq n \leq l$ with n even and $s_0A \otimes_A B_{(O-)s_0d_n(O+)}$ for $1 \leq n \leq l$ with n odd. The next fundamental theorem is an application of the nil-Brauer theory from [BWW23] to deduce character formulae for these special singular Soergel bimodules.

From now on, we use \mathbf{e}_n to denote the image of the idempotent (2.18) in the quotient \mathbf{cNB}_l of \mathbf{NB}_l . So \mathbf{e}_n is a homogeneous idempotent (possibly 0) in $\text{End}_{\mathbf{cNB}_l}(B^{*n})$. Let

$$\mathbf{f}_n := \Theta(\mathbf{e}_n), \quad (5.9)$$

which is a homogeneous idempotent (possibly 0) in $\text{End}_{A-A}(B^{\otimes n})$ defining a summand $[B^{\otimes n}, \mathbf{f}_n]$ of this bimodule. We set

$$B^{[n]} := q^{\binom{n}{2}} [B^{\otimes n}, \mathbf{f}_n]. \quad (5.10)$$

The similarity of this notation with (2.22) is deliberate.

Theorem 5.3. *For $0 \leq n \leq l$, we have that*

$$B^{[n]} \cong \begin{cases} B_{(O+)d_n(O+)} & \text{if } n \text{ is even} \\ s_0R^{O-} \otimes_{R^{O-}} B_{(O-)s_0d_n(O+)} & \text{if } n \text{ is odd.} \end{cases} \quad (5.11)$$

The characters of these bimodules are equal to the corresponding Kazhdan-Lusztig basis elements:

$$\text{ch}(B^{[n]}) = \begin{cases} b_{(O+)d_n(O+)} & \text{if } n \text{ is even} \\ s_0b_{(O-)s_0d_n(O+)} & \text{if } n \text{ is odd.} \end{cases} \quad (5.12)$$

In particular, these characters are independent of the characteristic of the ground field.

Proof. In $\mathfrak{gKar}(\mathbf{NB}_l)$, we have by (2.23) that

$$B^{*n} \cong \bigoplus_{i=0}^{\lfloor \frac{n}{2} \rfloor} [n-2i]_q! \left(\sum_{\lambda \in \text{Par}_{\neq t}(i \times (n-2i))} [\lambda_1 + 1]_q^2 \cdots [\lambda_i + 1]_q^2 \right) B^{(n-2i)}.$$

Applying $\mathfrak{gKar}(\Theta)$, the extension of Θ to the graded Karoubi envelopes, and using the definition (5.9), we deduce in $\mathfrak{gKar}(\mathbf{End}_{\mathfrak{B} \otimes \mathfrak{B}}(\mathfrak{Bim}(A)))$ that

$$B^{\otimes n} \cong \bigoplus_{i=0}^{\lfloor \frac{n}{2} \rfloor} [n-2i]_q! \left(\sum_{\lambda \in \text{Par}_{\neq t}(i \times (n-2i))} [\lambda_1 + 1]_q^2 \cdots [\lambda_i + 1]_q^2 \right) B^{[n-2i]}. \quad (5.13)$$

Since $\text{ch}(B^{\otimes n}) = b^{on}$ and the triangular transition matrix in the above decomposition is the same as in Corollary 4.6, we deduce that $B^{[n]}$ is a singular Soergel bimodule with character $b_{(O+)d_n(O+)}$ if n is even or $s_0b_{(O-)s_0d_n(O+)}$ if n is odd. Since Kazhdan-Lusztig basis elements are bar-invariant, it follows that $B^{[n]}$ is self-dual. By the homomorphism formula (Theorem 3.10) and Theorem 4.5, the graded dimension of the endomorphism algebra of $B^{[n]}$ is equal to $(b^{(n)}\eta_l, b^{(n)}\eta_l)_l \dim_q R^{O+}$. By (2.33) and (2.35), this is an element of $1 + q^{-1}\mathbb{N}[[q^{-1}]]$. Hence, the degree 0 component of the endomorphism algebra of $B^{[n]}$ is 1-dimensional. This implies that $B^{[n]}$ is indecomposable. Hence, it is isomorphic to $B_{(O+)d_n(O+)}$ or $s_0R^{O-} \otimes_{R^{O-}} B_{(O-)s_0d_n(O+)}$ by the definition of these bimodules (Theorem 3.11). \square

Corollary 5.4. *For $0 \leq n \leq \lfloor \frac{l}{2} \rfloor$, $\text{Hom}_{A-A}(A, B^{[2n]})$ is a free graded A -module of graded rank $q^{-n(l+t-1)} \binom{(l-t)/2}{n}_q$.*

Proof. This follows from the homomorphism formula (Theorem 3.10), using the identification of the character of $B^{[2n]}$ obtained in Theorem 5.3 and the combinatorics from Corollary 4.7. \square

5.3. Identification of the Grothendieck ring of \mathbf{cNB}_l . Theorem 5.3 implies that the homogeneous idempotents \mathbf{f}_n ($0 \leq n \leq l$) are primitive, and moreover that any primitive homogeneous idempotent in $\mathbf{End}_{\mathbf{eB} \otimes \mathbf{Bim}}(A)$ is equivalent to \mathbf{f}_n for a unique $0 \leq n \leq l$. Now we return to the cyclotomic nil-Brauer category \mathbf{cNB}_l .

Theorem 5.5. *For $0 \leq n \leq l$, \mathbf{e}_n is a primitive homogeneous idempotent in \mathbf{cNB}_l , and any primitive homogeneous idempotent in \mathbf{cNB}_l is equivalent to \mathbf{e}_n for a unique $0 \leq n \leq l$.*

Proof. By Theorem 2.6, we know already that the idempotents \mathbf{e}_n ($n \geq 0$) give a complete set of primitive homogeneous idempotents in \mathbf{NB}_l . It follows from general principles that the *non-zero* ones amongst their images give a full set of primitive homogeneous idempotents in \mathbf{cNB}_l . For $n > l$, we know that $\bar{\mathbf{e}}_n = 0$ by Lemma 2.12. To complete the proof of the theorem, it remains to show that \mathbf{e}_n is non-zero in \mathbf{cNB}_l for $0 \leq n \leq l$. This follows from Theorem 5.3 and the definition (5.9). \square

Corollary 5.6. *The monoidal functor $\Theta : \mathbf{cNB}_l \rightarrow \mathbf{End}_{\mathbf{eB} \otimes \mathbf{Bim}}(A)$ induces a $\mathbb{Z}[q, q^{-1}]$ -linear algebra isomorphism $K_0(\mathbf{gKar}(\mathbf{cNB}_l)) \xrightarrow{\sim} K_0(\mathbf{gKar}(\mathbf{End}_{\mathbf{eB} \otimes \mathbf{Bim}}(A)))$ taking $[B^{(n)}]$ to $[B^{[n]}]$ for $0 \leq n \leq l$.*

5.4. Diagrams for primitive idempotents. Recall the bimodule homomorphisms \mathbf{v}_n and \mathbf{r}_n from (4.37) and (4.39). The following theorem gives an explicit diagrammatic description of the homogeneous primitive idempotents \mathbf{f}_n ($0 \leq n \leq l$) in $\mathbf{End}_{\mathbf{eB} \otimes \mathbf{Bim}}(A)$.

Theorem 5.7. $\mathbf{f}_n = \begin{cases} \mathbf{r}_n \circ \mathbf{v}_n & \text{if } 0 \leq n \leq l \\ 0 & \text{if } n > l. \end{cases}$

Proof. By (5.9) and (2.18), we have that $\mathbf{f}_n = \mathbf{r}_n \circ \Theta \left(\begin{array}{c} \mathbf{f} \\ n \end{array} \right)$. Thus, \mathbf{f}_n is the composition of a degree $2 \binom{n}{2}$ endomorphism after a degree $-2 \binom{n}{2}$ endomorphism of $B^{\otimes n}$. It follows that $\mathbf{f}_n = 0$ for $n > l$ since, by Lemma 4.18, any degree $-2 \binom{n}{2}$ endomorphism of $B^{\otimes n}$ is 0 when $n > l$. It is also obvious that $\mathbf{f}_0 = \mathbf{r}_0 \circ \mathbf{v}_0$. Finally, suppose that $1 \leq n \leq l$. Since $\mathbf{f}_n \neq 0$, the degree $-2 \binom{n}{2}$ endomorphism $\Theta \left(\begin{array}{c} \mathbf{f} \\ n \end{array} \right)$ is non-zero. Also \mathbf{v}_n is of degree $-2 \binom{n}{2}$. This is the lowest non-zero degree of $\mathbf{End}_{A-A}(B^{\otimes n})$, and it is 1-dimensional in this degree by Lemma 4.18 again. Hence, $\mathbf{v}_n = c \Theta \left(\begin{array}{c} \mathbf{f} \\ n \end{array} \right)$ and $\mathbf{r}_n \circ \mathbf{v}_n = c \mathbf{f}_n$ for some $c \in \mathbb{k}$. To show that $c = 1$, we use that \mathbf{f}_n is an idempotent so its only non-zero eigenvalue is 1. Thus, it suffices to observe that $\mathbf{r}_n \circ \mathbf{v}_n$ has an eigenvector of eigenvalue 1: the vector $\mathbf{r}_n(1 \otimes \cdots \otimes 1)$ is fixed by $\mathbf{r}_n \circ \mathbf{v}_n$ by Corollary 4.26. \square

Remark 5.8. Here is some additional commentary on the theory underlying Theorem 5.7. Recall that $\mathbf{v}_n = \check{\mathbf{v}}_n \circ \hat{\mathbf{v}}_n$. Thus the idempotent \mathbf{f}_n factors as a composition of $(r_n \circ \check{\mathbf{v}}_n)$ with $\hat{\mathbf{v}}_n$. When n is even, the object being factored through is the singular Bott–Samelson bimodule M associated to the sequence of parabolic subgroups $[O+ \supset O\hat{n}+ \subset O\hat{n}\pm \supset O\hat{n}+ \subset O+]$. This sequence is a reduced expression for the double coset $(O+)d_n(O+)$ in the sense of [EK23] (see also [Wil08, Ch 1.3]). Consequently, by [Wil08, Proof of Theorem 5.4.2], this bimodule M has $B_{(O+)d_n(O+)}$ as a direct summand. Thus it is reasonable to expect that the idempotent \mathbf{f}_n , whose image is isomorphic to $B_{(O+)d_n(O+)}$ by Theorem 5.3, should factor through M . Indeed, the opposite composition $\hat{\mathbf{v}}_n \circ (\mathbf{r}_n \circ \check{\mathbf{v}}_n)$ should yield the idempotent endomorphism of M whose image is $B_{(O+)d_n(O+)}$. Similar statements can be made when n is odd.

REFERENCES

- [Bao19] Huanchen Bao. A categorification of Hecke algebras with parameters 1 and v . *J. Algebra*, 537:26–38, 2019.
 [BSWW18] Huanchen Bao, Peng Shan, Weiqiang Wang, and Ben Webster. Categorification of quantum symmetric pairs I. *Quantum Topol.*, 9:643–714, 2018.

- [BW18a] Huanchen Bao and Weiqiang Wang. Canonical bases arising from quantum symmetric pairs. *Invent. Math.*, 213(3):1099–1177, 2018.
- [BW18b] Huanchen Bao and Weiqiang Wang. A new approach to Kazhdan-Lusztig theory of type B via quantum symmetric pairs. *Astérisque*, 402:vii+134, 2018.
- [BW18c] Collin Berman and Weiqiang Wang. Formulae of t -divided powers in $U_q(\mathfrak{sl}_2)$. *J. Pure Appl. Algebra*, 222(9):2667–2702, 2018.
- [BWW23] Jonathan Brundan, Weiqiang Wang, and Ben Webster. Nil-Brauer categorifies the split t -quantum group of rank one. [arXiv:2305.05877](https://arxiv.org/abs/2305.05877), 2023.
- [BWW24] Jonathan Brundan, Weiqiang Wang, and Ben Webster. The nil-Brauer category. *Ann. Rep. Theory*, 1:21–58, 2024.
- [Dem73] Michel Demazure. Invariants symétriques entiers des groupes de Weyl et torsion. *Invent. Math.*, 21:287–301, 1973.
- [EK23] Ben Elias and Hankyung Ko. A singular Coxeter presentation. *Proc. Lond. Math. Soc. (3)*, 126(3):923–996, 2023.
- [EKLP23] Ben Elias, Hankyung Ko, Nicolàs Libedinsky, and Leonardo Patimo. Demazure operators for double cosets. [arXiv:2307.1502](https://arxiv.org/abs/2307.1502), 2023.
- [EKLP24] Ben Elias, Hankyung Ko, Nicolàs Libedinsky, and Leonardo Patimo. Singular light leaves. [arXiv:2401.03053](https://arxiv.org/abs/2401.03053), 2024.
- [Eli] Ben Elias. Gaitsgory’s central sheaves via the diagrammatic Hecke category. [arXiv:1811.06188](https://arxiv.org/abs/1811.06188).
- [Eli16] Ben Elias. The two-color Soergel calculus. *Compos. Math.*, 152(2):327–398, 2016.
- [Eli17] Ben Elias. Quantum Satake in type A . Part I. *J. Comb. Algebra*, 1(1):63–125, 2017.
- [EMTW20] Ben Elias, Shotaro Makisumi, Ulrich Thiel, and Geordie Williamson. *Introduction to Soergel Bimodules*, volume 5 of *RSME Springer Series*. Springer, Cham, 2020.
- [ESW17] Ben Elias, Noah Snyder, and Geordie Williamson. On cubes of Frobenius extensions. In *Representation Theory—Current Trends and Perspectives*, EMS Ser. Congr. Rep., pages 171–186. Eur. Math. Soc., Zürich, 2017.
- [EW16] Ben Elias and Geordie Williamson. Soergel calculus. *Represent. Theory*, 20:295–374, 2016.
- [KL10] Mikhail Khovanov and Aaron D. Lauda. A categorification of quantum $\mathfrak{sl}(n)$. *Quantum Topol.*, 1:1–92, 2010.
- [Let99] Gail Letzter. Symmetric pairs for quantized enveloping algebras. *J. Algebra*, 220(2):729–767, 1999.
- [Mac15] Ian G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford Classic Texts in the Physical Sciences. The Clarendon Press, Oxford University Press, New York, second edition, 2015.
- [Rou08] Raphaël Rouquier. 2-Kac-Moody algebras. [arXiv:0812.5023](https://arxiv.org/abs/0812.5023), 2008.
- [Soe97] Wolfgang Soergel. Kazhdan-Lusztig-Polynome und eine Kombinatorik für Kipp-Moduln. *Represent. Theory*, 1:37–68, 1997.
- [Wil08] Geordie Williamson. *Singular Soergel Bimodules*. PhD thesis, Albert-Ludwigs-Universität Freiburg, 2008.
- [Wil11] Geordie Williamson. Singular Soergel bimodules. *Int. Math. Res. Not. IMRN*, 20:4555–4632, 2011.
- [Wil23] Geordie Williamson. Erratum for “Singular Soergel bimodules”. [arXiv:1010.1283](https://arxiv.org/abs/1010.1283), 2023.

(E.B.) DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA, USA
Email address: ebodish@mit.edu

(J.B.) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR, USA
URL: pages.uoregon.edu/brundan, *ORCID:* orcid.org/0009-0009-2793-216X
Email address: brundan@uoregon.edu

(B.E.) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR, USA
Email address: belias@uoregon.edu