

SEMI-INFINITE HIGHEST WEIGHT CATEGORIES

JONATHAN BRUNDAN AND CATHARINA STROPPEL

Dedicated to Jens Carsten Jantzen on the occasion of his 70th birthday.

ABSTRACT. We develop axiomatics of highest weight categories and quasi-hereditary algebras in order to incorporate two semi-infinite situations which are in Ringel duality with each other; the underlying posets are either *upper finite* or *lower finite*. We also consider various more general sorts of stratified categories. In the upper finite cases, we give an alternative characterization of these categories in terms of based quasi-hereditary algebras and based stratified algebras, which are certain locally unital algebras possessing triangular bases.

CONTENTS

1. Introduction	1
2. Some finiteness properties on Abelian categories	9
3. Generalizations of highest weight categories	22
4. Tilting modules and semi-infinite Ringel duality	49
5. Generalizations of quasi-hereditary algebras	83
6. Examples	101
References	117

1. INTRODUCTION

Highest weight categories were introduced by Cline, Parshall and Scott [CPS1] in order to provide an axiomatic framework encompassing a number of important examples which had previously arisen in representation theory. In the first part of this article, we give a detailed exposition of two semi-infinite variants, which we call *lower finite* and *upper finite highest weight categories*. Lower finite highest weight categories were already included in the original work of Cline, Parshall and Scott, although they did not use this language. Well-known examples include the category $\mathcal{R}ep(G)$ of finite-dimensional rational representations of a (connected) reductive algebraic group. On the other hand, the upper finite highest weight categories studied here do not fit into the locally Artinian framework of [CPS1]. Nevertheless, there are many examples of upper finite highest weight categories already in the literature, often of a diagrammatic nature, and an appropriate axiomatic framework was sketched out by Elias and Losev in [ELos, §6.1.2]. There are plenty of subtleties, so a full treatment seems desirable.

Then, in the next part, we extend Ringel duality to the semi-infinite setting:

$$\left\{ \begin{array}{c} \text{lower finite} \\ \text{highest weight categories} \end{array} \right\} \xleftrightarrow{\text{Ringel duality}} \left\{ \begin{array}{c} \text{upper finite} \\ \text{highest weight categories} \end{array} \right\}.$$

Other approaches to “semi-infinite Ringel duality” exist in the literature, but these typically require the existence of a \mathbb{Z} -grading; e.g., see [Soe] (in a Lie algebra setting) and also [Maz2]. We avoid this by working with finite-dimensional comodules over a coalgebra in the lower finite case, and with locally finite-dimensional modules over a locally finite-dimensional locally unital algebra in the upper finite case. Another approach to semi-infinite Ringel duality based around pseudo-compact topological algebras was

This article is based upon work done while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California during the Spring 2018 semester. It was supported by the National Science Foundation grant DMS-1700905 and by the HCM in Bonn.

initiated by Marko and Zubkov [MZ]. However, their theory requires some additional finiteness assumptions which are not satisfied in important examples including all non-semisimple categories of the form $\mathcal{R}ep(G)$ for a reductive group G ; see Corollary 4.28, Remark 4.31 and Remark 4.23.

Finally, as an application of semi-infinite Ringel duality, we give an elementary algebraic characterization of upper finite highest weight categories, showing that any such category is equivalent to the category of locally finite-dimensional modules over an *upper finite based quasi-hereditary algebra*. This is an algebraic formulation of the notion of object-adapted cellular category from [ELau, Def. 2.1], and a generalization of the based quasi-hereditary algebras of [KM, Def. 2.4]. As well as Ringel duality, the proof of this characterization uses a construction from [AST] to construct bases for endomorphism algebras of tilting objects. The observation that the bases arising from [AST] are object-adapted cellular bases was made already by Elias and several others, and appears in recent work of Andersen [And].

Throughout the article, we systematically develop the entire theory in the more general setting of what we call ε -stratified categories. The idea of this definition is due to Ágoston, Dlab and Lukács: in [ADL, Def. 1.3] one finds the notion of a stratified algebra of type ε ; the category of finite-dimensional left modules over such a finite-dimensional algebra is an example of a ε -stratified category in our sense. The various other generalizations of highest weight category that have been considered in existing literature fit naturally into our ε -stratified framework.

To explain the contents of the paper in more detail, we start by explaining our precise setup in the finite-dimensional case, since even here it does not seem to have appeared explicitly elsewhere in the literature. Consider a *finite Abelian category*, that is, a category \mathcal{R} equivalent to the category $A\text{-mod}_{\text{fd}}$ of finite-dimensional left A -modules for some finite-dimensional algebra A over an algebraically closed field \mathbb{k} . A *stratification* of \mathcal{R} is a quintuple $(\mathbf{B}, L, \rho, \Lambda, \leq)$ consisting of a set \mathbf{B} , a *labelling function* L such that $\{L(b) \mid b \in \mathbf{B}\}$ is a full set of pairwise inequivalent irreducible objects of \mathcal{R} , and a *stratification function* $\rho : \mathbf{B} \rightarrow \Lambda$ for a poset (Λ, \leq) .

Given a stratification, let $P(b)$ (resp., $I(b)$) be a projective cover (resp., injective hull) of $L(b)$. For $\lambda \in \Lambda$, let $\mathcal{R}_{\leq \lambda}$ (resp., $\mathcal{R}_{< \lambda}$) be the Serre subcategory of \mathcal{R} generated by the irreducibles $L(b)$ for $b \in \mathbf{B}$ with $\rho(b) \leq \lambda$ (resp., $\rho(b) < \lambda$). Define the *stratum* \mathcal{R}_λ to be the Serre quotient $\mathcal{R}_{\leq \lambda}/\mathcal{R}_{< \lambda}$ with quotient functor $j^\lambda : \mathcal{R}_{\leq \lambda} \rightarrow \mathcal{R}_\lambda$. For $b \in \mathbf{B}_\lambda := \rho^{-1}(\lambda)$, let $L_\lambda(b) := j^\lambda L(b)$. These give a full set of pairwise inequivalent irreducible objects in \mathcal{R}_λ . Still for $b \in \mathbf{B}_\lambda$, let $P_\lambda(b)$ (resp., $I_\lambda(b)$) be a projective cover (resp., injective hull) of $L_\lambda(b)$ in \mathcal{R}_λ .

The functor j^λ has a left adjoint $j_!^\lambda$ and a right adjoint j_*^λ . We refer to these as the *standardization* and *costandardization functors*, respectively, following the language of [LW, §2]. Then we introduce the *standard*, *proper standard*, *costandard* and *proper costandard objects* of \mathcal{R} for $\lambda \in \Lambda$ and $b \in \mathbf{B}_\lambda$:

$$\Delta(b) := j_!^\lambda P_\lambda(b), \quad \bar{\Delta}(b) := j_!^\lambda L_\lambda(b), \quad \nabla(b) := j_*^\lambda I_\lambda(b), \quad \bar{\nabla}(b) := j_*^\lambda L_\lambda(b). \quad (1.1)$$

Equivalently, $\Delta(b)$ (resp., $\nabla(b)$) is the largest quotient of $P(b)$ (resp., the largest subobject of $I(b)$) that belongs to $\mathcal{R}_{\leq \lambda}$, and $\bar{\Delta}(b)$ (resp., $\bar{\nabla}(b)$) is the largest quotient of $\Delta(b)$ (resp., the largest subobject of $\nabla(b)$) such that all composition factors apart from its irreducible head (resp., its irreducible socle) belong to $\mathcal{R}_{< \lambda}$.

Fix a *sign function* $\varepsilon : \Lambda \rightarrow \{\pm\}$ and define the ε -*standard* and ε -*costandard objects*

$$\Delta_\varepsilon(b) := \begin{cases} \Delta(b) & \text{if } \varepsilon(\rho(b)) = + \\ \bar{\Delta}(b) & \text{if } \varepsilon(\rho(b)) = - \end{cases}, \quad \nabla_\varepsilon(b) := \begin{cases} \bar{\nabla}(b) & \text{if } \varepsilon(\rho(b)) = + \\ \nabla(b) & \text{if } \varepsilon(\rho(b)) = - \end{cases}. \quad (1.2)$$

By a Δ_ε -*flag* (resp., a ∇_ε -*flag*) of an object of \mathcal{R} , we mean a (necessarily finite) filtration whose sections are of the form $\Delta_\varepsilon(b)$ (resp., $\nabla_\varepsilon(b)$) for $b \in \mathbf{B}$. Then we call \mathcal{R} an ε -*stratified category* if one of the following equivalent properties holds:

($P\Delta_\varepsilon$) For every $b \in \mathbf{B}$, the projective object $P(b)$ has a Δ_ε -flag with sections $\Delta_\varepsilon(c)$ for $c \in \mathbf{B}$ with $\rho(c) \geq \rho(b)$.

($I\nabla_\varepsilon$) For every $b \in \mathbf{B}$, the injective object $I(b)$ has a ∇_ε -flag with sections $\nabla_\varepsilon(c)$ for $c \in \mathbf{B}$ with $\rho(c) \geq \rho(b)$.

The fact that these two properties are indeed equivalent was established in [ADL, Th. 2.2] (under slightly more restrictive hypotheses than here), extending the earlier work of Dlab [Dla1]. We give a self-contained proof in Theorem 3.5; see also §6.1 for some elementary examples. An equivalent statement is as follows.

Theorem 1.1 (Dlab, ...). *Let \mathcal{R} be a finite Abelian category equipped with a stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ and $\varepsilon : \Lambda \rightarrow \{\pm\}$ be a sign function. Then \mathcal{R} is ε -stratified if and only if \mathcal{R}^{op} is $(-\varepsilon)$ -stratified.*

If the stratification function $\rho : \mathbf{B} \rightarrow \Lambda$ is a bijection, i.e., each stratum \mathcal{R}_λ has a unique irreducible object (up to isomorphism), then we can use ρ to identify \mathbf{B} with Λ , and denote the various distinguished objects simply by $L(\lambda), P(\lambda), \Delta_\varepsilon(\lambda), \dots$ for $\lambda \in \Lambda$ instead of by $L(b), P(b), \Delta_\varepsilon(b), \dots$ for $b \in \mathbf{B}$. When $(P\Delta_\varepsilon)$ – $(I\nabla_\varepsilon)$ hold in this situation, we instead call \mathcal{R} an ε -highest weight category with weight poset (Λ, \leq) and labelling function L . The notion of ε -highest weight category generalizes the original notion of highest weight category from [CPS1]: a (finite) highest weight category in the sense of *loc. cit.* is an ε -stratified category in which each stratum \mathcal{R}_λ is actually simple, i.e., equivalent to Vec_{fd} . This stronger assumption means not only that ρ is a bijection but also that $L_\lambda(\lambda) = P_\lambda(\lambda) = I_\lambda(\lambda)$, hence, $\Delta(\lambda) = \bar{\Delta}(\lambda)$ and $\nabla(\lambda) = \bar{\nabla}(\lambda)$ for each $\lambda \in \Lambda$. Consequently, in highest weight categories, the sign function ε plays no role and may be omitted entirely, and the above properties simplify to the following:

($P\Delta$) Each $P(\lambda)$ has a Δ -flag with sections $\Delta(\mu)$ for $\mu \geq \lambda$.

($I\nabla$) Each $I(\lambda)$ has a ∇ -flag with sections $\nabla(\mu)$ for $\mu \geq \lambda$.

In fact, in this context, the equivalence of $(P\Delta)$ and $(I\nabla)$ was established already in [CPS1]. Moreover, in *loc. cit.*, it is shown that $A\text{-mod}_{\text{fd}}$ is a highest weight category if and only if A is a quasi-hereditary algebra.

The next important special cases arise when ε is the constant function $+$ or $-$. The idea of a $+$ -stratified category originated in the work of Dlab [Dla1] already mentioned, and in another work of Cline, Parshall and Scott [CPS2]. In particular, the “standardly stratified categories” of [CPS2, Def. 2.2.1] are $+$ -stratified categories.

We say that a finite Abelian category \mathcal{R} equipped with a stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ is a *fully stratified category* if it is both a $+$ -stratified category and a $-$ -stratified category; in that case, it is ε -stratified for all choices of the sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$. Such categories arise as categories of modules over the fully stratified algebras introduced in a remark after [ADL, Def. 1.3]. In fact, these sorts of algebras and categories have appeared several times elsewhere in the literature but under different names: they are called “weakly properly stratified” in [Fri1], “exactly properly stratified” in [CouZ], and “standardly stratified” in [LW]. The latter seems a particularly confusing choice since it clashes with the established notion from [CPS2] but we completely agree with the sentiment of [LW, Rem. 2.2]: fully stratified categories have a well-behaved structure theory. One reason for this is that all of the standardization and costandardization functors in a fully stratified category are exact. We note also that any ε -stratified category with duality is automatically fully stratified; see Corollary 3.21 for a precise statement.

We use the language *fibered highest weight category* in place of fully stratified category when the stratification function ρ is a bijection. Equivalently, a fibered highest weight category is a category which is ε -highest weight for all choices of the sign function ε . Such categories arise as the categories of finite-dimensional modules over the *properly stratified algebras* introduced in [Dla2]. It is perhaps worth pointing out that any finite Abelian category can be given the structure of a fully stratified category in a trivial way

Finite-dimensional algebra A	Finite Abelian category $A\text{-mod}_{\text{fd}}$
Quasi-hereditary algebra	Highest weight category
ε -Quasi-hereditary algebra	ε -Highest weight category
Properly stratified algebra	Fibered highest weight category
ε -Stratified algebra	ε -Stratified category
Stratified algebra	Fully stratified category

TABLE 1. Dictionary between algebras and categories

taking the poset Λ to be a singleton. Fibered highest weight categories are at the other extreme with Λ being as big as possible.

Table 1 gives a dictionary between the various different types of finite Abelian category \mathcal{R} discussed so far and the language we adopt for the underlying finite-dimensional algebras A such that \mathcal{R} is equivalent to $A\text{-mod}_{\text{fd}}$. Some of this language is non-standard; see Remark 3.8 for further discussion.

There are many classical examples of highest weight categories, including blocks of the BGG category \mathcal{O} for a semisimple Lie algebra, the classical Schur algebra and Donkin's generalized Schur algebras introduced in [Don2], and many more examples arising from categories of perverse sheaves with stratifications of geometric origin [BBD]. Further examples of fully stratified categories and fibered highest weight categories which are not highest weight arise in the context of categorification. This includes the pioneering examples of categorified tensor products of finite dimensional irreducible representations for the quantum group attached to \mathfrak{sl}_k from [FKS] (in particular Remark 2.5 therein), and the categorified induced cell modules for Hecke algebras from [MS, 6.5]. Building on these examples and the subsequent work of Webster [Web1], [Web2], Losev and Webster [LW] formulated the important axiomatic definition of a *tensor product categorification*. These are fully stratified categories which have been used to give a categorical interpretation of Lusztig's construction of tensor product of based modules for a quantum group.

The device of incorporating the sign function ε into the definition of ε -stratified or ε -highest weight category seems to be quite convenient as it streamlines many of the subsequent definitions and proofs. It also leads to some interesting new possibilities when it comes to the “tilting theory” which we discuss next.

Assume \mathcal{R} is an ε -stratified category as above. An ε -tilting object is an object of \mathcal{R} which has both a Δ_ε -flag and a ∇_ε -flag. Isomorphism classes of indecomposable ε -tilting objects are parametrized in a canonical way by the set \mathbf{B} ; see Theorem 4.2. The construction of these objects is a non-trivial generalization of Ringel's classical construction via iterated extensions of standard objects: in general one takes a mixture of extensions of standard objects on the top for positive strata and extensions of costandard objects on the bottom for negative strata. We denote the indecomposable ε -tilting objects by $\{T_\varepsilon(b) \mid b \in \mathbf{B}\}$.

Now let T be an ε -tilting generator, i.e., an ε -tilting object in which every $T_\varepsilon(b)$ appears at least once as a summand. If $\varepsilon = +$ or $-$ (the constant functions) then T is a *tilting* or *cotilting module*, respectively, for the underlying finite-dimensional algebra in the general sense of tilting theory; for more general ε , T is an example of a *Wakamatsu tilting module* as defined in [Rei, §4.1]. The *Ringel dual* of \mathcal{R} relative to T is the category $\mathcal{R}' := B\text{-mod}_{\text{fd}}$ where $B := \text{End}_{\mathcal{R}}(T)^{\text{op}}$ (so that T is a right B -module). The isomorphism classes of irreducible objects in \mathcal{R}' are in natural bijection with the isomorphism classes of indecomposable summands of T , hence, they may be indexed by the same set \mathbf{B} that labels the irreducibles in \mathcal{R} . We denote them by $\{L'(b) \mid b \in \mathbf{B}\}$. Let

$$F := \text{Hom}_{\mathcal{R}}(T, ?) : \mathcal{R} \rightarrow \mathcal{R}',$$

$$G := \text{Cohom}_{\mathcal{R}}(T, ?) = \text{Hom}_{\mathcal{R}}(?, T)^* : \mathcal{R} \rightarrow \mathcal{R}'.$$

These are the *Ringel duality functors*. The following theorem is well known for highest weight categories (where it is due to Ringel [Rin] and Happel [Hap]) and for $+/-$ and $-$ -stratified categories (where it is developed in the framework of standardly stratified algebras in [AHLU]). We prove it for general ε -stratified categories in Theorem 4.10.

Theorem 1.2 (Ringel, Happel, ...). *Let \mathcal{R}' be the Ringel dual of \mathcal{R} relative to an ε -tilting generator T as above. Let $-\varepsilon : \Lambda \rightarrow \{\pm\}$ be the negation of the original sign function ε .*

- (1) *The quintuple $(\mathbf{B}, L', \rho, \Lambda, \geq)$ is a stratification of \mathcal{R}' making it into a $(-\varepsilon)$ -stratified category with weight poset (Λ, \geq) , that is, the opposite of the poset used for \mathcal{R} . Moreover, each stratum $\mathcal{R}'_\lambda := \mathcal{R}'_{\geq \lambda} / \mathcal{R}'_{> \lambda}$ of \mathcal{R}' is equivalent to the corresponding stratum $\mathcal{R}_\lambda := \mathcal{R}_{\leq \lambda} / \mathcal{R}_{< \lambda}$ of \mathcal{R} .*
- (2) *The functor F defines an equivalence of categories between the category of $\nabla_{-\varepsilon}$ -filtered objects in \mathcal{R} and the category of $\Delta_{-\varepsilon}$ -filtered objects in \mathcal{R}' . It sends ε -tilting objects (resp., injective objects) in \mathcal{R} to projective objects (resp., $(-\varepsilon)$ -tilting objects) in \mathcal{R}' .*
- (3) *The functor G defines an equivalence of categories between the category of Δ_{ε} -filtered objects in \mathcal{R} and the category of $\nabla_{-\varepsilon}$ -filtered objects in \mathcal{R}' . It sends ε -tilting objects (resp., projective objects) in \mathcal{R} to injective objects (resp., $(-\varepsilon)$ -tilting objects) in \mathcal{R}' .*
- (4) *Assume that \mathcal{R}_λ is of finite global dimension for all strata λ with $\varepsilon(\lambda) = -$ (resp., $\varepsilon(\lambda) = +$). Then the total derived functor $\mathbb{R}F : D^b(\mathcal{R}) \rightarrow D^b(\mathcal{R}')$ (resp., $\mathbb{L}G : D^b(\mathcal{R}) \rightarrow D^b(\mathcal{R}')$) is an equivalence between the bounded derived categories.*

In the setup of the theorem, let P be a projective generator for \mathcal{R} . Then $T' := GP$ is a $(-\varepsilon)$ -tilting generator for \mathcal{R}' such that $A := \text{End}_{\mathcal{R}}(P)^{\text{op}} \cong \text{End}_{\mathcal{R}'}(T')^{\text{op}}$. Since \mathcal{R} is equivalent to $A\text{-mod}_{\text{fd}}$, this shows that \mathcal{R} is equivalent to the Ringel dual $(\mathcal{R}')'$ of \mathcal{R}' relative to T' . Thus, the original category \mathcal{R} can be recovered from its Ringel dual \mathcal{R}' . This statement can be interpreted as a *double centralizer property*: starting from $\mathcal{R} = A\text{-mod}_{\text{fd}}$ so that T is an (A, B) -bimodule, and taking the projective generator P to be the left regular A -module so that $A \cong \text{End}_A(P)^{\text{op}}$, the (B, A) -bimodule $T' = GP$ is isomorphic to the dual T^* of T . Now Theorem 1.2(3) implies that $A \cong \text{End}_B(T^*)^{\text{op}}$.

We do not consider here derived equivalences in the case of infinite global dimension, but instead refer to [PS], where this and involved t -structures are treated in detail by generalizing the classical theory of co(resolving) subcategories. This requires the use of certain coderived and contraderived categories in place of ordinary derived categories.

Now we shift our attention to the semi-infinite case, which is really the main topic of the article. Following [EGNO], a *locally finite Abelian category* is a category that is equivalent to the category $\text{comod}_{\text{fd}}\text{-}C$ of finite-dimensional right comodules over some coalgebra C . Let \mathcal{R} be such a category. A *lower finite stratification* of \mathcal{R} is a quintuple $(\mathbf{B}, L, \rho, \Lambda, \leq)$ consisting of a set \mathbf{B} , a function L labelling a full set $\{L(b) \mid b \in \mathbf{B}\}$ of pairwise inequivalent irreducible objects, a stratification function $\rho : \mathbf{B} \rightarrow \Lambda$ required now to have finite fibers $\mathbf{B}_\lambda := \rho^{-1}(\lambda)$, and a lower finite poset (Λ, \leq) (i.e., the intervals $(-\infty, \mu]$ are finite for all $\mu \in \Lambda$). Fix also a sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$. For any lower set (i.e., ideal of the poset) Λ^\downarrow in Λ , we can consider the Serre subcategory \mathcal{R}^\downarrow of \mathcal{R} generated by the objects $\{L(b) \mid b \in \mathbf{B}^\downarrow\}$ where $\mathbf{B}^\downarrow := \rho^{-1}(\Lambda^\downarrow)$. The restriction of the stratification of \mathcal{R} gives a stratification $(\mathbf{B}^\downarrow, L, \rho, \Lambda, \leq)$ of \mathcal{R}^\downarrow . We say that \mathcal{R} is a *lower finite ε -stratified category* if \mathcal{R}^\downarrow is a finite Abelian category that is ε -stratified in the earlier sense for every finite lower set Λ^\downarrow of Λ ; cf. Definition 3.50. By the same procedure one also obtains definitions of lower finite ε -highest weight, lower finite fully stratified, lower finite fibered highest weight, and lower finite highest weight categories.

In a lower finite ε -stratified category \mathcal{R} , there are ε -standard and ε -costandard objects $\Delta_\varepsilon(b)$ and $\nabla_\varepsilon(b)$; they are the same as the ε -standard and ε -costandard objects of the Serre subcategory \mathcal{R}^\downarrow defined from any finite lower set Λ^\downarrow containing $\rho(b)$. As well as

(finite) Δ_ε - and ∇_ε -flags, one can consider certain infinite ∇_ε -flags in objects of the ind-completion $\text{Ind}(\mathcal{R})$ (which is the category $\text{comod-}C$ of all right C -comodules in the case that $\mathcal{R} = \text{comod}_{\text{fd}}C$). We refer to these as *ascending ∇_ε -flags*; see Definition 3.52 for the precise formulation. Theorem 3.56 establishes a homological criterion for an object to possess an ascending ∇_ε -flag similar to the well-known criterion for good filtrations in rational representations of reductive groups [Jan1, Prop. II.4.16]. From this, it follows that the injective hull $I(b)$ of $L(b)$ in $\text{Ind}(\mathcal{R})$ has an ascending ∇_ε -flag. Moreover, the multiplicity of $\nabla_\varepsilon(c)$ as a section of such a flag satisfies

$$(I(b) : \nabla_\varepsilon(c)) = [\Delta_\varepsilon(c) : L(b)],$$

generalizing BGG reciprocity. This leads to alternative “global” characterizations of lower finite ε -stratified and fully stratified categories; see Theorems 3.60 and 3.63. The latter involves an Ext^2 -vanishing condition which first appeared in work of Dlab and Ringel [DR].

In a lower finite ε -stratified category, there are also ε -tilting objects. Isomorphism classes of the indecomposable ones are labelled by \mathbf{B} just like in the finite case. In fact, for $b \in \mathbf{B}$ the corresponding indecomposable ε -tilting object of \mathcal{R} is the same as the object $T_\varepsilon(b)$ of the Serre subcategory \mathcal{R}^\downarrow defined from any finite lower set Λ^\downarrow containing $\rho(b)$. By an *ε -tilting generator* for \mathcal{R} , we now mean an object $T = \bigoplus_{i \in I} T_i \in \text{Ind}(\mathcal{R})$ with a given decomposition as a direct sum of ε -tilting objects $T_i \in \mathcal{R}$ such that each $T_\varepsilon(b)$ appears at least once as a summand of T . Then the *Ringel dual* \mathcal{R}' of \mathcal{R} relative to T is the category $A\text{-mod}_{\text{fd}}$ of locally finite-dimensional left modules over the locally finite-dimensional locally unital algebra

$$A = \left(\bigoplus_{i,j \in I} \text{Hom}_{\mathcal{R}}(T_i, T_j) \right)^{\text{op}},$$

where the op denotes that multiplication in A is the opposite of composition in \mathcal{R} ; see Definition 4.24. Saying that A is *locally unital* means that $A = \bigoplus_{i,j \in I} e_i A e_j$ where $\{e_i | i \in I\}$ are the mutually orthogonal idempotents defined by the identity endomorphisms of each T_i , and *locally finite-dimensional* means that $\dim e_i A e_j < \infty$ for all $i, j \in I$. A *locally finite-dimensional module* is an A -module $V = \bigoplus_{i \in I} e_i V$ with $\dim e_i V < \infty$ for each i . As $e_i A e_j = \text{Hom}_{\mathcal{R}}(T_i, T_j)$ is finite-dimensional, each left ideal $A e_j$ is a locally finite-dimensional projective module.

This brings us to the notion of an *upper finite ε -stratified category*, whose definition may be discovered by considering the nature of the categories \mathcal{R}' that can arise as Ringel duals of lower finite ε -stratified categories. We refer to Definition 3.34 for the intrinsic formulation; there are also upper finite counterparts of ε -highest weight, fully stratified, fibered highest weight and highest weight categories. Starting from \mathcal{R} that is a lower finite ε -stratified category as above, the Ringel dual \mathcal{R}' comes equipped with an *upper finite stratification* $(\mathbf{B}, L', \rho, \Lambda, \succ)$ making it into an upper finite $(-\varepsilon)$ -stratified category; see Theorem 4.25 which extends parts (1) and (2) of Theorem 1.2.

In general, in an upper finite ε -stratified category, the underlying poset is required to be upper finite, i.e., all of the intervals $[\lambda, \infty)$ are finite. There are ε -standard and ε -costandard objects, but now these can have infinite length (although composition multiplicities in such objects are finite). On the other hand, the indecomposable projectives and injectives do still have finite Δ_ε -flags and ∇_ε -flags, exactly like in $(P\Delta_\varepsilon)$ and $(I\nabla_\varepsilon)$. Perhaps the most interesting feature is that one can still make sense of ε -tilting objects. These are objects possessing certain infinite flags: both an *ascending Δ_ε -flag* and a *descending ∇_ε -flag*; see Definition 3.35. This allows us to define the *Ringel dual* of an upper finite ε -stratified category relative to an ε -tilting generator T : it is the category $\text{comod}_{\text{fd}}C$ for the coalgebra $C := \text{Coend}_{\mathcal{R}}(T)$ that is the continuous dual of the opposite endomorphism algebra $B := \text{End}_{\mathcal{R}}(T)^{\text{op}}$; see Theorem 4.27 which extends parts (1)

and (3) of Theorem 1.2. This makes sense because B is a pseudo-compact topological algebra; see Lemma 2.10.

Again there are double centralizer properties. For \mathcal{R}' arising as the Ringel dual of a lower finite ε -stratified category \mathcal{R} relative to $T = \bigoplus_{i \in I} T_i$, the indecomposable $(-\varepsilon)$ -tilting objects in \mathcal{R}' are the images of the indecomposable injective objects of \mathcal{R} under

$$F := \bigoplus_{i \in I} \text{Hom}_{\mathcal{R}}(T_i, ?) : \mathcal{R} \rightarrow \mathcal{R}'$$

and, given a $(-\varepsilon)$ -tilting generator T' for \mathcal{R}' , the Ringel dual $(\mathcal{R}')'$ of \mathcal{R}' relative to T' is equivalent to the original category \mathcal{R} ; see Corollary 4.29 and also §6.2 for an explicit example. Similarly, for \mathcal{R}' arising as the Ringel dual of an upper finite ε -stratified category relative to T , the indecomposable $(-\varepsilon)$ -tilting objects of \mathcal{R}' are the images of the indecomposable projective objects of \mathcal{R} under $G := \text{Cohom}_{\mathcal{R}}(T, ?)$ and, given a $(-\varepsilon)$ -tilting generator $T' = \bigoplus_{i \in I} T'_i$ for \mathcal{R}' , the Ringel dual $(\mathcal{R}')'$ of \mathcal{R}' relative to T' is equivalent to \mathcal{R} ; see Corollary 4.30.

In §5.1, we apply semi-infinite Ringel duality together with arguments from [AST] to give an elementary algebraic characterization of upper finite highest weight categories in terms of upper finite *based quasi-hereditary algebras*. In the finite-dimensional setting, these are the based quasi-hereditary algebras defined by Kleshchev and Muth in [KM], who proved that their definition of based quasi-hereditary algebra is equivalent to the original definition of quasi-hereditary algebra from [CPS1]; we have streamlined the definition a little further here. Our more general algebras are locally finite-dimensional locally unital algebras rather than unital algebras. Viewing them instead as finite-dimensional categories, that is, small \mathbb{k} -linear categories with finite-dimensional morphism spaces, the definition translates into something equivalent to the notion of an *object-adapted cellular category* which was introduced already by Elias and Lauda [ELau, Def. 2.1]. (In turn, the Elias-Lauda definition evolved from work of Westbury [Wes], who extended the definition of cellular algebra due to Graham and Lehrer [GL] from finite-dimensional algebras to finite-dimensional categories.)

We say that a fully stratified category is *tilting-rigid* if there is a bijection $\nu : \mathbf{B} \rightarrow \mathbf{B}$ such that $T_+(b) \cong T_-(\nu(b))$ for all $b \in \mathbf{B}$; see Definition 4.36. In the finite case, \mathcal{R} is tilting-rigid if and only if it is Gorenstein with strata that are quasi-Frobenius (then ν encodes their Nakayama permutations); see Theorem 4.39 which generalizes [CM, Th. 2.2]. The situation is even better if in addition all of the strata are symmetric, since in that case the tilting objects $T_\varepsilon(b)$ are isomorphic for all choices of the sign function ε so that they may all be denoted by $T(b)$. Most of the naturally-occurring examples of fully stratified categories are tilting-rigid with symmetric strata, including the tensor product categorifications from [LW] mentioned earlier. For us, the key point about the tilting-rigid hypothesis is that the Ringel dual of a tilting-rigid fully stratified category is again a tilting-rigid fully stratified category; see Theorem 4.42. This is important in §5.3, when we introduce notions of *based stratified algebras* and *based properly stratified algebras*; see Definitions 5.20 and 5.21. These have a similar flavor to the fibered object-adapted cellular categories of [ELau, Def. 2.17]. We show that the category of locally finite-dimensional modules over an upper finite based stratified algebra (resp., upper finite based properly stratified algebra) is an upper finite fully stratified (resp., fibered highest weight) category, and conversely any such category which is also tilting-rigid with symmetric strata can be realized in this way.

The definition of an upper finite based stratified algebra A involves certain basic finite-dimensional algebras A_λ ($\lambda \in \Lambda$) which provide explicit realizations of the strata. Their direct sum $\bigoplus_{\lambda \in \Lambda} A_\lambda$ is a locally unital algebra which plays the role of “Cartan subalgebra”, although in general it is not a subalgebra of A . The assumption that the algebras A_λ are basic can in fact be dropped entirely. On doing that one obtains a weaker notion which we call an algebra with a *triangular basis*; see Definition 5.26. Our




TABLE 2. Upper finite algebras and categories

understanding of this definition was influenced by the recent preprint [GRS] in which the authors introduce the closely-related notion of an algebra with a *weak triangular decomposition*; up to a choice of basis, this is the same as an algebra with a triangular basis in our sense in which all distinguished idempotents are special. It is still the case that the category of locally finite-dimensional modules over such an algebra is an upper finite fully stratified category, just like for based stratified algebras. This observation is due to Gao, Rui and Song [GRS, Th. 3.5]; we give a slightly different proof in Theorem 5.28. Gao, Rui and Song also discuss some interesting examples arising from cyclotomic quotients of the affine Brauer and oriented Brauer categories and their q -analogs.

For many of the naturally occurring algebras A with a triangular basis, the upper and lower halves of the basis span a pair of *opposite Borel subalgebras* A^\flat and A^\sharp ; this includes all of the level one cyclotomic quotients from [GRS] but not the ones of higher level. In Definition 5.31, we formalize this idea with the final notion of an algebra with a *triangular decomposition*. The first author came upon essentially this definition originally from considerations involving the oriented Brauer category and its q -analog; see [Rey], [Bru] and also [RS], which applies a similar approach in the context of the Brauer category. A closely related notion of *triangular category* was developed independently by Sam and Snowden [SS] in order to study these and other examples; see also [CouZ]. In the presence of a triangular decomposition, the “Cartan subalgebra” $\bigoplus_{\lambda \in \Lambda} A_\lambda$ may be identified with $A^\circ := A^\flat \cap A^\sharp$, so that now it is actually a subalgebra of A , and the standardization/costandardization functors can be realized as parabolic induction/coinduction functors. In Theorem 5.35, we explain a general construction to make any algebra with a triangular decomposition into a based stratified algebra. If A° is semisimple, as is the case for the examples arising from the (oriented) Brauer category in characteristic zero but not in positive characteristic, this produces a based quasi-hereditary algebra. There are other advantages to having a triangular decomposition rather than merely a triangular basis, e.g., see [SS] where triangular decompositions are used to show that many of the motivating examples are Noetherian.

Table 2 summarizes some of the connections established between these various types of algebras and their module categories. In the main body of the text, we also discuss a parallel situation involving *essentially finite* rather than upper finite algebras

and categories. For example, the finite-dimensional graded algebras with a triangular decomposition studied in [HN], [BT] fit naturally into our more general framework of algebras with an essentially finite triangular decomposition; see Remark 5.33.

As we have already mentioned, the category $\mathcal{R} := \mathcal{R}ep(G)$ for a reductive group G is the archetypical example of a lower finite highest weight category. Its Ringel dual \mathcal{R}' is an upper finite highest weight category. This case has been studied in particular by Donkin (e.g., see [Don2], [Don3]), but Donkin's approach involves truncating to a finite-dimensional algebra from the outset. The double centralizer property allowing \mathcal{R} to be reconstructed from \mathcal{R}' in this case can be interpreted as a shadow of the Tannakian formalism; see Theorem 6.11. Other important examples of semi-infinite Ringel duality come from blocks of category \mathcal{O} over an affine Lie algebra: in negative levels one obtains lower finite highest weight categories, while positive levels produce the upper finite ones which are their Ringel duals. These and several other prominent examples are outlined in §§6.3–6.7.

We would finally like to remark that our semi-infinite versions of highest weight categories should not be confused with the affine highest weight categories of [Kle], and our based quasi-hereditary algebras are not affine quasi-hereditary algebras in the sense of [Kle]. The latter are special examples of affine cellular algebras introduced in [Xi], [KX]. They are not covered by our setup since we require that strata can be realized by finite-dimensional algebras over an algebraically closed field. To incorporate them, one would need to develop the theory here over more general commutative ground rings as suggested in Remark 5.7.

Acknowledgements. The first author would like to thank Ben Elias, Alexander Kleshchev and Ivan Losev for many illuminating discussions. In particular, the fact that Ringel duality could be extended to upper finite highest weight categories was originally explained to this author by Losev. The second author would like to thank Henning Andersen, Shrawan Kumar and Wolfgang Soergel for several useful discussions on topics related to this paper. The authors also thank Tomoyuki Arakawa, Peter Fiebig and Julian Külshammer for helpful comments, and Kevin Coulembier for pointing out a mistake in the treatment of ind-completions in §2.3 of the first version of this article.

2. SOME FINITENESS PROPERTIES ON ABELIAN CATEGORIES

We fix an algebraically closed field \mathbb{k} . All algebras, categories, functors, etc. will be assumed to be linear over \mathbb{k} . We write \otimes for $\otimes_{\mathbb{k}}$. The naive terms *direct limit* and *inverse limit* will be used for small filtered colimits and limits, respectively. We begin by introducing some language for Abelian categories with various finiteness properties; see Table 3.

2.1. Finite and locally finite Abelian categories. According to [EGNO, Def. 1.8.5], a *finite Abelian category* is a category that is equivalent to the category $A\text{-mod}_{\text{fd}}$ of finite-dimensional (left) modules over some finite-dimensional algebra A . We refer to a choice for the algebra A here as an *algebra realization* of \mathcal{R} . Note that the opposite category is also a finite Abelian category as it is equivalent to the category $A^{\text{op}}\text{-mod}_{\text{fd}} = \text{mod}_{\text{fd}}\text{-}A$ due to the existence of the contravariant equivalence

$$?^* : A\text{-mod}_{\text{fd}} \rightarrow \text{mod}_{\text{fd}}\text{-}A \quad (2.1)$$

taking a finite-dimensional left A -module to the linear dual viewed as a right A -module in the natural way.

A finite Abelian category can also be characterized as a category which is equivalent to the category $\text{comod}_{\text{fd}}\text{-}C$ of finite-dimensional (right) comodules over some finite-dimensional coalgebra C . To explain this in more detail, recall that the dual $A := C^*$ of a finite-dimensional coalgebra C has a natural algebra structure with multiplication




TABLE 3. Finiteness properties

$A \otimes A \rightarrow A$ that is the dual of the comultiplication $C \rightarrow C \otimes C$; for this, one needs to use the canonical isomorphism

$$C^* \otimes C^* \rightarrow (C \otimes C)^*, \quad f \otimes g \mapsto (v \otimes w \mapsto f(v)g(w)) \quad (2.2)$$

to identify $C^* \otimes C^*$ with $(C \otimes C)^*$. Then any right C -comodule can be viewed as a left A -module with action defined from $av := \sum_{i=1}^n a(c_i)v_i$ assuming here that the structure map $\eta : V \rightarrow V \otimes C$ sends $v \mapsto \sum_{i=1}^n v_i \otimes c_i$. Conversely, the C -comodule structure on V can be recovered uniquely from the action of A . Thus, the categories $\text{comod}_{\text{fd}} C$ and $A\text{-mod}_{\text{fd}}$ are isomorphic.

A *locally finite Abelian category* is a category \mathcal{R} that is equivalent to $\text{comod}_{\text{fd}} C$ for a (not necessarily finite-dimensional) coalgebra C . We refer to a choice of C as a *coalgebra realization* of \mathcal{R} . The following result of Takeuchi gives an intrinsic characterization of locally finite Abelian categories; see [Tak] and [EGNO, Th. 1.9.15]. It is a version of [Gab, Th. IV.4] adapted to our situation. Note Takeuchi’s original paper uses the language “locally finite Abelian” slightly differently (following [Gab]) but his formulation of the result is equivalent to the one here (which follows [EGNO, Def. 1.8.1]). In *loc. cit.* it is shown moreover that C can be chosen so that it is *pointed*, i.e., all of its irreducible comodules are one-dimensional; in that case, C is unique up to isomorphism.

Lemma 2.1. *An essentially small category \mathcal{R} is a locally finite Abelian category if and only if it is Abelian, all of its objects are of finite length, and all of its morphism spaces are finite-dimensional.*

In view of Lemma 2.1, one could also define a locally finite Abelian category to be a category that is equivalent to $A\text{-mod}_{\text{fd}}$ for a (not necessarily finite-dimensional) unital algebra A , but we prefer to work in terms of comodules since this language facilitates the passage to the ind-completion. To explain this in more detail, consider the locally finite Abelian category

$$\mathcal{R} = \text{comod}_{\text{fd}} C.$$

Fix a full set of pairwise inequivalent irreducible objects $\{L(b) \mid b \in \mathbf{B}\}$ in \mathcal{R} . By Schur’s Lemma, we have that $\text{End}_{\mathcal{R}}(L(b)) = \mathbb{k}$ for each $b \in \mathbf{B}$. Note that the opposite category \mathcal{R}^{op} is again a locally finite Abelian category, and a coalgebra realization for it is given by the opposite coalgebra C^{cop} . This follows because there is a contravariant equivalence

$$?^* : \text{comod}_{\text{fd}} C \rightarrow C\text{-comod}_{\text{fd}} \quad (2.3)$$

sending a finite-dimensional right comodule to the dual vector space viewed as a left comodule in the natural way: if v_1, \dots, v_n is a basis for V , with dual basis f_1, \dots, f_n for V^* , and the structure map $V \rightarrow V \otimes C$ sends $v_j \mapsto \sum_{i=1}^n v_i \otimes c_{i,j}$ then the dual’s structure map $V^* \rightarrow C \otimes V^*$ sends $f_i \mapsto \sum_{j=1}^n c_{i,j} \otimes f_j$. Since we have that $C\text{-comod}_{\text{fd}} \cong \text{comod}_{\text{fd}} C^{\text{cop}}$, we deduce that \mathcal{R}^{op} is equivalent to $\text{comod}_{\text{fd}} C^{\text{cop}}$.

In general, \mathcal{R} need not have enough injectives or projectives. To get injectives, we pass to the *ind-completion* $\text{Ind}(\mathcal{R})$; see e.g. [KS, §6.1]. For $V, W \in \text{Ind}(\mathcal{R})$, we write $\text{Ext}_{\mathcal{R}}^n(V, W)$, or sometimes $\text{Ext}_C^n(V, W)$, for $\text{Ext}_{\text{Ind}(\mathcal{R})}^n(V, W)$; it may be computed via an injective resolution of W in the ind-completion. This convention is unambiguous due to [KS, Th. 15.3.1]; see also [Cou3, Th. 2.2.1]. One can also consider the right derived functors $\mathbb{R}^n F$ of a left exact functor $F : \text{Ind}(\mathcal{R}) \rightarrow \mathcal{R}'$ to an Abelian category \mathcal{R}' .

Let $\text{comod-}C$ be the category of all right C -comodules. Every comodule is the union (hence, the direct limit) of its finite-dimensional subcomodules. Moreover, a comodule V is compact, i.e., the functor $\text{Hom}_C(V, ?)$ commutes with direct limits, if and only if it is finite-dimensional. Using this, [KS, Cor. 6.3.5] implies that the canonical functor $\text{Ind}(\mathcal{R}) \rightarrow \text{comod-}C$ is an equivalence of categories. This means that one can work with $\text{comod-}C$ in place of $\text{Ind}(\mathcal{R})$, as we do in the next few paragraphs.

The category $\text{comod-}C$ is a Grothendieck category: it is Abelian, it possesses all small coproducts, direct colimits of monomorphisms are monomorphisms, and there is a generator. A generating family may be obtained by choosing representatives for the isomorphism classes of finite-dimensional C -comodules. By the general theory of Grothendieck categories, every C -comodule has an injective hull. We use the notation $I(b)$ to denote an injective hull of $L(b)$. The right regular comodule decomposes as

$$C \cong \bigoplus_{b \in \mathbf{B}} I(b)^{\oplus \dim L(b)}. \quad (2.4)$$

By Baer's criterion for Grothendieck categories (e.g., see [KS, Prop. 8.4.7]), arbitrary direct sums of injectives are injective. It follows that an injective hull of $V \in \text{comod-}C$ comes from an injective hull of its socle: if $\text{soc } V \cong \bigoplus_{s \in S} L(b_s)$ then $\bigoplus_{s \in S} I(b_s)$ is an injective hull of V .

In any Abelian category, we write $[V : L]$ for the *composition multiplicity* of an irreducible object L in an object V . By definition, this is the supremum of sizes of the sets $\{i = 1, \dots, n \mid V_i/V_{i-1} \cong L\}$ over all finite filtrations $0 = V_0 < V_1 < \dots < V_n = V$; possibly, $[V : L] = \infty$. Composition multiplicity is additive on short exact sequences. For any right C -comodule V , we have by Schur's Lemma that

$$[V : L(b)] = \dim \text{Hom}_C(V, I(b)). \quad (2.5)$$

When C is infinite-dimensional, the map (2.2) is not an isomorphism, but one can still use it to make the dual vector space $B := C^*$ into a unital algebra. Since C is the union of its finite-dimensional subcoalgebras, the algebra B is the inverse limit of its finite-dimensional quotients, i.e., the canonical homomorphism $B \rightarrow \varprojlim (B/J)$ is an isomorphism where the limit is over all two-sided ideals J of B of finite codimension. These two-sided ideals J form a base of neighborhoods of 0 making B into a *pseudo-compact topological algebra*; see [Gab, Ch. IV] or [Sim, Def. 2.4]. We refer to the topology on B defined in this way as the *profinite topology*. The coalgebra C can be recovered from B as the *continuous dual*

$$B^* := \{f \in B^* \mid f \text{ vanishes on some two-sided ideal } J \text{ of finite codimension}\}. \quad (2.6)$$

It has a natural coalgebra structure dual to the algebra structure on B . This is discussed further in [Sim, §3]; see also [EGNO, §1.12] where B^* is called the *finite dual*. We note that any left ideal I of B of finite codimension contains a two-sided ideal J of finite codimension, namely, $J := \text{Ann}_B(B/I)$. So, in the definition (2.6) of continuous dual, “two-sided ideal J of finite codimension” can be replaced by “left ideal I of finite codimension”. Similarly for right ideals.

Any right C -comodule V is naturally a left B -module by the same construction as in the finite-dimensional case. We deduce that the category $\text{comod-}C$ of all right C -comodules is isomorphic to the full subcategory $B\text{-mod}_{\text{ds}}$ of $B\text{-mod}$ consisting of all *discrete* left B -modules, that is, all B -modules which are the unions of their finite-dimensional submodules. In particular, $\text{comod}_{\text{fd}}\text{-}C$ and $B\text{-mod}_{\text{fd}}$ are identified under

this construction. This means that any locally finite Abelian category may be realized as the category of finite-dimensional modules over an algebra which is pseudo-compact with respect to the profinite topology; see also [Sim, §3].

The definition of the left C -comodule structure on the linear dual V^* of a right C -comodule V in (2.3) required V to be finite-dimensional in order for it to make sense. If V is an infinite-dimensional right C -comodule, it can be viewed equivalently as a discrete left module over the dual algebra $B := C^*$. Then its dual V^* is a *pseudo-compact right B -module*, that is, a B -module isomorphic to the inverse limit of its finite-dimensional quotients. Viewing pseudo-compact modules as topological B -modules with respect to the profinite topology (i.e., submodules of finite codimension form a basis of neighborhoods of 0), we obtain the category $\text{mod}_{\text{pc}}\text{-}B$ of all pseudo-compact right B -modules and continuous B -module homomorphisms. The functor (2.3) extends to

$$?^* : B\text{-mod}_{\text{ds}} \rightarrow \text{mod}_{\text{pc}}\text{-}B. \quad (2.7)$$

This is a contravariant equivalence with quasi-inverse given by the functor

$$?^* : \text{mod}_{\text{pc}}\text{-}B \rightarrow B\text{-mod}_{\text{ds}} \quad (2.8)$$

taking $V \in \text{mod}_{\text{pc}}\text{-}B$ to its *continuous dual*

$$V^* := \{f \in V^* \mid f \text{ vanishes on some submodule of } V \text{ of finite codimension}\}.$$

We are using subtly different notation here ($?^*$ vs. $?^*$), but confusion seldom arises due to context.

We record one more basic lemma about comodules over a coalgebra.

Lemma 2.2. *Suppose that C is a coalgebra and $B := C^*$ is its dual algebra. For any right C -comodule V , composing with the counit $\epsilon : C \rightarrow \mathbb{k}$ defines an isomorphism of left B -modules $\alpha_V : \text{Hom}_C(V, C) \xrightarrow{\sim} V^*$. When $V = C$, the right regular comodule, this map gives an algebra isomorphism $\text{End}_C(C)^{\text{op}} \cong B$.*

Proof. Let $\eta : V \rightarrow V \otimes C$ be the comodule structure map. To show that α_V is an isomorphism, one checks that the map $\beta_V : V^* \rightarrow \text{Hom}_C(V, C)$, $f \mapsto (f \otimes \text{id}) \circ \eta$ is its two-sided inverse; cf. [Sim, Lem. 4.9]. It remains to show that $\alpha_C : \text{End}_C(C)^{\text{op}} \xrightarrow{\sim} B$ is an algebra homomorphism: for $f, g \in B$ we have that

$$\begin{aligned} \alpha_C(\beta_C(g) \circ \beta_C(f)) &= \epsilon \circ (g \otimes \text{id}) \circ \eta \circ (f \otimes \text{id}) \circ \eta \\ &= (g \otimes \text{id}) \circ (\text{id} \otimes \epsilon) \circ \eta \circ (f \otimes \text{id}) \circ \eta = g \circ (f \otimes \text{id}) \circ \eta = fg. \quad \square \end{aligned}$$

2.2. Locally unital algebras. We are going to work with certain Abelian categories which are not locally finite, but which nevertheless have some well-behaved finiteness properties. We will define these in the next subsection. First we must review some basic notions about locally unital algebras. These ideas originate in the work of Mitchell [Mit].

A *locally unital algebra* is an associative (but not necessarily unital) algebra A equipped with a distinguished system $\{e_i \mid i \in I\}$ of mutually orthogonal idempotents such that

$$A = \bigoplus_{i,j \in I} e_i A e_j.$$

We say A is *locally finite-dimensional* if each subspace $e_i A e_j$ is finite-dimensional.

A *locally unital homomorphism* (resp., *isomorphism*) between two locally unital algebras A and B is an algebra homomorphism (resp., isomorphism) which takes distinguished idempotents to distinguished idempotents. We say that A is an *idempotent contraction* of B , or B is an *idempotent expansion* of A , if there is an algebra isomorphism $A \xrightarrow{\sim} B$ sending each distinguished idempotent in A to a sum of distinguished idempotents in B . Usually when we use this language it will be the case that $B = A$ and the isomorphism $A \rightarrow B$ is the identity function; then $A = \bigoplus_{i,j \in \hat{I}} \hat{e}_i A \hat{e}_j$ is an idempotent expansion of $A = \bigoplus_{i,j \in I} e_i A e_j$ if each of the idempotents \hat{e}_i ($i \in \hat{I}$) is a finite sum of the idempotents e_j ($j \in I$).

For a locally unital algebra A , an A -module means a left module V as usual such that $V = \bigoplus_{i \in I} e_i V$. A vector $v \in V$ is *homogeneous* if $v \in e_i V$ for some $i \in I$. A module V is

- *locally finite-dimensional* if $\dim e_i V < \infty$ for all $i \in I$;
- *finitely generated* if $V = Av_1 + \cdots + Av_n$ for vectors $v_1, \dots, v_n \in V$ (which may be assumed to be homogeneous) or, equivalently, it is a quotient of the finitely generated projective A -module $Ae_{i_1} \oplus \cdots \oplus Ae_{i_n}$ for $i_1, \dots, i_n \in I$ and $n \in \mathbb{N}$;
- *finitely presented* if there is an exact sequence

$$Ae_{j_1} \oplus \cdots \oplus Ae_{j_m} \longrightarrow Ae_{i_1} \oplus \cdots \oplus Ae_{i_n} \longrightarrow V \longrightarrow 0$$

for $i_1, \dots, i_n, j_1, \dots, j_m \in I$ and $m, n \in \mathbb{N}$.

Let $A\text{-mod}$ (resp., $A\text{-mod}_{\text{fd}}$, resp., $A\text{-mod}_{\text{fg}}$, resp., $A\text{-mod}_{\text{fp}}$) be the category of all A -modules (resp., the locally finite-dimensional ones, resp., the finitely generated ones, resp., the finitely presented ones). Similarly, we define the categories $\text{mod-}A$, $\text{mod}_{\text{fd-}}A$, $\text{mod}_{\text{fg-}}A$ and $\text{mod}_{\text{fp-}}A$ of right modules.

Remark 2.3. Any locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$ can be viewed as a category with object set I and $\text{Hom}_{\mathcal{A}}(j, i) = e_i A e_j$, with the idempotent $e_i \in A$ corresponding to the identity endomorphism $1_i \in \text{End}_{\mathcal{A}}(i)$. Conversely, any small category \mathcal{A} (\mathbb{k} -linear, of course) gives rise to a corresponding locally unital algebra A which we call the *path algebra* of \mathcal{A} . In these terms, locally finite-dimensional locally unital algebras correspond to *finite-dimensional categories*, that is, small categories all of whose morphism spaces are finite-dimensional. The notion of idempotent expansion of the algebra A becomes the notion of *thickening* of the category \mathcal{A} , which is a sort of “partial Karoubi envelope”. Also, a left A -module (resp., a locally finite-dimensional left A -module) is the same as a \mathbb{k} -linear functor from \mathcal{A} to the category $\mathcal{V}ec$ (resp., $\mathcal{V}ec_{\text{fd}}$) of vector spaces (resp., finite-dimensional vector spaces); right A -modules are functors to $\mathcal{V}ec^{\text{op}}$.

Lemma 2.4. *An essentially small category \mathcal{R} is equivalent to $A\text{-mod}$ for some locally unital algebra A if and only if \mathcal{R} is Abelian, it possesses all small coproducts, and it has a projective generating family, i.e., there is a family $(P_i)_{i \in I}$ of compact projective objects such that $V \neq 0 \Rightarrow \text{Hom}_{\mathcal{R}}(P_i, V) \neq 0$ for some $i \in I$.*

Proof. This is similar to [Fre, Ex. 5.F]. One shows that \mathcal{R} is equivalent to $A\text{-mod}$ for the locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$ defined by setting $e_i A e_j := \text{Hom}_{\mathcal{R}}(P_i, P_j)$ with multiplication that is the opposite of composition in \mathcal{R} . The canonical equivalence $\mathcal{R} \rightarrow A\text{-mod}$ is given by the functor $\bigoplus_{i \in I} \text{Hom}_{\mathcal{R}}(P_i, ?)$. \square

Lemma 2.5. *Let A be a locally unital algebra. An A -module V is compact if and only if it is finitely presented. Also, for projective modules, the notions of finitely presented and finitely generated coincide.*

Proof. This is well known for modules over a ring, and the usual proof in that setting carries over almost unchanged to the locally unital case. \square

Lemma 2.6. *Let A be a locally unital algebra. Any A -module is isomorphic to a direct limit of finitely presented A -modules.*

Proof. As any A -module is the union of its finitely generated submodules, it suffices to show that any finitely generated A -module V is isomorphic to a direct limit of finitely presented modules. But then V is a quotient of $P = Ae_{i_1} \oplus \cdots \oplus Ae_{i_n}$ by a submodule. This submodule is the union of its finitely generated submodules W , so we have that $V \cong P / \varinjlim W \cong \varinjlim P/W$. This is a direct limit of finitely presented modules. \square

The following lemma is fundamental. It is the analog of “adjointness of tensor and hom” in the locally unital setting; see e.g. [BD1, §2.1] for a fuller discussion.

Lemma 2.7. *Let $A = \bigoplus_{i,j \in I} e_i A e_j$ and $B = \bigoplus_{i,j \in J} f_i B f_j$ be locally unital algebras, and let $M = \bigoplus_{i \in I, j \in J} e_i M f_j$ be an (A, B) -bimodule.*

- (1) The functor $M \otimes_B ? : B\text{-mod} \rightarrow A\text{-mod}$ is left adjoint to $\bigoplus_{j \in J} \text{Hom}_A(M f_j, ?)$.
(2) The functor $? \otimes_A M : \text{mod-}A \rightarrow \text{mod-}B$ is left adjoint to $\bigoplus_{i \in I} \text{Hom}_B(e_i M, ?)$.

For any locally unital algebra A , there is a contravariant equivalence

$$?^{\otimes} : A\text{-mod}_{\text{lfid}} \rightarrow \text{mod}_{\text{lfid-}A} \quad (2.9)$$

sending a left module V to $V^{\otimes} := \bigoplus_{i \in I} (e_i V)^*$, viewed as a right module in the obvious way. The analogous functor $?^{\otimes} : \text{mod}_{\text{lfid-}A} \rightarrow A\text{-mod}_{\text{lfid}}$ gives a quasi-inverse. The contravariant functor (2.9) also makes sense on arbitrary left (or right) A -modules. It is no longer an equivalence, but we still have that

$$\text{Hom}_A(V, W^{\otimes}) \cong \text{Hom}_A(W, V^{\otimes}) \quad (2.10)$$

for any $V \in A\text{-mod}$ and $W \in \text{mod-}A$. To prove this, apply Lemma 2.7(1) to the (\mathbb{k}, A) -bimodule W to show that $\text{Hom}_A(V, W^{\otimes}) \cong (W \otimes_A V)^*$, then apply Lemma 2.7(2) to the (A, \mathbb{k}) -bimodule V to show that $(W \otimes_A V)^* \cong \text{Hom}_A(W, V^{\otimes})$.

Lemma 2.8. *The dual V^{\otimes} of a projective (left or right) A -module is an injective (right or left) A -module.*

Proof. Just like in the classic treatment of duality for vector spaces from [Mac, IV.2], (2.10) shows that the covariant functor $?^{\otimes} : A\text{-mod} \rightarrow (\text{mod-}A)^{\text{op}}$ is left adjoint to the exact covariant functor $?^{\otimes} : (\text{mod-}A)^{\text{op}} \rightarrow A\text{-mod}$. So it sends projective left A -modules to projectives in $(\text{mod-}A)^{\text{op}}$, which are injective right A -modules. \square

Now we assume that A is a locally unital algebra and $T \in A\text{-mod}_{\text{lfid}}$. We are going to give a self-contained account of the construction of a coalgebra $\text{Coend}_A(T)$ which is the continuous dual of the endomorphism algebra $\text{End}_A(T)^{\text{op}}$. This is the *coend construction* which is an essential ingredient in the proof of Lemma 2.1 as discussed for example in [EGNO, §1.10], although as usual we are using the language of algebras and modules rather than the language of categories and functors used there. To start with, let

$$B := \text{End}_A(T)^{\text{op}}, \quad (2.11)$$

which is a unital algebra. Then T is an (A, B) -bimodule and the dual T^{\otimes} is a (B, A) -bimodule. Let $T_i := e_i T$, so that $T = \bigoplus_{i \in I} T_i$ and $T^{\otimes} = \bigoplus_{i \in I} T_i^*$.

Lemma 2.9. *Suppose that $T = \bigoplus_{i \in I} T_i \in A\text{-mod}_{\text{lfid}}$ and $B := \text{End}_A(T)^{\text{op}}$ are as above. For any $V \in A\text{-mod}$, there is a natural isomorphism of right B -modules*

$$\text{Hom}_A(V, T) \xrightarrow{\sim} (T^{\otimes} \otimes_A V)^*, \quad \theta \mapsto (f \otimes v \mapsto f(\theta(v))). \quad (2.12)$$

In particular, taking $V = T$, we get that $(T^{\otimes} \otimes_A T)^ \cong B$ as (B, B) -bimodules.*

Proof. By Lemma 2.7 applied to the (A, \mathbb{k}) -bimodule T^{\otimes} , the functor $T^{\otimes} \otimes_A ?$ is left adjoint to $\bigoplus_{i \in I} \text{Hom}_{\mathbb{k}}(T_i^*, ?)$. Hence,

$$(T^{\otimes} \otimes_A V)^* = \text{Hom}_{\mathbb{k}}(T^{\otimes} \otimes_A V, \mathbb{k}) \cong \text{Hom}_A\left(V, \bigoplus_{i \in I} \text{Hom}_{\mathbb{k}}(T_i^*, \mathbb{k})\right) \cong \text{Hom}_A(V, T).$$

This is the natural isomorphism in the statement of the lemma. We leave it to the reader to check that it is a B -module homomorphism. \square

Continuing with this setup, let

$$C := T^{\otimes} \otimes_A T. \quad (2.13)$$

There is a unique way to make this into a coalgebra so that the bimodule isomorphism $B \xrightarrow{\sim} C^*$ from Lemma 2.9 is actually an algebra isomorphism (viewing the dual C^* of a coalgebra as an algebra as in the previous subsection). Explicitly, let $u_1^{(i)}, \dots, u_{d(i)}^{(i)}$ be a

basis for T_i and $v_1^{(i)}, \dots, v_{d(i)}^{(i)}$ be the dual basis for T_i^* . Let $c_{r,s}^{(i)} := v_s^{(i)} \otimes u_r^{(i)} \in C$. Then the comultiplication $\delta : C \rightarrow C \otimes C$ and counit $\epsilon : C \rightarrow \mathbb{k}$ satisfy

$$\delta(c_{r,s}^{(i)}) = \sum_{t=1}^{d(i)} c_{r,t}^{(i)} \otimes c_{t,s}^{(i)}, \quad \epsilon(c_{r,s}^{(i)}) = \delta_{r,s} \quad (2.14)$$

for each $i \in I$ and $1 \leq r, s \leq d(i)$. For the next lemma, recall the definition of continuous dual of a pseudo-compact topological algebra from (2.6).

Lemma 2.10. *The endomorphism algebra $B = \text{End}_A(T)^{\text{op}}$ of $T \in A\text{-mod}_{\text{lfid}}$ is a pseudo-compact topological algebra with respect to the profinite topology, i.e., B is isomorphic to $\varprojlim B/J$ where the inverse limit is over all two-sided ideals J of finite codimension. Moreover, the coalgebra C from (2.13) may be identified with the continuous dual B^* .*

Proof. This follows because $B \cong C^*$ as algebras. \square

Thus, the coalgebra C defined by (2.13) is identified with the continuous dual

$$\text{Coend}_A(T) := (\text{End}_A(T)^{\text{op}})^* \quad (2.15)$$

of B . Explicitly, using the formula (2.12), the element $c_{r,s}^{(i)} = v_s^{(i)} \otimes u_r^{(i)} \in C$ is identified with the function sending $\theta \in \text{End}_A(T)$ to $v_s(\theta(u_r))$.

Now consider the functor $T^{\otimes} \otimes_A ? : A\text{-mod} \rightarrow B\text{-mod}$. Since T is locally finite-dimensional, it takes finitely generated A -modules to finite-dimensional B -modules. Any A -module V is the union of its finitely generated submodules, and $T^{\otimes} \otimes_A ?$ commutes with direct limits, so we see that $T^{\otimes} \otimes_A V$ is actually a discrete B -module. Since $B \cong C^*$, the category $B\text{-mod}_{\text{ds}}$ is isomorphic to $\text{comod-}C$. So we have constructed a functor

$$T^{\otimes} \otimes_A ? : A\text{-mod} \rightarrow \text{comod-}C. \quad (2.16)$$

For $V \in A\text{-mod}$, the comodule structure map on $T^{\otimes} \otimes_A V$ is given explicitly by the formula

$$\eta : T^{\otimes} \otimes_A V \rightarrow T^{\otimes} \otimes_A V \otimes C, \quad v_s^{(i)} \otimes v \mapsto \sum_{r=1}^{d(i)} v_r^{(i)} \otimes v \otimes c_{r,s}^{(i)}. \quad (2.17)$$

Recall the definition of the functor $?^*$ from (2.8).

Lemma 2.11. *Suppose that $T = \bigoplus_{i \in I} T_i \in A\text{-mod}_{\text{lfid}}$, $B := \text{End}_A(T)^{\text{op}}$ and $C \cong B^*$ are as above. The functor $T^{\otimes} \otimes_A ?$ just constructed is isomorphic to*

$$G = \text{Cohom}_A(T, ?) := \text{Hom}_A(?, T)^* : A\text{-mod} \rightarrow \text{comod-}C, \quad (2.18)$$

and it is left adjoint to the functor

$$G_* = \bigoplus_{i \in I} \text{Hom}_C(T_i^*, ?) : \text{comod-}C \rightarrow A\text{-mod}. \quad (2.19)$$

Thus, (G, G_*) is an adjoint pair.

Proof. The fact that (2.16) is left adjoint to (2.19) follows by Lemma 2.7. To see that it is isomorphic to (2.18), take $V \in A\text{-mod}$ and consider the natural isomorphism $\text{Hom}_A(V, T) \cong (T^{\otimes} \otimes_A V)^*$ of right B -modules from Lemma 2.9. As $T^{\otimes} \otimes_A V$ is discrete, its dual is a pseudo-compact left B -module, hence, $\text{Hom}_A(V, T)$ is pseudo-compact too. Then we apply $*$, using that it is quasi-inverse to $*$, to get that $\text{Hom}_A(V, T)^* \in B\text{-mod}_{\text{ds}}$ is naturally isomorphic to $T^{\otimes} \otimes_A V$. \square

2.3. Schurian categories. By a *Schurian category*, we mean a category \mathcal{R} that is equivalent to $A\text{-mod}_{\text{lfid}}$ for a locally finite-dimensional locally unital algebra A . This non-standard terminology is considerably more restrictive than other usage of the same term elsewhere in the literature, where “Schurian category” is typically used to indicate a \mathbb{k} -linear category in which the endomorphism algebras of the indecomposable objects are one-dimensional¹ (e.g., see work of Roiter).

By an *algebra realization* of a Schurian category \mathcal{R} , we mean a locally finite-dimensional locally unital algebra A (together with the set I indexing its distinguished idempotents) such that \mathcal{R} is equivalent to $A\text{-mod}_{\text{lfid}}$. Now we assume that

$$\mathcal{R} = A\text{-mod}_{\text{lfid}}$$

and proceed to summarize some of the basic properties of such categories, referring to [BD1, §2] for a more detailed treatment. Let $\{L(b) \mid b \in \mathbf{B}\}$ be a full set of pairwise inequivalent irreducible objects of \mathcal{R} . Schur’s Lemma holds: we have that $\text{End}_{\mathcal{R}}(L(b)) = \mathbb{k}$ for each $b \in \mathbf{B}$. Note that the opposite category \mathcal{R}^{op} is also Schurian, and A^{op} gives an algebra realization for it. This follows because $\mathcal{R}^{\text{op}} = (A\text{-mod}_{\text{lfid}})^{\text{op}}$ is equivalent to $\text{mod}_{\text{lfid}}\text{-}A \cong (A^{\text{op}})\text{-mod}_{\text{lfid}}$ using (2.9).

Let \mathcal{R}_c be the (not necessarily Abelian) full subcategory of \mathcal{R} consisting of all compact objects, and $\text{Ind}(\mathcal{R}_c)$ be its ind-completion. The canonical functor $\text{Ind}(\mathcal{R}_c) \rightarrow A\text{-mod}$ is an equivalence of categories. To see this, we note that all finitely generated A -modules are locally finite-dimensional as A itself is locally finite-dimensional. Hence, finitely presented A -modules are locally finite-dimensional too, i.e. $A\text{-mod}_{\text{fg}}$ is a subcategory of $A\text{-mod}_{\text{lfid}}$. In view of Lemma 2.5, this is the category \mathcal{R}_c . It just remains to apply [KS, Cor. 6.3.5], using Lemma 2.6 when checking the required hypotheses.

The category $A\text{-mod}$ is a Grothendieck category. In particular, this means that every A -module has an injective hull in $A\text{-mod}$. Since every A -module is a quotient of a direct sum of projective A -modules of the form Ae_i , the category $A\text{-mod}$ also has enough projectives. It is *not* true that an arbitrary A -module has a projective cover, but we will see in Lemma 2.14 below that finitely generated A -modules do.

Like we did in §2.1, we write $\text{Ext}_{\mathcal{R}}^n(V, W)$, or sometimes $\text{Ext}_A^n(V, W)$, in place of $\text{Ext}_{\text{Ind}(\mathcal{R}_c)}^n(V, W)$ for any $V, W \in \text{Ind}(\mathcal{R}_c)$. This can be computed either from a projective resolution of V or from an injective resolution of W . We can also consider both right derived functors $\mathbb{R}^n F$ of a left exact functor $F : \text{Ind}(\mathcal{R}_c) \rightarrow \mathcal{R}'$ and left derived functors $\mathbb{L}_n G$ of a right exact functor $G : \text{Ind}(\mathcal{R}_c) \rightarrow \mathcal{R}'$. We provide an elementary proof of the following, but note it also follows from [KS, Th. 15.3.1].

Lemma 2.12. *For $V, W \in \mathcal{R}$ and $n \geq 0$, there is a natural isomorphism*

$$\text{Ext}_{\mathcal{R}}^n(V, W) \cong \text{Ext}_{\mathcal{R}^{\text{op}}}^n(W, V).$$

Proof. Using (2.9), we must show that $\text{Ext}_A^n(V, W) \cong \text{Ext}_A^n(W^{\otimes}, V^{\otimes})$ for locally finite-dimensional A -modules V and W . To compute $\text{Ext}_A^n(V, W)$, take a projective resolution

$$\cdots \longrightarrow P_1 \longrightarrow P_0 \longrightarrow V \longrightarrow 0$$

of V in $A\text{-mod}$. By Lemma 2.8, on applying the exact functor \otimes , we get an injective resolution

$$0 \longrightarrow V^{\otimes} \longrightarrow P_0^{\otimes} \longrightarrow P_1^{\otimes} \longrightarrow \cdots$$

of V^{\otimes} in $\text{mod}\text{-}A$. Since W is locally finite-dimensional, we can use (2.10) to see that $\text{Hom}_A(P_i, W) \cong \text{Hom}_A(W^{\otimes}, P_i^{\otimes})$ for each i . So $\text{Ext}_A^n(V, W) \cong \text{Ext}_A^n(W^{\otimes}, V^{\otimes})$. \square

¹Note also that the present usage is different from several recent papers of the first author: in [BD1], the phrase “locally Schurian” was used to describe the categories we now call “Schurian”; more precisely, in [BD1], a locally Schurian category referred to a category of the form $A\text{-mod}$ (rather than $A\text{-mod}_{\text{lfid}}$) for locally finite-dimensional locally unital algebras A . We could not use the phrase “Schurian” in *loc. cit.* since that was reserved for a more restrictive notion defined in [BLW, §2.1]; this more restrictive notion will be discussed in the next subsection, again using different language.

Let $I(b)$ be an injective hull of $L(b)$ in $A\text{-mod}$. The dual $(e_i A)^\otimes$ of the projective right A -module $e_i A$ is injective in $A\text{-mod}$. Since $\text{End}_A((e_i A)^\otimes)^{\text{op}} \cong \text{End}_A(e_i A) \cong e_i A e_i$, which is finite-dimensional, the injective module $(e_i A)^\otimes$ can be written as a finite direct sum of indecomposable injectives. To determine which ones, we compute its socle: we have that $\text{Hom}_A(L(b), (e_i A)^\otimes) \cong \text{Hom}_A(e_i A, L(b)^\otimes) \cong (L(b)^\otimes) e_i = (e_i L(b))^*$, hence,

$$(e_i A)^\otimes \cong \bigoplus_{b \in \mathbf{B}} I(b)^{\oplus \dim e_i L(b)}, \quad (2.20)$$

with all but finitely many summands on the right hand side being zero. In particular, this shows for fixed i that $\dim e_i L(b) = 0$ for all but finitely many $b \in \mathbf{B}$. Conversely, for fixed $b \in \mathbf{B}$, we can always choose $i \in I$ so that $e_i L(b) \neq 0$, and deduce that $I(b)$ is a summand of $(e_i A)^\otimes$. This means that each indecomposable injective $I(b)$ is a locally finite-dimensional left A -module.

Let $P(b)$ be the dual of the injective hull of the irreducible right A -module $L(b)^\otimes$. By dualizing the right module analog of the decomposition (2.20), we get also that

$$A e_i \cong \bigoplus_{b \in \mathbf{B}} P(b)^{\oplus \dim e_i L(b)}, \quad (2.21)$$

with all but finitely many summands being zero. In particular, $P(b)$ is projective in $A\text{-mod}$, hence, it is a projective cover of $L(b)$ in $A\text{-mod}$. The composition multiplicities of any A -module satisfy

$$[V : L(b)] = \dim \text{Hom}_A(V, I(b)) = \dim \text{Hom}_A(P(b), V). \quad (2.22)$$

Lemma 2.13. *For A as above, left A -module V is locally finite-dimensional if and only if $[V : L(b)] < \infty$ for all $b \in \mathbf{B}$.*

Proof. Note that V is locally finite-dimensional if and only if $\dim \text{Hom}_A(A e_i, V) < \infty$ for each $i \in I$. Using the decomposition (2.21), this is if and only if $\dim \text{Hom}_A(P(b), V) < \infty$ for each $b \in \mathbf{B}$. \square

There is a little more to be said about finitely generated modules. Recall from the previous subsection that a module is finitely generated if $V = A v_1 + \cdots + A v_n$ for homogeneous vectors $v_1, \dots, v_n \in V$. We say that V is *finitely cogenerated* if its dual is finitely generated. It is obvious from these definitions that $\text{Hom}_A(V, W)$ is finite-dimensional either if V is finitely generated and W is locally finite-dimensional, or if V is locally finite-dimensional and W is finitely cogenerated. The following checks that our naive definitions are consistent with the usual notions of finitely generated and cogenerated objects of Grothendieck categories.

Lemma 2.14. *For $V \in A\text{-mod}$, the following properties are equivalent:*

- (i) V is finitely generated;
- (ii) the radical $\text{rad } V$, i.e., the sum of its maximal proper submodules, is a superfluous submodule and $\text{hd } V := V/\text{rad } V$ is of finite length;
- (iii) V is a quotient of a finite direct sum of the modules $P(b)$ for $b \in \mathbf{B}$.

Moreover, any finitely generated V has a projective cover.

Proof. We have already observed that $P(b)$ is a projective cover of $L(b)$. The lemma follows by standard arguments given this and the decomposition (2.21). \square

Corollary 2.15. *For $V \in A\text{-mod}$, the following properties are equivalent:*

- (i) V is finitely cogenerated;
- (ii) $\text{soc } V$ is an essential submodule of finite length;
- (iii) V is isomorphic to a submodule of a finite direct sum of modules $I(b)$ for $b \in \mathbf{B}$.

We say that a locally finite-dimensional locally unital algebra $A = \bigoplus_{i, j \in I} e_i A e_j$ is *pointed* if A is a basic algebra, i.e., all of its irreducible modules are one-dimensional, and all of its distinguished idempotents $\{e_i \mid i \in I\}$ are primitive.

Lemma 2.16. *Let $A = \bigoplus_{i,j \in I} e_i A e_j$ be a locally finite-dimensional locally unital algebra. Pick an idempotent expansion $A = \bigoplus_{i,j \in \hat{I}} \hat{e}_i A \hat{e}_j$ such that for some subset $\mathbf{B} \subseteq \hat{I}$ the set $\{\hat{e}_b \mid b \in \mathbf{B}\}$ is a complete set of pairwise non-conjugate primitive idempotents in A . Let $B := \bigoplus_{a,b \in \mathbf{B}} \hat{e}_a A \hat{e}_b$. Then B is a pointed locally unital algebra that is Morita equivalent to A , and any such pointed locally unital algebra is isomorphic to B .*

Proof. It is clear that B is pointed. To see that A and B are Morita equivalent, note that the functor $A\text{-mod} \rightarrow B\text{-mod}, V \mapsto \bigoplus_{b \in \mathbf{B}} \hat{e}_b V$ is an equivalence of categories with quasi-inverse given by the functor $(\bigoplus_{b \in \mathbf{B}} A \hat{e}_b) \otimes_B ?$. Finally if B' another pointed locally unital algebra that is Morita equivalent to A , let $F : A\text{-mod} \rightarrow B'\text{-mod}$ be an equivalence of categories. Then we have that $B' = \bigoplus_{b \in \mathbf{B}} B'_b$ for left ideals $B'_b \cong F(A \hat{e}_b)$. So

$$B' \cong \left(\bigoplus_{a,b \in \mathbf{B}} \text{Hom}_{B'}(B'_a, B'_b) \right)^{\text{op}} \cong \bigoplus_{a,b \in \mathbf{B}} \text{Hom}_A(A \hat{e}_a, A \hat{e}_b) = \bigoplus_{a,b \in \mathbf{B}} \hat{e}_a A \hat{e}_b = B.$$

This proves the uniqueness. \square

Finally, we introduce some terminology which will not be needed until §5.5.

Definition 2.17. Let $A = \bigoplus_{i,j \in I} e_i A e_j$ be a locally finite-dimensional locally unital algebra. Let $\mathbf{S} \subseteq I$ be a subset. We say that a left A -module V is \mathbf{S} -free if there is a subset $X = \bigsqcup_{s \in \mathbf{S}} X(s) \subset V$ such that the following properties hold:

(LF1) $V = \bigoplus_{x \in X} Ax$.

(LF2) The homomorphism $Ae_s \rightarrow Ax, a \mapsto ax$ is an isomorphism for $x \in X(s)$.

Equivalently, there is a \mathbb{K} -submodule U of $eV := \bigoplus_{s \in \mathbf{S}} e_s V$ such that the multiplication map $Ae \otimes_{\mathbb{K}} U \rightarrow V$ is an isomorphism, where $Ae := \bigoplus_{s \in \mathbf{S}} Ae_s$ and $\mathbb{K} := \bigoplus_{s \in \mathbf{S}} \mathbb{k}e_s$.

Lemma 2.18. *Suppose that $A = \bigoplus_{i,j \in I} e_i A e_j$ is a locally finite-dimensional locally unital algebra and $\{e_b \mid b \in \mathbf{B}\}$ is a full set of pairwise non-conjugate primitive idempotents in A for some subset $\mathbf{B} \subseteq I$. Then every finitely generated projective left A -module is \mathbf{B} -free.*

Proof. Any finitely generated projective left A -module V decomposes as a finite direct sum of indecomposable projectives, and any indecomposable projective is isomorphic to Ae_b for some $b \in \mathbf{B}$. Hence, we can pick a finite subset $X = \bigsqcup_{b \in \mathbf{B}} X(b)$ so that $V = \bigoplus_{x \in X} Ax$ with $Ax \cong Ae_b$ for $x \in X(b)$. \square

There are obvious right module analogs of these notions.

2.4. Essentially finite Abelian categories. We say that a locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$ is *essentially finite-dimensional* if each right ideal $e_i A$ and each left ideal $A e_j$ is finite-dimensional. By an *essentially finite Abelian category*, we mean a category \mathcal{R} that is equivalent to $A\text{-mod}_{\text{fd}}$ for such an A . In that case, we refer to A as an *algebra realization* of \mathcal{R} . Note that \mathcal{R} is essentially finite Abelian if and only if \mathcal{R}^{op} is essentially finite Abelian. Moreover, if A is an algebra realization for \mathcal{R} then A^{op} is one for \mathcal{R}^{op} by the obvious contravariant equivalence $?^* : A\text{-mod}_{\text{fd}} \rightarrow \text{mod}_{\text{fd}}\text{-}A$.

Lemma 2.19. *An essentially small category \mathcal{R} is equivalent to $A\text{-mod}_{\text{fd}}$ for a locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$ such that each left ideal Ae_j (resp., each right ideal $e_i A$) is finite-dimensional if and only if \mathcal{R} is a locally finite Abelian category with enough projectives (resp., enough injectives).*

Proof. We just prove the result for left ideals and projectives; the parenthesized statement for right ideals and injectives follows by replacing \mathcal{R} and A with \mathcal{R}^{op} and A^{op} .

Suppose first that $A = \bigoplus_{i,j \in I} e_i A e_j$ is a locally unital algebra such that each left ideal Ae_j is finite-dimensional. Then $A\text{-mod}_{\text{fd}}$ is a locally finite Abelian category. It has enough projectives because the left ideals Ae_j are finite-dimensional.

Conversely, suppose \mathcal{R} is a locally finite Abelian category with enough projectives. Let $\{L(b) \mid b \in \mathbf{B}\}$ be a full set of pairwise inequivalent irreducible objects, and $P(b) \in \mathcal{R}$ a projective cover of $L(b)$. Define A to be the locally unital algebra $A = \bigoplus_{a,b \in \mathbf{B}} e_a A e_b$ where $e_a A e_b := \text{Hom}_{\mathcal{R}}(P(a), P(b))$ with multiplication that is the opposite of composition in \mathcal{R} . This is a pointed locally finite-dimensional locally unital algebra. As in the proof of Lemma 2.4, the functor $\bigoplus_{b \in \mathbf{B}} \text{Hom}_{\mathcal{R}}(P(b), ?)$ defines an equivalence $\mathcal{R} \rightarrow A\text{-mod}_{\text{fd}}$. It remains to note that the ideals $A e_b$ are finite-dimensional since they are the images under this functor of the projectives $P(b)$, which are of finite length. \square

Corollary 2.20. *An essentially small category \mathcal{R} is an essentially finite Abelian category if and only if it is a locally finite Abelian category with enough injectives and projectives.*

Essentially finite Abelian categories are almost as convenient to work with as finite Abelian categories since one can perform all of the usual constructions of homological algebra without needing to pass to the ind-completion.

Lemma 2.21. *For a category \mathcal{R} , the following are equivalent:*

- (i) \mathcal{R} is a finite Abelian category;
- (ii) \mathcal{R} is a Schurian category with only finitely many isomorphism classes of irreducible objects;
- (iii) \mathcal{R} is an essentially finite Abelian category with only finitely many isomorphism classes of irreducible objects;
- (iv) \mathcal{R} is a locally finite Abelian category with only finitely many isomorphism classes of irreducible objects and either enough projectives or enough injectives;
- (v) \mathcal{R} is both a locally finite Abelian category and a Schurian category.

Proof. Clearly, (i) implies (ii) and (iii). The implication (ii) \Rightarrow (i) follows on considering a pointed algebra realization of \mathcal{R} . The implication (iii) \Rightarrow (iv) follows from Corollary 2.20. The implication (iv) \Rightarrow (i) follows from Lemma 2.19. Clearly (ii) and (iv) together imply (v). Finally, to see that (v) implies (ii), it suffices to note that a Schurian category with infinitely many isomorphism classes of irreducible objects cannot be locally finite Abelian: the direct sum of infinitely many non-isomorphic irreducibles is a well-defined object of \mathcal{R} but it is not of finite length. \square

Essentially finite Abelian categories with infinitely many isomorphism classes of irreducible objects are *not* Schurian categories. However they are closely related as we explain next.

- If \mathcal{R} is essentially finite Abelian, we define its *Schurian envelope* $\text{Env}(\mathcal{R})$ to be the full subcategory of $\text{Ind}(\mathcal{R})$ consisting of all objects that have finite composition multiplicities.
- If \mathcal{R} is Schurian, let $\text{Fin}(\mathcal{R})$ be the full subcategory of \mathcal{R} consisting of all objects of finite length.

We say that a Schurian category \mathcal{R} is *Cartan-bounded* if its Cartan matrix C has only finitely many non-zero entries in every row and column, where by Cartan matrix we mean the matrix

$$(\dim \text{Hom}_{\mathcal{R}}(P(a), P(b)))_{a,b \in \mathbf{B}} = (\dim \text{Hom}_{\mathcal{R}}(I(a), I(b)))_{a,b \in \mathbf{B}}, \quad (2.23)$$

where \mathbf{B} is labelling indecomposable projectives and injectives in the usual way.

Lemma 2.22. *If \mathcal{R} is an essentially finite Abelian category then $\text{Env}(\mathcal{R})$ is a Cartan-finite Schurian category, and conversely if \mathcal{R} is a Cartan-finite Schurian category then $\text{Fin}(\mathcal{R})$ is an essentially finite Abelian category. Moreover, Env and Fin are inverses in the sense that $\text{Fin}(\text{Env}(\mathcal{R}))$ is equivalent to \mathcal{R} for any essentially finite Abelian \mathcal{R} , and $\text{Env}(\text{Fin}(\mathcal{R}))$ is equivalent to \mathcal{R} for any Cartan-finite Schurian \mathcal{R} :*

$$\left(\begin{array}{c} \text{Essentially finite} \\ \text{Abelian categories} \end{array} \right) \begin{array}{c} \xrightarrow{\text{Env}} \\ \xleftarrow{\text{Fin}} \end{array} \left(\begin{array}{c} \text{Cartan-finite} \\ \text{Schurian categories} \end{array} \right).$$

Proof. This is easy to see in terms of an algebra realization: if $\mathcal{R} = A\text{-mod}_{\text{fd}}$ for an essentially finite-dimensional locally unital algebra A then $\text{Env}(\mathcal{R}) = A\text{-mod}_{\text{fd}}$, so it is Schurian. Since the indecomposable injectives and projectives in $\text{Env}(\mathcal{R})$ are the same as in \mathcal{R} , they have finite length. Conversely, using Lemma 2.16, we may assume that $\mathcal{R} = A\text{-mod}_{\text{fd}}$ for a *pointed* locally finite-dimensional locally unital algebra, such that all of the indecomposable injectives and projectives are of finite length. Since A is pointed, this means equivalently that all of the left ideals Ae_i and right ideals e_iA are finite-dimensional. Hence, A is essentially finite-dimensional, and $\text{Fin}(\mathcal{R}) = A\text{-mod}_{\text{fd}}$ is essentially finite Abelian. \square

2.5. Recollement. We conclude the section with some reminders about “recollement” in our algebraic setting; see [BBD, §1.4] or [CPS1, §2] for further background. We need this here only for Abelian categories \mathcal{R} satisfying finiteness properties as developed above. The recollement formalism provides us with an adjoint triple of functors $(i^*, i, i^!)$ associated to the inclusion $i : \mathcal{R}^\perp \rightarrow \mathcal{R}$ of a Serre subcategory, and an adjoint triple of functors $(j_!, j, j_*)$ associated to the projection $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ onto a Serre quotient category, with the image of i being the kernel of j . These functors will play an essential role in all subsequent arguments.

First suppose that \mathcal{R} is any Abelian category. Assume that we are given a full set $\{L(b) \mid b \in \mathbf{B}\}$ of pairwise inequivalent irreducible objects. Let \mathbf{B}^\perp be a subset of \mathbf{B} and \mathcal{R}^\perp be the full subcategory of \mathcal{R} consisting of all the objects V such that $[V : L(b)] \neq 0 \Rightarrow b \in \mathbf{B}^\perp$. This is a Serre subcategory of \mathcal{R} with irreducible objects $\{L^\perp(b) \mid b \in \mathbf{B}^\perp\}$ defined by $L^\perp(b) := L(b)$.

Lemma 2.23. *In the above setup, the inclusion functor $i : \mathcal{R}^\perp \rightarrow \mathcal{R}$ has a left adjoint i^* and a right adjoint $i^!$:*

$$\begin{array}{ccc} & i^! & \\ \curvearrowright & & \curvearrowleft \\ \mathcal{R}^\perp & \xrightarrow{i} & \mathcal{R} \\ \curvearrowleft & & \curvearrowright \\ & i^* & \end{array}$$

The counit of one of these adjunctions and the unit of the other give isomorphisms:

$$i^* \circ i \xrightarrow{\sim} \text{Id}_{\mathcal{R}^\perp} \xrightarrow{\sim} i^! \circ i.$$

In particular, i is fully faithful.

Proof. This is straightforward. Explicitly, i^* (resp., $i^!$) sends an object of \mathcal{R} to the largest quotient (resp., subobject) that belongs to \mathcal{R}^\perp . \square

Now we are going to pass to the *Serre quotient* $\mathcal{R}^\dagger := \mathcal{R}/\mathcal{R}^\perp$. This is an Abelian category equipped with an exact *quotient functor* $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ satisfying the following universal property: if $h : \mathcal{R} \rightarrow \mathcal{C}$ is any exact functor to an Abelian category \mathcal{C} with $hL(b) = 0$ for all $b \in \mathbf{B}^\perp$, then there is a unique functor $\bar{h} : \mathcal{R}^\dagger \rightarrow \mathcal{C}$ such that $h = \bar{h} \circ j$. The irreducible objects in \mathcal{R}^\dagger are $\{L^\dagger(b) \mid b \in \mathbf{B}^\dagger\}$ where $\mathbf{B}^\dagger := \mathbf{B} \setminus \mathbf{B}^\perp$ and $L^\dagger(b) := jL(b)$. For a fuller discussion of these statements, see e.g. [Gab].

The quotient functor j need not have a left or a right adjoint in general, so we need to impose some additional hypotheses. We first assume that \mathcal{R} is finite Abelian, essentially finite Abelian or Schurian. Then one can understand j rather explicitly as an idempotent truncation functor and it always has both a left and right adjoint:

Lemma 2.24. *Suppose that \mathcal{R} is finite Abelian, essentially finite Abelian or Schurian, $\mathbf{B} = \mathbf{B}^\perp \sqcup \mathbf{B}^\dagger$, and $i : \mathcal{R}^\perp \rightarrow \mathcal{R}$ and $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger = \mathcal{R}/\mathcal{R}^\perp$ are as above. Then \mathcal{R}^\perp and \mathcal{R}^\dagger are of the same type (finite Abelian, essentially finite Abelian or Schurian) as \mathcal{R} . Moreover, the quotient functor $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ has a left adjoint $j_!$ and a right adjoint j_* :*

$$\begin{array}{ccc}
& & j_* \\
& \curvearrowright & \\
\mathcal{R} & \xrightarrow{j} & \mathcal{R}^\dagger \\
& \curvearrowleft & \\
& & j_!
\end{array}$$

The counit of one of the adjunctions and the unit of the other give isomorphisms:

$$j \circ j_* \xrightarrow{\sim} \text{Id}_{\mathcal{R}^\dagger} \xrightarrow{\sim} j \circ j_!$$

In particular, $j_!$ and j_* are fully faithful.

Proof. Fix a pointed algebra realization

$$A = \bigoplus_{a,b \in \mathbf{B}} e_a A e_b$$

of \mathcal{R} , so A is finite-dimensional, essentially finite-dimensional or locally finite-dimensional according to whether \mathcal{R} is finite Abelian, essentially finite Abelian or Schurian. Let

$$A^\downarrow = \bigoplus_{a,b \in \mathbf{B}^\downarrow} \bar{e}_a A^\downarrow \bar{e}_b := A / (e_c \mid c \in \mathbf{B}^\uparrow), \quad A^\uparrow := \bigoplus_{a,b \in \mathbf{B}^\uparrow} e_a A e_b,$$

where \bar{x} denotes the canonical image of $x \in A$ under the quotient map $A \twoheadrightarrow A^\downarrow$. Then it is clear that \mathcal{R}^\downarrow is equivalent to $A^\downarrow\text{-mod}_{\text{fd}}$ in the finite Abelian or essentially finite Abelian cases, and to $A^\downarrow\text{-mod}_{\text{lfid}}$ in the Schurian case. As A^\downarrow satisfies the same finiteness properties as A , we deduce that \mathcal{R}^\downarrow is of the same type as \mathcal{R} .

The quotient category \mathcal{R}^\uparrow is realized by the algebra A^\uparrow , and the quotient functor j becomes the functor that sends an A -module V to

$$jV := \bigoplus_{a \in \mathbf{B}^\uparrow} e_a V \quad (2.24)$$

with A^\uparrow acting by restricting the action of A . We deduce that \mathcal{R}^\uparrow is again of the same type as \mathcal{R} . Since j is isomorphic to $\bigoplus_{b \in \mathbf{B}^\uparrow} \text{Hom}_A(Ae_b, -)$, it has the left adjoint

$$j_! := \left(\bigoplus_{b \in \mathbf{B}^\uparrow} Ae_b \right) \otimes_{A^\uparrow} ? : A^\uparrow\text{-mod} \rightarrow A\text{-mod} \quad (2.25)$$

thanks to Lemma 2.7(1). From this, it is clear that the unit of adjunction $\text{Id}_{\mathcal{R}^\uparrow} \rightarrow j \circ j_!$ is an isomorphism. On the other hand, j is also isomorphic to the tensor functor $(\bigoplus_{b \in \mathbf{B}^\uparrow} e_b A) \otimes_{A^\uparrow} ?$, so Lemma 2.7(1) also gives that j has the right adjoint

$$j_* := \bigoplus_{a \in \mathbf{B}} \text{Hom}_{A^\uparrow} \left(\bigoplus_{b \in \mathbf{B}^\uparrow} e_b A e_a, ? \right) : A^\uparrow\text{-mod} \rightarrow A\text{-mod}. \quad (2.26)$$

Again using this we see that the counit $j \circ j_* \rightarrow \text{Id}_{\mathcal{R}^\uparrow}$ is an isomorphism. \square

The situation when \mathcal{R} is locally finite Abelian is more complicated. Continuing with the above notation, it follows immediately from Lemma 2.1 that the Serre subcategory \mathcal{R}^\downarrow and the quotient category \mathcal{R}^\uparrow are locally Schurian too. The following lemma explains how to obtain an explicit coalgebra realization of \mathcal{R}^\downarrow starting from one for \mathcal{R} .

Lemma 2.25. *Suppose that $\mathcal{R} = \text{comod}_{\text{fd}}\text{-}C$ for a coalgebra C . Let C^\downarrow be the largest right coideal of C belonging to \mathcal{R}^\downarrow . Then C^\downarrow is a subcoalgebra of C . Moreover, \mathcal{R}^\downarrow consists of all $V \in \text{comod}_{\text{fd}}\text{-}C$ such that the image of the structure map $\eta : V \rightarrow V \otimes C$ is contained in $V \otimes C^\downarrow$, i.e., we have that $\mathcal{R}^\downarrow = \text{comod}_{\text{fd}}\text{-}C^\downarrow$.*

Proof. For a right comodule V with structure map $\eta : V \rightarrow V \otimes C$, we can consider $V \otimes C$ as a right comodule with structure map $\text{id} \otimes \delta$. The coassociative and counit axioms imply that η is an injective homomorphism of right comodules. We deduce that all irreducible subquotients of V belong to \mathcal{R}^\downarrow if and only if $\eta(V) \subseteq V \otimes C^\downarrow$. Applying this with $V = C^\downarrow$ shows that C^\downarrow is a subcoalgebra. Applying it to $V \in \mathcal{R}$ shows that $V \in \mathcal{R}^\downarrow$ if and only if $\eta(V) \subseteq V \otimes C^\downarrow$. \square

For locally finite Abelian \mathcal{R} , the quotient category \mathcal{R}^\dagger can also be realized explicitly as a category of comodules: if $\mathcal{R} = \text{comod}_{\text{fd}}\text{-}C$ then $\mathcal{R}^\dagger = \text{comod}_{\text{fd}}\text{-}eCe$ for an idempotent $e \in C^*$ and the quotient functor j becomes the idempotent truncation functor defined by e . This is reviewed in detail in [Nav]. It follows that the extension $j : \text{Ind}(\mathcal{R}) \rightarrow \text{Ind}(\mathcal{R}^\dagger)$ of j to the ind-completions always has a right adjoint j_* with $j \circ j_* \cong \text{Id}_{\text{Ind}(\mathcal{R}^\dagger)}$. However, this adjoint does not necessarily take objects of \mathcal{R}^\dagger to objects of \mathcal{R} , so that the original functor $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ need not have a right adjoint itself. For left adjoints, the situation is even a bit worse since one should really pass to the pro-completions. For our purposes, though, it will always be sufficient to impose the stronger condition from (i) of the following lemma; this ensures that both adjoints exist without any need to pass to ind- or pro-completions.

Lemma 2.26. *Suppose that \mathcal{R} is locally finite Abelian, and let $\mathbf{B}^\dagger \subseteq \mathbf{B}$ and $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be as above. Then the following are equivalent:*

- (i) $L(b)$ has an injective hull $I(b)$ and a projective cover $P(b)$ in \mathcal{R} for all $b \in \mathbf{B}^\dagger$;
- (ii) \mathcal{R}^\dagger is essentially finite Abelian and the quotient functor $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ has a left adjoint $j_!$ and a right adjoint j_* :

$$\begin{array}{ccc} & j_* & \\ \curvearrowright & & \curvearrowleft \\ \mathcal{R} & \xrightarrow{j} & \mathcal{R}^\dagger \\ \curvearrowleft & & \curvearrowright \\ & j_! & \end{array}$$

When these properties hold, there are isomorphisms $j \circ j_* \cong \text{Id}_{\mathcal{R}^\dagger} \cong j \circ j_!$ just like in Lemma 2.24.

Proof. (i) \Rightarrow (ii): Let $j_* : \text{Ind}(\mathcal{R}^\dagger) \rightarrow \text{Ind}(\mathcal{R})$ be the right adjoint of $j : \text{Ind}(\mathcal{R}) \rightarrow \text{Ind}(\mathcal{R}^\dagger)$ as in [Nav]. For $b \in \mathbf{B}^\dagger$, let $I^\dagger(b)$ be an injective hull of $L^\dagger(b)$ in $\text{Ind}(\mathcal{R}^\dagger)$. By adjunction properties, $j_* I^\dagger(b)$ is an injective hull of $L(b)$ in $\text{Ind}(\mathcal{R})$, hence, $j_* I^\dagger(b) \cong I(b)$ which has finite length by assumption. From $j \circ j_* \cong \text{Id}_{\text{Ind}(\mathcal{R}^\dagger)}$, we deduce that $I^\dagger(b) \cong j I(b)$ is of finite length too, so $I^\dagger(b) \in \mathcal{R}^\dagger$ and \mathcal{R}^\dagger has enough injectives. We have shown that j_* takes $I^\dagger(b)$ to $I(b) \in \mathcal{R}$, hence using left exactness we deduce that it takes any object of finite length to an object of finite length. This means that the restriction $j_* : \mathcal{R}^\dagger \rightarrow \mathcal{R}$ is well-defined and gives a right adjoint to $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$. The dual argument shows that \mathcal{R}^\dagger has enough projectives and that $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ has a left adjoint $j_! : \mathcal{R}^\dagger \rightarrow \mathcal{R}$. Finally we deduce that \mathcal{R}^\dagger is essentially finite Abelian due to Corollary 2.20.

(ii) \Rightarrow (i): We can take $I(b) := j_* I^\dagger(b)$ and $P(b) := j_! P^\dagger(b)$ where $I^\dagger(b)$ is an injective hull and $P^\dagger(b)$ is a projective cover of $L^\dagger(b)$ in \mathcal{R}^\dagger . \square

In the locally finite Abelian or Schurian cases, we may use the same notations $i, i^*, i^!$ for the natural extensions of these functors to the ind-completions $\text{Ind}(\mathcal{R}), \text{Ind}(\mathcal{R}^\dagger)$ or $\text{Ind}(\mathcal{R}_c), \text{Ind}(\mathcal{R}_c^\dagger)$, respectively. Similarly, we will use the notations $j, j_*, j_!$ for the extensions of these to the appropriate ind-completions, assuming the equivalent conditions from Lemma 2.26 hold in the locally finite Abelian case.

Lemma 2.27. *Continuing with the above setup, assume either that \mathcal{R} is finite Abelian, essentially finite Abelian, or Schurian, or that \mathcal{R} is locally finite Abelian and the equivalent conditions from Lemma 2.26 hold. For $b \in \mathbf{B}^\dagger$, let $P(b)$ (resp., $I(b)$) and $P^\dagger(b)$ (resp., $I^\dagger(b)$) be a projective cover (resp., injective hull) of $L(b)$ in \mathcal{R} and a projective cover (resp., injective hull) of $L^\dagger(b)$ in \mathcal{R}^\dagger . Then we have that*

$$jP(b) \cong P^\dagger(b), \quad jI(b) \cong I^\dagger(b), \quad j_! P^\dagger(b) \cong P(b), \quad j_* I^\dagger(b) \cong I(b).$$

Moreover, the adjunction gives isomorphisms

$$\text{Hom}_{\mathcal{R}}(P(b), j_* V) \cong \text{Hom}_{\mathcal{R}^\dagger}(P^\dagger(b), V), \quad \text{Hom}_{\mathcal{R}}(j_! V, I(b)) \cong \text{Hom}_{\mathcal{R}^\dagger}(V, I^\dagger(b)) \quad (2.27)$$

for $V \in \mathcal{R}^\dagger$, hence, $[V : L^\dagger(b)] = [j_* V : L(b)] = [j_! V : L(b)]$ for all $b \in \mathbf{B}^\dagger$.

Proof. Take $b \in \mathbf{B}^\dagger$. By adjunction properties, $j_!P^\dagger(b)$ is a projective cover of $L(b)$ in \mathcal{R} , so it is isomorphic to $P(b)$. Hence, $j(j_!P^\dagger(b)) \cong P^\dagger(b) \cong jP(b)$; similarly for injectives. The remaining assertions follow. \square

3. GENERALIZATIONS OF HIGHEST WEIGHT CATEGORIES

In this section, we define the various generalizations of highest weight categories and derive some of their fundamental properties in the four settings of finite Abelian, essentially finite Abelian, Schurian, and locally finite Abelian categories. The important definitions in the section are Definitions 3.7, 3.34 and 3.50. The reader new to these ideas may find it helpful to assume initially that all of the strata are simple in the sense of Lemma 3.4, when the definitions specialize to the notions of finite, essentially finite, upper finite and lower finite highest weight categories, respectively.

3.1. Stratifications and the associated standard and costandard objects. Let (Λ, \leq) be a poset. It is *interval finite* (resp., *upper finite*, resp., *lower finite*) if the interval $[\lambda, \mu] := \{\nu \in \Lambda \mid \lambda \leq \nu \leq \mu\}$ (resp., $[\lambda, \infty) := \{\nu \in \Lambda \mid \lambda \leq \nu\}$, resp., $(-\infty, \mu] := \{\nu \in \Lambda \mid \nu \leq \mu\}$) is finite for all $\lambda, \mu \in \Lambda$. A *lower set* (resp., *upper set*) means a subset Λ^\downarrow (resp., Λ^\uparrow) such that $\mu \leq \lambda \in \Lambda^\downarrow \Rightarrow \mu \in \Lambda^\downarrow$ (resp., $\mu \geq \lambda \in \Lambda^\uparrow \Rightarrow \mu \in \Lambda^\uparrow$).

A *stratification function* $\rho : \mathbf{B} \rightarrow \Lambda$ is a function from a set \mathbf{B} to a poset (Λ, \leq) such that all of the fibers $\mathbf{B}_\lambda := \rho^{-1}(\lambda)$ are finite. We often use other obvious notations like $\mathbf{B}_{\leq \lambda} := \bigcup_{\mu \leq \lambda} \mathbf{B}_\mu$, $\mathbf{B}_{< \lambda} := \bigcup_{\mu < \lambda} \mathbf{B}_\mu$, etc..

A *stratification* of an Abelian category \mathcal{R} is a quintuple $(\mathbf{B}, L, \rho, \Lambda, \leq)$ consisting of a set \mathbf{B} , a function L labelling a full set $\{L(b) \mid b \in \mathbf{B}\}$ of pairwise inequivalent irreducible objects in \mathcal{R} , and a stratification function $\rho : \mathbf{B} \rightarrow \Lambda$ for the poset (Λ, \leq) . In the case that ρ is a bijection, one can use it to identify \mathbf{B} with Λ , writing $L(\lambda)$ instead of $L(b)$; similarly for all of the other families of objects indexed by the set \mathbf{B} to be introduced shortly.

Given a stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ of \mathcal{R} and $\lambda \in \Lambda$, let $\mathcal{R}_{\leq \lambda}$ and $\mathcal{R}_{< \lambda}$ be the Serre subcategories of \mathcal{R} associated to the subsets $\mathbf{B}_{\leq \lambda}$ and $\mathbf{B}_{< \lambda}$ of \mathbf{B} , respectively. We denote the inclusion functors by

$$i_{\leq \lambda} : \mathcal{R}_{\leq \lambda} \rightarrow \mathcal{R}, \quad i_{< \lambda} : \mathcal{R}_{< \lambda} \rightarrow \mathcal{R}, \quad (3.1)$$

The left and right adjoints of $i_{\leq \lambda}$ are $i_{\leq \lambda}^*$ and $i_{\leq \lambda}^\dagger$ as in Lemma 2.23. We say that the stratification is

- (F) a *finite stratification* if \mathcal{R} is a finite Abelian category (so that \mathbf{B} is a finite set);
- (EF) an *essentially finite stratification* if \mathcal{R} is an essentially finite Abelian category and the poset Λ is interval finite;
- (LF) a *lower finite stratification* if \mathcal{R} is a locally finite Abelian category and the poset Λ is lower finite;
- (UF) an *upper finite stratification* if \mathcal{R} is a Schurian category and the poset Λ is upper finite.

In these four cases, the induced stratifications of the subcategories $\mathcal{R}_{< \lambda}$ and $\mathcal{R}_{\leq \lambda}$ are automatically of the same type.

By an *admissible stratification*, we mean a stratification of one of the above four types such that the following axiom is satisfied when in type (LF) (it holds automatically for the other types):

- (A) The irreducible object $L(b)$ has both a projective cover and an injective hull in $\mathcal{R}_{\leq \rho(b)}$ for all $b \in \mathbf{B}$.

This is a significant restriction on the sorts of lower finite Abelian categories that can be considered; for example, the category $\mathcal{R}ep(\mathbb{G}_a)$ of rational representations of the additive group does not have this property. Using Lemma 2.21 together with Lemma 2.26 in the

lower finite case, we deduce for $\lambda \in \Lambda$ that the quotient category $\mathcal{R}_\lambda := \mathcal{R}_{\leq \lambda} / \mathcal{R}_{< \lambda}$ is finite Abelian in all cases. Let

$$j^\lambda : \mathcal{R}_{\leq \lambda} \rightarrow \mathcal{R}_\lambda \quad (3.2)$$

be the quotient functor. The objects

$$\{L_\lambda(b) := j^\lambda L(b) \mid b \in \mathbf{B}_\lambda\} \quad (3.3)$$

give a full set of pairwise inequivalent irreducible objects in \mathcal{R}_λ . Moreover, we are in a recollement situation as in Lemmas 2.23, 2.24 and 2.26:

$$\begin{array}{ccc} & \overset{i_{< \lambda}^!}{\curvearrowright} & \overset{j_\#^\lambda}{\curvearrowright} \\ \mathcal{R}_{< \lambda} & \xrightarrow{i_{< \lambda}} & \mathcal{R}_{\leq \lambda} & \xrightarrow{j^\lambda} & \mathcal{R}_\lambda \\ & \underset{i_{< \lambda}^*}{\curvearrowright} & \underset{j_!^\lambda}{\curvearrowright} \end{array} \quad (3.4)$$

Let $P_\lambda(b)$ be a projective cover and $I_\lambda(b)$ be an injective hull of $L_\lambda(b)$ in \mathcal{R}_λ . By Lemma 2.27, these are isomorphic to the images of the projective cover and injective hull of $L(b)$ in $\mathcal{R}_{\leq \lambda}$, respectively. Finally, define *standard*, *costandard*, *proper standard* and *proper costandard objects* $\Delta(b), \nabla(b), \bar{\Delta}(b)$ and $\bar{\nabla}(b)$ according to (1.1).

Lemma 3.1. *Suppose we are given an admissible stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ of \mathcal{R} . Take $b \in \mathbf{B}$ and set $\lambda := \rho(b)$.*

- (1) *The standard object $\Delta(b)$ is a projective cover of $L(b)$ in $\mathcal{R}_{\leq \lambda}$. The proper standard object $\bar{\Delta}(b)$ is the largest quotient of $\Delta(b)$ such that all composition factors of $\text{rad } \bar{\Delta}(b)$ are of the form $L(c)$ for $c \in \mathbf{B}_{< \lambda}$.*
- (2) *The costandard object $\nabla(b)$ is an injective hull of $L(b)$ in $\mathcal{R}_{\leq \lambda}$. The proper costandard object $\bar{\nabla}(b)$ is the largest subobject of $\nabla(b)$ such that all composition factors of $\bar{\nabla}(b) / \text{soc } \bar{\nabla}(b)$ are of the form $L(c)$ for $c \in \mathbf{B}_{< \lambda}$.*

Proof. We just check (1) since (2) is similar. We have that $\Delta(b)$ is a projective cover of $L(b)$ in $\mathcal{R}_{\leq \lambda}$ by Lemma 2.27. It remains to prove the statement about $\bar{\Delta}(b)$. Assume $[\bar{\Delta}(b) : L(c)] \neq 0$. Since $\bar{\Delta}(b) \in \mathcal{R}_{\leq \lambda}$, we have $\rho(c) \leq \rho(b)$. If $\rho(c) = \rho(b)$ then

$$[\bar{\Delta}(b) : L(c)] = [j^\lambda \bar{\Delta}(b) : j^\lambda L(c)] = [L_\lambda(b) : L_\lambda(c)] = \delta_{b,c}.$$

Thus, $\bar{\Delta}(b)$ is such a quotient of $\Delta(b)$. To show that it is the largest such quotient, it suffices to show that the kernel K of $\Delta(b) \twoheadrightarrow \bar{\Delta}(b)$ is finitely generated with head that only involves irreducibles $L(c)$ with $\rho(c) = \rho(b)$. To see this, apply the right exact functor $j_!^\lambda$ to a short exact sequence $0 \rightarrow \hat{K} \rightarrow P_\lambda(b) \rightarrow L_\lambda(b) \rightarrow 0$ to get an epimorphism $j_!^\lambda \hat{K} \twoheadrightarrow K$. Then observe that $j_!^\lambda \hat{K}$ is finitely generated as $j_!^\lambda$ is a left adjoint, and its head only involves irreducibles $L(c)$ with $\rho(c) = \rho(b)$. The latter assertion follows because $\text{Hom}_{\mathcal{R}}(j_!^\lambda \hat{K}, L(c)) \cong \text{Hom}_{\mathcal{R}_\lambda}(\hat{K}, j^\lambda L(c))$ for $c \in \mathbf{B}_{\leq \lambda}$. \square

Corollary 3.2. *We have that $\dim \text{Hom}_{\mathcal{R}}(\Delta(b), \bar{\nabla}(c)) = \dim \text{Hom}_{\mathcal{R}}(\bar{\Delta}(b), \nabla(c)) = \delta_{b,c}$ for all $b, c \in \mathbf{B}$.*

Lemma 3.3. *Suppose that we are given an admissible stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ of \mathcal{R} , and in addition that \mathcal{R} possesses a contravariant autoequivalence ${}^\vee$ which preserves isomorphism classes of irreducibles. Then we have that $P(b)^\vee \cong I(b)$, $I(b)^\vee \cong P(b)$, $\Delta(b)^\vee \cong \nabla(b)$, $\bar{\Delta}(b)^\vee \cong \bar{\nabla}(b)$, $\nabla(b)^\vee \cong \Delta(b)$ and $\bar{\nabla}(b)^\vee \cong \bar{\Delta}(b)$ for all $b \in \mathbf{B}$.*

Proof. Since $L(b)^\vee \cong L(b)$, we have that $I(b)^\vee \cong P(b)$ and $P(b)^\vee \cong I(b)$. Then the statements about $\Delta(b)^\vee, \bar{\Delta}(b)^\vee, \bar{\nabla}(b)^\vee$ and $\nabla(b)^\vee$ follow using Lemma 3.1. \square

For $\lambda \in \Lambda$, we say that the stratum \mathcal{R}_λ is *simple* if it is equivalent to the category Vec_{fd} of finite-dimensional vector spaces.

Lemma 3.4. *The following are equivalent:*

- (i) *all of the strata are simple;*

- (ii) ρ is a bijection and $\Delta(\lambda) = \bar{\Delta}(\lambda)$ for all $\lambda \in \Lambda$;
- (iii) ρ is a bijection and $\text{Hom}_{\mathcal{R}}(\Delta(\lambda), \nabla(\lambda))$ is one-dimensional;
- (iv) ρ is a bijection and $\nabla(\lambda) = \bar{\nabla}(\lambda)$ for all $\lambda \in \Lambda$.

Proof. (i) \Rightarrow (ii): Take $\lambda \in \Lambda$. As the stratum \mathcal{R}_λ is simple, $\mathbf{B}_\lambda = \{b_\lambda\}$ is a singleton and $P_\lambda(b_\lambda) = L_\lambda(b_\lambda)$. We deduce that ρ is a bijection and $\Delta(b_\lambda) = \bar{\Delta}(b_\lambda)$.

(ii) \Rightarrow (iii): This follows because $\nabla(\lambda)$ is the injective hull of $L(\lambda)$ in $\mathcal{R}_{\leq \lambda}$.

(iii) \Rightarrow (iv): This follows because $\Delta(\lambda)$ is the projective cover of $L(\lambda)$ in $\mathcal{R}_{\leq \lambda}$.

(iv) \Rightarrow (i): Take $\lambda \in \Lambda$. Then \mathcal{R}_λ has just one irreducible object (up to isomorphism), namely, $j^\lambda \bar{\nabla}(\lambda)$. Since this equals $j^\lambda \nabla(\lambda)$, it is also projective. Hence, \mathcal{R}_λ is simple. \square

Given a *sign function* $\varepsilon : \Lambda \rightarrow \{\pm\}$, we introduce the ε -*standard* and ε -*costandard* objects $\Delta_\varepsilon(b)$ and $\nabla_\varepsilon(b)$ as in (1.2). Corollary 3.2 implies that

$$\dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), \nabla_\varepsilon(c)) = \delta_{b,c} \quad (3.5)$$

for $b, c \in \mathbf{B}$. A Δ_ε -*flag* of $V \in \mathcal{R}$ means a finite filtration $0 = V_0 < V_1 < \dots < V_n = V$ with sections $V_m/V_{m-1} \cong \Delta_\varepsilon(b_m)$ for $b_m \in \mathbf{B}$. Similarly, we define ∇_ε -*flags*. We denote the exact subcategories of \mathcal{R} consisting of all objects with a Δ_ε -flag or a ∇_ε -flag by $\Delta_\varepsilon(\mathcal{R})$ and $\nabla_\varepsilon(\mathcal{R})$, respectively.

A Δ -*flag* (resp., $\bar{\nabla}$ -*flag*) is a Δ_ε -flag (resp., ∇_ε -flag) in the special case that $\varepsilon = +$. A $\bar{\Delta}$ -*flag* (resp., $\bar{\nabla}$ -*flag*) is a Δ_ε -flag (resp., ∇_ε -flag) in the special case that $\varepsilon = -$. We denote the exact subcategories of \mathcal{R} consisting of all objects with a Δ -flag, a $\bar{\Delta}$ -flag, a ∇ -flag or a $\bar{\nabla}$ -flag by $\Delta(\mathcal{R})$, $\bar{\Delta}(\mathcal{R})$, $\nabla(\mathcal{R})$ and $\bar{\nabla}(\mathcal{R})$, respectively.

3.2. Finite and essentially finite ε -stratified categories. Throughout this subsection, \mathcal{R} is a finite or essentially finite Abelian category equipped with a finite or essentially finite stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Also $\varepsilon : \Lambda \rightarrow \{\pm\}$ denotes a sign function. Let $P(b)$ and $I(b)$ be a projective cover and an injective hull of $L(b)$, respectively. We also need the objects from (1.1)–(1.2). Consider the following two properties:

- $(\widehat{P\Delta}_\varepsilon)$ For each $b \in \mathbf{B}$, there exists a projective object P_b admitting a Δ_ε -flag with $\Delta_\varepsilon(b)$ at the top and other sections $\Delta_\varepsilon(c)$ for $c \in \mathbf{B}$ with $\rho(c) \geq \rho(b)$.
- $(\widehat{I\nabla}_\varepsilon)$ For each $b \in \mathbf{B}$, there exists an injective object I_b admitting a ∇_ε -flag with $\nabla_\varepsilon(b)$ at the bottom and other sections $\nabla_\varepsilon(c)$ for $c \in \mathbf{B}$ with $\rho(c) \geq \rho(b)$.

It is trivial to see that the property $(P\Delta_\varepsilon)$ formulated in the introduction implies $(\widehat{P\Delta}_\varepsilon)$, and similarly $(I\nabla_\varepsilon)$ implies $(\widehat{I\nabla}_\varepsilon)$. The seemingly weaker properties $(\widehat{P\Delta}_\varepsilon)$ – $(\widehat{I\nabla}_\varepsilon)$ are often easier to check in concrete examples. The essence of the following fundamental theorem appeared originally in [ADL], extending earlier work of Dlab [Dla1].

Theorem 3.5. *The four properties $(\widehat{P\Delta}_\varepsilon)$, $(\widehat{I\nabla}_\varepsilon)$, $(P\Delta_\varepsilon)$ and $(I\nabla_\varepsilon)$ are equivalent. When these properties hold, the standardization functor $j_!^\lambda$ is exact whenever $\varepsilon(\lambda) = -$, and the costandardization functor j_*^λ is exact whenever $\varepsilon(\lambda) = +$.*

Remark 3.6. When all strata are simple, the properties $(\widehat{P\Delta}_\varepsilon)$ – $(\widehat{I\nabla}_\varepsilon)$ may be written more succinctly as the following:

- $(\widehat{P\Delta})$ For each $\lambda \in \Lambda$, there exists a projective object P_λ admitting a Δ -flag with $\Delta(\lambda)$ at the top and other sections of the form $\Delta(\mu)$ for $\mu \in \Lambda$ with $\mu \geq \lambda$.
- $(\widehat{I\nabla})$ For each $\lambda \in \Lambda$, there exists an injective object I_λ admitting a ∇ -flag with $\nabla(\lambda)$ at the bottom and other sections of the form $\nabla(\mu)$ for $\mu \in \Lambda$ with $\mu \geq \lambda$.

Theorem 3.5 shows that these are equivalent to the properties $(P\Delta)$ – $(I\nabla)$ from the introduction, as was explained originally by Cline, Parshall and Scott in [CPS1].

We postpone the proof of Theorem 3.5 until a little later in the subsection. It is important because it justifies the next key definition (ε S) and its variations (FS), (ε HW), (FHW) and (HW).

Definition 3.7. Let \mathcal{R} be an Abelian category equipped with a finite (resp., essentially finite) stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$.

- (ε S) We say that \mathcal{R} is a *finite* (resp., *essentially finite*) ε -*stratified category* if one of the equivalent properties $(\widehat{P\Delta}_\varepsilon)$ – $(\widehat{I\nabla}_\varepsilon)$ holds for a given choice of sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$.
- (FS) We say \mathcal{R} is a *finite* (resp., *essentially finite*) *fully stratified category* if one of these properties holds for all choices of sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$.
- (ε HW) We say \mathcal{R} is a *finite* (resp., *essentially finite*) ε -*highest weight category* if the stratification function ρ is a bijection, i.e., each stratum has a unique irreducible object (up to isomorphism), and one of the equivalent properties $(\widehat{P\Delta}_\varepsilon)$ – $(\widehat{I\nabla}_\varepsilon)$ holds for a given choice of sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$.
- (FHW) We say \mathcal{R} is a *finite* (resp., *essentially finite*) *fibered highest weight category* if the stratification function ρ is a bijection and one of these properties holds for all choices of sign function.
- (HW) We say \mathcal{R} is a *finite* (resp., *essentially finite*) *highest weight category* if all of the strata are simple (cf. Lemma 3.4) and one of the equivalent properties $(\widehat{P\Delta})$ – $(\widehat{I\nabla})$ holds.

Remark 3.8. The language “fibered highest weight” in Definition 3.7 is a departure from the existing literature, where such categories are usually referred to as *properly stratified categories*; this terminology goes back to the work of Dlab [Dla2]. A recent exposition which takes a more traditional viewpoint than here can be found in [CouZ]. In particular, in [CouZ, Def. 2.7.4], one finds five types of finite-dimensional algebra A defined in terms of properties of the category $A\text{-mod}_{\text{fd}}$, namely, standardly stratified algebras, exactly standardly stratified algebras, strongly stratified algebras, properly stratified algebras, and quasi-hereditary algebras. In our preferred language, these are $+$ -stratified algebras, stratified algebras, $+$ -quasi-hereditary algebras, properly stratified algebras, and quasi-hereditary algebras, respectively, as in Table 1 from the introduction. For further reference to the original literature, [CouZ, §A.2] is helpful.

We can view $\{L(b) \mid b \in \mathbf{B}\}$ equivalently as a full set of pairwise inequivalent irreducible objects in \mathcal{R}^{op} . The stratification of \mathcal{R} is also one of \mathcal{R}^{op} . The indecomposable projectives and injectives in \mathcal{R}^{op} are $I(b)$ and $P(b)$, while the $(-\varepsilon)$ -standard and $(-\varepsilon)$ -costandard objects in \mathcal{R}^{op} are $\nabla_\varepsilon(b)$ and $\Delta_\varepsilon(b)$, respectively. So we can reinterpret Theorem 3.5 as the following.

Theorem 3.9. \mathcal{R} is ε -stratified, fully stratified, ε -highest weight, fibered highest weight or highest weight if and only if \mathcal{R}^{op} is $(-\varepsilon)$ -stratified, fully stratified, $(-\varepsilon)$ -highest weight, fibered highest weight or highest weight, respectively.

Now we must prepare for the proof Theorem 3.5. The main step in the argument will be provided by the *homological criterion* for ∇_ε -flags from the next Theorem 3.11. In turn, the proof of this criterion reduces to the following lemma which treats a key special case. The reader wanting to work fully through the proofs should look also at this point at the lemmas in §3.4 below.

Lemma 3.10. Assume that \mathcal{R} is an Abelian category equipped with a finite or essentially finite stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ and sign function ε , such that property $(\widehat{P\Delta}_\varepsilon)$ holds. Let λ be a maximal element of Λ with respect to the ordering \leq , and $V \in \mathcal{R}$ be an object satisfying the following properties:

- (i) $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), V) = 0$ for all $b \in \mathbf{B}$;
- (ii) $\text{soc } V \cong L(b_1) \oplus \cdots \oplus L(b_n)$ for $b_1, \dots, b_n \in \mathbf{B}_\lambda$.

Then V belongs to $\mathcal{R}_{\leq \lambda}$ (so that it makes sense to apply the functor j^λ to it), and

$$V \cong \begin{cases} j_*^\lambda(j^\lambda V) & \text{if } \varepsilon(\lambda) = +, \\ \nabla(b_1) \oplus \cdots \oplus \nabla(b_n) & \text{if } \varepsilon(\lambda) = -. \end{cases} \quad (3.6)$$

Moreover, in the case $\varepsilon(\lambda) = +$, the functor j_*^λ is exact. Hence, in both cases, we have that $V \in \nabla_\varepsilon(\mathcal{R})$.

Proof (assuming lemmas from §3.4 below). We first prove (3.6) in case $\varepsilon(\lambda) = -$. Let $W := \nabla(b_1) \oplus \cdots \oplus \nabla(b_n)$. By the maximality of λ and Lemma 3.46, this is an injective hull of $\text{soc } V$. So there is a short exact sequence $0 \rightarrow V \rightarrow W \rightarrow W/V \rightarrow 0$. For any $a \in \mathbf{B}$, we apply $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), ?)$ and use property (i) to get a short exact sequence

$$0 \longrightarrow \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), V) \xrightarrow{f} \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), W) \longrightarrow \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), W/V) \longrightarrow 0. \quad (3.7)$$

If $\rho(a) \neq \lambda$ then $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), W) = 0$ as none of the composition factors of $\Delta_\varepsilon(a)$ are constituents of $\text{soc } W$. If $\rho(a) = \lambda$ then $\Delta_\varepsilon(a) = \bar{\Delta}(a)$ and any homomorphism $\bar{\Delta}(a) \rightarrow W$ must factor through the unique irreducible quotient $L(a)$ of $\bar{\Delta}(a)$. So its image is contained in $\text{soc } W \subseteq V$, showing that f is an isomorphism. These arguments show that $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), W/V) = 0$ for all $a \in \mathbf{B}$. We deduce that $\text{soc}(W/V) = 0$, hence, $W/V = 0$, which is what we needed.

Now consider (3.6) when $\varepsilon(\lambda) = +$. By Lemma 3.46 again, the injective hull of V is $\nabla(b_1) \oplus \cdots \oplus \nabla(b_n)$, which is an object of $\mathcal{R}_{\leq \lambda}$. Hence, $V \in \mathcal{R}_{\leq \lambda}$. The unit of adjunction gives us a morphism $g : V \rightarrow W := j_*^\lambda(j^\lambda V)$. Since g becomes an isomorphism when we apply j^λ , its kernel belongs to $\mathcal{R}_{< \lambda}$. In view of property (2), we deduce that $\ker g = 0$, so g is a monomorphism. Hence, we can identify V with a subobject of W . To show that g is an epimorphism as well, we apply $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), ?)$ to $0 \rightarrow V \rightarrow W \rightarrow W/V \rightarrow 0$ to get the short exact sequence (3.7). By adjunction, the middle morphism space is isomorphic to $\text{Hom}_{\mathcal{R}_\lambda}(j^\lambda \Delta_\varepsilon(a), j^\lambda V)$, which is zero if $\rho(a) < \lambda$. If $\rho(a) = \lambda$ then $\Delta_\varepsilon(a) = \Delta(a)$ is the projective cover of $L(a)$ in \mathcal{R} by Lemma 3.46, and $j^\lambda \Delta_\varepsilon(a)$ is the projective cover of $L_\lambda(a)$ in \mathcal{R}_λ . We deduce that both the first and second morphism spaces in (3.7) are of the same dimension $[V : L(a)] = [j^\lambda V : L_\lambda(a)]$, so f must be an isomorphism. Therefore $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), W/V) = 0$ for all $a \in \mathbf{B}$, hence, $V = W$ and (3.6) is proved.

To complete the proof, we must show that j_*^λ is exact when $\varepsilon(\lambda) = +$. For this, we use induction on composition length to show that j_*^λ is exact on any short exact sequence $0 \rightarrow K \rightarrow X \rightarrow Q \rightarrow 0$ in \mathcal{R}_λ . For the induction step, suppose we are given such an exact sequence with $K, Q \neq 0$. By induction, $j_*^\lambda K$ and $j_*^\lambda Q$ both have filtrations with sections $\bar{\nabla}(b)$ for $b \in \mathbf{B}_\lambda$. Hence, by Lemma 3.48, we have that $\text{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(b), j_*^\lambda K) = \text{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(b), j_*^\lambda Q) = 0$ for all $n \geq 1$ and $b \in \mathbf{B}$. As it is a right adjoint, j_*^λ is left exact, so there is an exact sequence

$$0 \longrightarrow j_*^\lambda K \longrightarrow j_*^\lambda X \longrightarrow j_*^\lambda Q. \quad (3.8)$$

Let $Y := j_*^\lambda X / j_*^\lambda K$, so that there is a short exact sequence

$$0 \longrightarrow j_*^\lambda K \longrightarrow j_*^\lambda X \longrightarrow Y \longrightarrow 0. \quad (3.9)$$

To complete the argument, it suffices to show that $Y \cong j_*^\lambda Q$. To establish this, we show that Y satisfies both of the properties (i) and (ii); then, by the previous paragraph and exactness of j^λ , we get that $Y \cong j_*^\lambda(j^\lambda Y) \cong j_*^\lambda(X/K) \cong j_*^\lambda Q$, and we are done. To see that Y satisfies (i), we apply $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), ?)$ to (3.9) to get an exact sequence

$$\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), j_*^\lambda X) \longrightarrow \text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), Y) \longrightarrow \text{Ext}_{\mathcal{R}}^2(\Delta_\varepsilon(b), j_*^\lambda K).$$

The first Ext^1 is zero by Lemma 3.47. Since we already know that the Ext^2 term is zero, $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), Y) = 0$. To see that Y satisfies (ii), note comparing (3.8)–(3.9) that $Y \hookrightarrow j_*^\lambda Q$, and $\text{soc } j_*^\lambda Q$ is of the desired form by what we know about its $\bar{\nabla}_\varepsilon$ -flag. \square

Theorem 3.11 (Homological criterion for Δ_ε -flags). *Assume that \mathcal{R} is an Abelian category equipped with a finite or essentially finite stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ and sign function ε , such that property $(\widehat{P}\Delta_\varepsilon)$ holds. For $V \in \mathcal{R}$, the following properties are equivalent:*

- (i) $V \in \nabla_\varepsilon(\mathcal{R})$;
- (ii) $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), V) = 0$ for all $b \in \mathbf{B}$;

(iii) $\text{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(b), V) = 0$ for all $b \in \mathbf{B}$ and $n \geq 1$.

If these properties hold, the multiplicity $(V : \nabla_\varepsilon(b))$ of $\nabla_\varepsilon(b)$ as a section of a ∇_ε -flag of V is well-defined independent of the choice of flag, as it equals $\dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V)$.

Proof (assuming lemmas from §3.4 below). (iii) \Rightarrow (ii): Trivial.

(i) \Rightarrow (iii) and the final assertion of the lemma: These follow directly from Lemma 3.48.

(ii) \Rightarrow (i): Assume that V satisfies (ii). We prove that it has a ∇_ε -flag by induction on

$$d(V) := \sum_{b \in \mathbf{B}} \dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) \in \mathbb{N}.$$

The base case when $d(V) = 0$ is trivial as we have then that $V = 0$. For the induction step, let $\lambda \in \Lambda$ be minimal such that $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) \neq 0$ for some $b \in \mathbf{B}$. The Serre subcategory $\mathcal{R}_{\leq \lambda}$ with the induced (finite or essentially finite) stratification also satisfies $(\widehat{P\Delta}_\varepsilon)$ thanks to Lemma 3.45(2). Let $W := i_{\leq \lambda}^! V$. Because W is a subobject of V , we have by the minimality of λ that $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), W) \neq 0$ only if $b \in \mathbf{B}_\lambda$. Hence, $\text{soc } W \cong L(b_1) \oplus \cdots \oplus L(b_n)$ for $b_1, \dots, b_n \in \mathbf{B}_\lambda$. Thus, W satisfies the hypothesis (ii) from Lemma 3.10 (with V and \mathcal{R} there replaced by W and $\mathcal{R}_{\leq \lambda}$).

Now let $Q := V/W$. Take any $b \in \mathbf{B}$ and apply $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), ?)$ to the short exact sequence $0 \rightarrow W \rightarrow V \rightarrow Q \rightarrow 0$ to get the exact sequence

$$\begin{aligned} 0 &\longrightarrow \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), W) \longrightarrow \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) \longrightarrow \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), Q) \\ &\longrightarrow \text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), W) \longrightarrow 0 \longrightarrow \text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), Q) \longrightarrow \text{Ext}_{\mathcal{R}}^2(\Delta_\varepsilon(b), W). \end{aligned}$$

By the definition of W , the socle of Q has no constituent $L(b)$ for $b \in \mathbf{B}_{\leq \lambda}$. So, for $b \in \mathbf{B}_{\leq \lambda}$ the space $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), Q)$ is zero, and we get that $\text{Ext}_{\mathcal{R}_{\leq \lambda}}^1(\Delta_\varepsilon(b), W) \cong \text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), W) = 0$ for all such b . This verifies hypothesis (i) from Lemma 3.10. So now we can appeal to the lemma to deduce that $W \in \nabla_\varepsilon(\mathcal{R}_{\leq \lambda})$. Hence, $W \in \nabla_\varepsilon(\mathcal{R})$.

In view of Lemma 3.48, we get that $\text{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(b), W) = 0$ for all $n \geq 1$ and $b \in \mathbf{B}$. So, by the above exact sequence again, we get that $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), Q) = 0$ for all $b \in \mathbf{B}$, and moreover $d(Q) = d(V) - d(W) < d(V)$. Finally we appeal to the induction hypothesis to deduce that $Q \in \Delta_\varepsilon(\mathcal{R})$. Since we already know that $W \in \Delta_\varepsilon(\mathcal{R})$, this shows that $V \in \Delta_\varepsilon(\mathcal{R})$. \square

Corollary 3.12. *In the setup of Theorem 3.11, multiplicities in a ∇_ε -flag of $I(b)$ satisfy $(I(b) : \nabla_\varepsilon(c)) = [\Delta_\varepsilon(c) : L(b)]$.*

Corollary 3.13. *For \mathcal{R} as in Theorem 3.11, let $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ be a short exact sequence. If U and V have ∇_ε -flags then so does W .*

Proof of Theorem 3.5. Suppose that \mathcal{R} satisfies $(\widehat{P\Delta}_\varepsilon)$. Since $V = I(b)$ is injective, it satisfies the hypothesis of Theorem 3.11(ii). Hence, by that theorem, $I(b)$ has a ∇_ε -flag and the multiplicity $(I(b) : \nabla_\varepsilon(c))$ of $\nabla_\varepsilon(c)$ as a section of any such flag is given by

$$(I(b) : \nabla_\varepsilon(c)) = \dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(c), I(b)) = [\Delta_\varepsilon(c) : L(b)].$$

This is zero unless $\rho(b) \leq \rho(c)$. Also the bottom section must be $\nabla_\varepsilon(b)$ since $I(b)$ has socle $L(b)$. Thus, we have verified that \mathcal{R} satisfies $(I\nabla_\varepsilon)$. Moreover, Lemma 3.10 shows that j_*^λ is exact whenever $\varepsilon(\lambda) = +$, giving half of final assertion made in the statement of the theorem we are trying to prove.

Repeating the arguments in the previous paragraph but with \mathcal{R} replaced by \mathcal{R}^{op} and ε replaced with $-\varepsilon$ show that $(\widehat{I\nabla}_\varepsilon)$ implies $(P\Delta_\varepsilon)$ and that $j_!^\lambda$ is exact whenever $\varepsilon(\lambda) = -$. Since $(P\Delta_\varepsilon) \Rightarrow (\widehat{P\Delta}_\varepsilon)$ and $(I\nabla_\varepsilon) \Rightarrow (\widehat{I\nabla}_\varepsilon)$, this completes the proof. \square

So now Theorem 3.5 is proved and Definition 3.7 is in place. In the remainder of the subsection, we are going to develop some further fundamental properties of these sorts of category. We start off in the most general setup with \mathcal{R} being a finite or essentially

finite ε -stratified category. Again some of the proofs that follow invoke parts of the lemmas from §3.4. From Lemma 3.44 and the dual statement, deduce that

$$\mathrm{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), \Delta_\varepsilon(c)) = \mathrm{Ext}_{\mathcal{R}}^1(\nabla_\varepsilon(c), \nabla_\varepsilon(b)) = 0 \quad (3.10)$$

for $b, c \in \mathbf{B}$ with $\rho(b) \not\leq \rho(c)$. By “dual statement” here, we mean that one takes Lemma 3.44 with \mathcal{R} replaced by $\mathcal{R}^{\mathrm{op}}$ and ε by $-\varepsilon$, which we may do due to Theorem 3.9 and Lemma 2.12, then applies the contravariant isomorphism between \mathcal{R} and $\mathcal{R}^{\mathrm{op}}$. In a similar way, the following theorem follows immediately as it is the dual statement to Theorem 3.11.

Theorem 3.14 (Homological criterion for ∇_ε -flags). *Assume that \mathcal{R} is a finite or essentially finite ε -stratified category. For $V \in \mathcal{R}$, the following properties are equivalent:*

- (i) $V \in \Delta_\varepsilon(\mathcal{R})$;
- (ii) $\mathrm{Ext}_{\mathcal{R}}^1(V, \nabla_\varepsilon(b)) = 0$ for all $b \in \mathbf{B}$;
- (iii) $\mathrm{Ext}_{\mathcal{R}}^n(V, \nabla_\varepsilon(b)) = 0$ for all $b \in \mathbf{B}$ and $n \geq 1$.

Assuming that these properties hold, the multiplicity $(V : \Delta_\varepsilon(b))$ of $\Delta_\varepsilon(b)$ as a section of a Δ_ε -flag of V is well-defined independent of the choice of flag, as it equals $\dim \mathrm{Hom}_{\mathcal{R}}(V, \nabla_\varepsilon(b))$.

Corollary 3.15. $(P(b) : \Delta_\varepsilon(c)) = [\nabla_\varepsilon(c) : L(b)]$.

Corollary 3.16. *Let $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ be a short exact sequence in a finite or essentially finite ε -stratified category. If V and W have Δ_ε -flags then so does U .*

The following results about truncation to lower and upper sets are extremely useful; some aspects of them were already used in the proof of Theorem 3.11.

Theorem 3.17 (Truncation to lower sets). *Assume that \mathcal{R} is a finite or essentially finite ε -stratified category. Suppose that Λ^\downarrow is a lower set in Λ . Let $\mathbf{B}^\downarrow := \rho^{-1}(\Lambda^\downarrow)$ and $i : \mathcal{R}^\downarrow \rightarrow \mathcal{R}$ be the corresponding Serre subcategory of \mathcal{R} with the induced stratification. Then \mathcal{R}^\downarrow is itself a finite or essentially finite ε -stratified category according to whether Λ^\downarrow is finite or infinite. Moreover:*

- (1) *The distinguished objects in \mathcal{R}^\downarrow satisfy $L^\downarrow(b) \cong L(b)$, $P^\downarrow(b) \cong i^*P(b)$, $I^\downarrow(b) \cong i^!I(b)$, $\Delta^\downarrow(b) \cong \Delta(b)$, $\bar{\Delta}^\downarrow(b) \cong \bar{\Delta}(b)$, $\nabla^\downarrow(b) \cong \nabla(b)$ and $\bar{\nabla}^\downarrow(b) \cong \bar{\nabla}(b)$ for $b \in \mathbf{B}^\downarrow$.*
- (2) *i^* sends short exact sequences of objects in $\Delta_\varepsilon(\mathcal{R})$ to short exact sequences of objects in $\Delta_\varepsilon(\mathcal{R}^\downarrow)$, with $i^*\Delta(b) \cong \Delta^\downarrow(b)$ and $i^*\bar{\Delta}(b) \cong \bar{\Delta}^\downarrow(b)$ for $b \in \mathbf{B}^\downarrow$ and $i^*\Delta(b) = i^*\bar{\Delta}(b) = 0$ for $b \notin \mathbf{B}^\downarrow$.*
- (3) $\mathrm{Ext}_{\mathcal{R}}^n(V, iW) \cong \mathrm{Ext}_{\mathcal{R}^\downarrow}^n(i^*V, W)$ for $V \in \Delta_\varepsilon(\mathcal{R})$, $W \in \mathcal{R}^\downarrow$ and all $n \geq 0$.
- (4) *$i^!$ sends short exact sequences of objects in $\nabla_\varepsilon(\mathcal{R})$ to short exact sequences of objects in $\nabla_\varepsilon(\mathcal{R}^\downarrow)$, with $i^!\nabla(b) \cong \nabla^\downarrow(b)$ and $i^!\bar{\nabla}(b) \cong \bar{\nabla}^\downarrow(b)$ for $b \in \mathbf{B}^\downarrow$ and $i^!\nabla(b) = i^!\bar{\nabla}(b) = 0$ for $b \notin \mathbf{B}^\downarrow$.*
- (5) $\mathrm{Ext}_{\mathcal{R}}^n(iV, W) \cong \mathrm{Ext}_{\mathcal{R}^\downarrow}^n(V, i^!W)$ for $V \in \mathcal{R}^\downarrow$, $W \in \nabla_\varepsilon(\mathcal{R})$ and all $n \geq 0$.
- (6) $\mathrm{Ext}_{\mathcal{R}}^n(iV, iW) \cong \mathrm{Ext}_{\mathcal{R}^\downarrow}^n(V, W)$ for $V, W \in \mathcal{R}^\downarrow$ and $n \geq 0$.

Proof. Apart from (6), this follows by Lemma 3.45 and its dual. To prove (6), by the same argument as used to prove Lemma 3.45(4), it suffices to show that $(\mathbb{L}_n i^*)V = 0$ for $V \in \mathcal{R}^\downarrow$ and $n \geq 1$. Since any such V has finite length it suffices to consider an irreducible object in \mathcal{R}^\downarrow , i.e., we must show that $(\mathbb{L}_n i^*)L(b) = 0$ for $b \in \mathbf{B}^\downarrow$ and $n \geq 1$. Take a short exact sequence $0 \rightarrow K \rightarrow \Delta_\varepsilon(b) \rightarrow L(b) \rightarrow 0$ and apply i^* and Lemma 3.45(3) to get

$$0 \longrightarrow (\mathbb{L}_1 i^*)L(b) \longrightarrow i^*K \longrightarrow i^*\Delta_\varepsilon(b) \longrightarrow i^*L(b) \longrightarrow 0.$$

But $K, \Delta_\varepsilon(b)$ and $L(b)$ all lie in \mathcal{R}^\downarrow so i^* is the identity on them. We deduce that $(\mathbb{L}_1 i^*)L(b) = 0$. Degree shifting easily gives the result for $n > 1$. \square

Theorem 3.18 (Truncation to upper sets). *Assume that \mathcal{R} is a finite or essentially finite ε -stratified category. Suppose that Λ^\uparrow is an upper set in Λ . Let $\mathbf{B}^\uparrow := \rho^{-1}(\Lambda^\uparrow)$ and $j : \mathcal{R} \rightarrow \mathcal{R}^\uparrow$ be the corresponding Serre quotient category of \mathcal{R} with the induced*

stratification. Then \mathcal{R}^\dagger is itself a finite or essentially finite ε -stratified category according to whether \mathbf{B}^\dagger is finite or infinite. Moreover:

- (1) For $b \in \mathbf{B}^\dagger$, the distinguished objects $L^\dagger(b)$, $P^\dagger(b)$, $I^\dagger(b)$, $\Delta^\dagger(b)$, $\bar{\Delta}^\dagger(b)$, $\nabla^\dagger(b)$ and $\bar{\nabla}^\dagger(b)$ in \mathcal{R}^\dagger are isomorphic to the images under j of the corresponding objects of \mathcal{R} .
- (2) We have that $jL(b) = j\Delta(b) = j\bar{\Delta}(b) = j\nabla(b) = j\bar{\nabla}(b) = 0$ if $b \notin \mathbf{B}^\dagger$.
- (3) $\text{Ext}_{\mathcal{R}}^n(V, j_*W) \cong \text{Ext}_{\mathcal{R}^\dagger}^n(jV, W)$ for $V \in \mathcal{R}, W \in \nabla_\varepsilon(\mathcal{R}^\dagger)$ and all $n \geq 0$.
- (4) j_* sends short exact sequences of objects in $\nabla_\varepsilon(\mathcal{R}^\dagger)$ to short exact sequences of objects in $\nabla_\varepsilon(\mathcal{R})$, with $j_*\nabla^\dagger(b) \cong \nabla(b)$, $j_*\bar{\nabla}^\dagger(b) \cong \bar{\nabla}(b)$ and $j_*I^\dagger(b) \cong I(b)$ for $b \in \mathbf{B}^\dagger$.
- (5) $\text{Ext}_{\mathcal{R}}^n(j_!V, W) \cong \text{Ext}_{\mathcal{R}^\dagger}^n(V, jW)$ for $V \in \Delta_\varepsilon(\mathcal{R}^\dagger), W \in \mathcal{R}$ and all $n \geq 0$.
- (6) $j_!$ sends short exact sequences of objects in $\Delta_\varepsilon(\mathcal{R}^\dagger)$ to short exact sequences of objects in $\Delta_\varepsilon(\mathcal{R})$, with $j_!\Delta^\dagger(b) \cong \Delta(b)$, $j_!\bar{\Delta}^\dagger(b) \cong \bar{\Delta}(b)$ and $j_!P^\dagger(b) = P(b)$ for $b \in \mathbf{B}^\dagger$.

Proof. Apart from (4) and (6), this follows from Lemma 3.49 and its dual. For (4) and (6), it suffices to prove (4), since (6) is the equivalent dual statement. The descriptions of $j_*\nabla^\dagger(b)$, $j_*\bar{\nabla}^\dagger(b)$ and $j_*I^\dagger(b)$, follow from Lemma 3.49(1). It remains to prove the exactness. We can actually show slightly more, namely, that $(\mathbb{R}^1 j_*)V = 0$ for $V \in \nabla_\varepsilon(\mathcal{R}^\dagger)$ and $n \geq 1$. Take $V \in \nabla_\varepsilon(\mathcal{R}^\dagger)$. Consider a short exact sequence $0 \rightarrow V \rightarrow I \rightarrow Q \rightarrow 0$ in \mathcal{R}^\dagger with I injective. Apply the left exact functor j_* and consider the resulting long exact sequence:

$$0 \longrightarrow j_*V \longrightarrow j_*I \longrightarrow j_*Q \longrightarrow (\mathbb{R}^1 j_*)V \longrightarrow 0.$$

As V has a ∇_ε -flag, we can use (3) to see that $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), j_*V) \cong \text{Hom}_{\mathcal{R}^\dagger}(j\Delta_\varepsilon(b), V)$ and $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), j_*V) \cong \text{Ext}_{\mathcal{R}^\dagger}^1(j\Delta_\varepsilon(b), V)$ for every $b \in \mathbf{B}$. Hence, Theorem 3.11, j_*V has a ∇_ε -flag with

$$(j_*V : \nabla_\varepsilon(b)) = \dim \text{Hom}_{\mathcal{R}}(j\Delta_\varepsilon(b), V) = \begin{cases} (V : \nabla_\varepsilon^\dagger(b)) & \text{if } b \in \mathbf{B}^\dagger, \\ 0 & \text{otherwise.} \end{cases}$$

Both I and Q have ∇_ε -flags too, so we get similar statements for j_*I and j_*Q . Since $(I : \nabla_\varepsilon^\dagger(b)) = (V : \nabla_\varepsilon^\dagger(b)) + (Q : \nabla_\varepsilon^\dagger(b))$ by the exactness of the original sequence, we deduce that $0 \rightarrow j_*V \rightarrow j_*I \rightarrow j_*Q \rightarrow 0$ is exact. Hence, $(\mathbb{R}^1 j_*)V = 0$. This proves the result for $n = 1$. The result for $n > 1$ follows by a degree shifting argument. \square

Corollary 3.19. *Let notation be as in Theorem 3.18 and set $\mathbf{B}^\downarrow := \mathbf{B} \setminus \mathbf{B}^\dagger$.*

- (1) For $V \in \nabla_\varepsilon(\mathcal{R})$, there is a short exact sequence $0 \rightarrow K \rightarrow V \xrightarrow{\gamma} j_*(jV) \rightarrow 0$ where γ comes from the unit of adjunction, $j_*(jV)$ has a ∇_ε -flag with sections $\nabla_\varepsilon(b)$ for $b \in \mathbf{B}^\dagger$, and K has a ∇_ε -flag with sections $\nabla_\varepsilon(c)$ for $c \in \mathbf{B}^\downarrow$.
- (2) For $V \in \Delta_\varepsilon(\mathcal{R})$, there is a short exact sequence $0 \rightarrow j_!(jV) \xrightarrow{\delta} V \rightarrow Q \rightarrow 0$ where δ comes from the counit of adjunction, $j_!(jV)$ has a Δ_ε -flag with sections $\Delta_\varepsilon(b)$ for $b \in \mathbf{B}^\dagger$ and Q has a Δ_ε -flag with sections $\Delta_\varepsilon(c)$ for $c \in \mathbf{B}^\downarrow$.

Proof. We prove only (1), since (2) is just the dual statement. Using (3.10), we can order the ∇_ε -flag of V to get a short exact sequence $0 \rightarrow K \rightarrow V \rightarrow Q \rightarrow 0$ such that K has a ∇_ε -flag with sections $\nabla_\varepsilon(b)$ for $b \in \mathbf{B}^\downarrow$ and Q has a ∇_ε -flag with sections $\nabla_\varepsilon(c)$ for $c \in \mathbf{B}^\dagger$. For each $b \in \mathbf{B}^\dagger$, the unit of adjunction $\nabla_\varepsilon(b) \rightarrow j_*(j\nabla_\varepsilon(b))$ is an isomorphism; this follows from Theorem 3.18(4) using the observation that it becomes an isomorphism on applying j . Since j_* sends short exact sequences of objects in $\nabla_\varepsilon(\mathcal{R}^\dagger)$ to short exact sequences, we deduce that the unit of adjunction $Q \rightarrow j_*(jQ)$ is an isomorphism too. It remains to note that $jV \cong jQ$, hence, $j_*(jV) \cong j_*(jQ)$. \square

We proceed to discuss some of the additional features which show up when in one of the more refined settings (FS), (ε HW), (FHW) and (HW). By Theorem 3.9, \mathcal{R} is a fully stratified category (resp., fibered highest weight category) if and only if so is \mathcal{R}^{op} . The

following lemma shows that fully stratified categories in our terminology are the same as the “standardly stratified categories” defined by Losev and Webster in [LW, §2].

Lemma 3.20. *Given a stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ of \mathcal{R} , the following are equivalent:*

- (i) \mathcal{R} is a fully stratified category;
- (ii) \mathcal{R} is ε -stratified for every choice of sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$;
- (iii) \mathcal{R} is ε -stratified and $(-\varepsilon)$ -stratified for some choice of sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$;
- (iv) \mathcal{R} is ε -stratified for some $\varepsilon : \Lambda \rightarrow \{\pm\}$ and all of its standardization and costandardization functors are exact;
- (v) \mathcal{R} is a $+$ -stratified category and each $\Delta(b)$ has a $\bar{\Delta}$ -flag with sections $\bar{\Delta}(c)$ for c with $\rho(c) = \rho(b)$;
- (vi) \mathcal{R} is a $-$ -stratified category and each $\nabla(b)$ has a $\bar{\nabla}$ -flag with sections $\bar{\nabla}(c)$ for c with $\rho(c) = \rho(b)$.

Proof. (i) \Rightarrow (ii) \Rightarrow (iii): Obvious.

(iii) \Rightarrow (iv): Take ε as in (iii) so that \mathcal{R} is ε -stratified. The standardization functor $j_!^\lambda$ is exact when $\varepsilon(\lambda) = -$ by the last part of Theorem 3.5. Also \mathcal{R} is $(-\varepsilon)$ -stratified, so the same result gives that $j_!^\lambda$ is exact when $\varepsilon(\lambda) = +$. Similarly, all of the costandardization functors are exact too.

(iv) \Rightarrow (v): Applying the exact standardization functor $j_!^\lambda$ to a composition series of $P_\lambda(b)$, we deduce that $\Delta(b)$ has a $\bar{\Delta}$ -flag with sections $\bar{\Delta}(c)$ for c with $\rho(c) = \rho(b)$. Similarly, applying j_*^λ , we get that $\nabla(b)$ has a $\bar{\nabla}$ -flag with sections $\bar{\nabla}(c)$ for c with $\rho(c) = \rho(b)$.

To show that \mathcal{R} is $+$ -stratified, we check that each $I(b)$ has a $\bar{\nabla}$ -flag with sections $\bar{\nabla}(c)$ for c with $\rho(c) \geq \rho(b)$. This is immediate if $\varepsilon(b) = +$ since we are assuming that \mathcal{R} is ε -stratified. If $\varepsilon(b) = -$ then $I(b)$ has a ∇ -flag with sections $\nabla(c)$ for c with $\rho(c) \geq \rho(b)$. Hence, by the previous paragraph, it also has the required sort of $\bar{\nabla}$ -flag.

(v) \Rightarrow (i): We just need to show that \mathcal{R} is $-$ -stratified. We know that each $P(b)$ has a Δ -flag with sections $\Delta(c)$ for c with $\rho(c) \geq \rho(b)$. Now use the given $\bar{\Delta}$ -flags of each $\Delta(c)$ to see that each $P(b)$ also has the appropriate sort of $\bar{\Delta}$ -flag.

(v) \Leftrightarrow (vi): This follows from the above using the observation made earlier that \mathcal{R} is fully stratified if and only if \mathcal{R}^{op} is fully stratified. \square

Corollary 3.21. *If \mathcal{R} is an ε -stratified category with a contravariant autoequivalence which preserves isomorphism classes of irreducible objects, then \mathcal{R} is fully stratified. Moreover, if \mathcal{R} is an ε -highest weight category with a contravariant autoequivalence preserving isomorphism classes of irreducible objects, then \mathcal{R} is fibered highest weight.*

Proof. Since \mathcal{R} is ε -stratified, \mathcal{R}^{op} is $(-\varepsilon)$ -stratified. Using Lemma 3.3, we deduce that \mathcal{R} is $(-\varepsilon)$ -stratified. This verifies Lemma 3.20(iii) and the first claim follows. The second is then obvious. \square

Lemma 3.22. *Suppose that \mathcal{R} is a finite or essentially finite fully stratified category. For $b, c \in \mathbf{B}$ and $n \geq 0$, we have that*

$$\text{Ext}_{\mathcal{R}}^n(\bar{\Delta}(b), \bar{\nabla}(c)) \cong \begin{cases} \text{Ext}_{\mathcal{R}_\lambda}^n(L(b), L(c)) & \text{if } \lambda = \mu \\ 0 & \text{otherwise,} \end{cases}$$

where $\lambda := \rho(b)$ and $\mu := \rho(c)$.

Proof. Choose ε so that $\varepsilon(\lambda) = -$, hence, $\bar{\Delta}(b) = \Delta_\varepsilon(b)$. By Lemma 3.20, \mathcal{R} is ε -stratified, so we can apply Theorem 3.17(4) with $\mathcal{R}^\dagger = \mathcal{R}_{\leq \mu}$ to deduce that

$$\text{Ext}_{\mathcal{R}}^n(\bar{\Delta}(b), \bar{\nabla}(c)) \cong \text{Ext}_{\mathcal{R}_{\leq \mu}}^n(i_{\leq \mu}^* \bar{\Delta}(b), \bar{\nabla}(c)).$$

This is zero unless $\lambda \leq \mu$. If $\lambda \leq \mu$ it is $\text{Ext}_{\mathcal{R}_{\leq \mu}}^n(\bar{\Delta}(b), \bar{\nabla}(c))$. Now we change ε so that $\varepsilon(\mu) = +$, hence, $\bar{\nabla}(c) = \nabla_\varepsilon(c)$. Then by Theorem 3.18(3) with $\mathcal{R} = \mathcal{R}_{\leq \mu}$ and $\mathcal{R}^\dagger = \mathcal{R}_\mu$ we get that $\text{Ext}_{\mathcal{R}_{\leq \mu}}^n(\bar{\Delta}(b), \bar{\nabla}(c)) \cong \text{Ext}_{\mathcal{R}_\mu}^n(j^\mu \bar{\Delta}(b), L(c))$. This is zero unless $\lambda = \mu$, when $j^\mu \bar{\Delta}(b) = L(b)$ and we are done. \square

The next results are concerned with global dimension.

Lemma 3.23. *Let \mathcal{R} be a finite ε -stratified category.*

- (1) *All $V \in \Delta_\varepsilon(\mathcal{R})$ are of finite projective dimension if and only if all negative strata² have finite global dimension.*
- (2) *All $V \in \nabla_\varepsilon(\mathcal{R})$ are of finite injective dimension if and only if all positive strata have finite global dimension.*

Proof. As the two parts are dual statements, it suffices to prove (1). Replacing Λ by the finite set $\rho(\mathbf{B})$ if necessary, we may assume that $|\Lambda| < \infty$.

First assume that all negative strata have finite global dimension. By [Wei, Ex. 4.1.2], it suffices to show that $\text{pd } \Delta_\varepsilon(b) < \infty$ for each $b \in \mathbf{B}$. We proceed by downwards induction on the partial order on the finite poset Λ . Take any $\lambda \in \Lambda$ and consider $\Delta_\varepsilon(b)$ for $b \in \mathbf{B}_\lambda$, assuming that $\text{pd } \Delta_\varepsilon(c) < \infty$ for each $c \in \mathbf{B}_{>\lambda}$. We first observe that there is a short exact sequence $0 \rightarrow Q \rightarrow P(b) \rightarrow \Delta(b) \rightarrow 0$ such that Q has a Δ_ε -flag with sections $\Delta_\varepsilon(c)$ for $c \in \mathbf{B}_{>\lambda}$. If $\varepsilon(\lambda) = +$ this follows immediately from $(P\Delta_\varepsilon)$; if $\varepsilon(\lambda) = -$ one also needs to use (3.10) to see that a Δ_ε -flag in $P(b)$ can be ordered so that the sections $\bar{\Delta}(c)$ with $c \in \mathbf{B}_\lambda$ appear above the sections with $c \in \mathbf{B}_{>\lambda}$. By the induction hypothesis, Q has finite projective dimension, hence, so does $\Delta(b)$. This verifies the induction step in the case that $\varepsilon(\lambda) = +$. Instead, suppose that $\varepsilon(\lambda) = -$, i.e., $\Delta_\varepsilon(b) = \bar{\Delta}(b)$. Let $0 \rightarrow P_n \rightarrow \cdots \rightarrow P_0 \rightarrow L_\lambda(b) \rightarrow 0$ be a finite projective resolution of $L_\lambda(b)$ in the stratum \mathcal{R}_λ . Applying $j_!^\lambda$, which is exact thanks to Theorem 3.5, we obtain an exact sequence $0 \rightarrow V_n \rightarrow \cdots \rightarrow V_0 \rightarrow \bar{\Delta}(b) \rightarrow 0$ such that each V_m is a direct sum of standard objects $\Delta(c)$ for $c \in \mathbf{B}_\lambda$. The result already established plus [Wei, Ex. 4.1.3] implies that $\text{pd } V_m < \infty$ for each m . Arguing like in the proof of [Wei, Th. 4.3.1], we deduce that $\text{pd } \bar{\Delta}(b) < \infty$ too.

Conversely, suppose that $\text{pd } \Delta_\varepsilon(b) < \infty$ for all $b \in \mathbf{B}$. Take $\lambda \in \Lambda$ with $\varepsilon(\lambda) = -$. To show that \mathcal{R}_λ has finite global dimension, it suffices to show that there is some $d(\lambda) \geq 0$ such that $\text{Ext}_{\mathcal{R}_\lambda}^n(L_\lambda(b), W) = 0$ for all $n > d(\lambda)$, $b \in \mathbf{B}_\lambda$ and $W \in \mathcal{R}_\lambda$. By Theorems 3.18(3) and 3.17(3), we have that

$$\text{Ext}_{\mathcal{R}_\lambda}^n(L_\lambda(b), W) \cong \text{Ext}_{\mathcal{R}_{\leq \lambda}}^n(\Delta_\varepsilon(b), j_*^\lambda W) \cong \text{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(b), i_{\leq \lambda}(j_*^\lambda W)).$$

So we can take $d(\lambda) = \max\{\text{pd } \Delta_\varepsilon(b) \mid b \in \mathbf{B}_\lambda\}$. \square

The case when all strata are positive (respectively negative) will be of great importance.

Corollary 3.24. *If \mathcal{R} is a finite +-stratified (resp., --stratified) category then all $V \in \Delta(\mathcal{R})$ (resp., $V \in \nabla(\mathcal{R})$) are of finite projective (resp., injective) dimension.*

Corollary 3.25. *Suppose that \mathcal{R} is a finite ε -stratified category. If \mathcal{R} is of finite global dimension then all of its strata are of finite global dimension too.*

Proof. Lemma 3.23(1) implies that all negative strata have finite global dimension, and Lemma 3.23(2) implies that all positive strata have finite global dimension. \square

Corollary 3.26. *Suppose that \mathcal{R} is either a finite +-stratified category or a finite --stratified category. If all of the strata are of finite global dimension then \mathcal{R} is of finite global dimension.*

Proof. We just explain this in the case that \mathcal{R} is --stratified; the argument in the +-stratified case is similar. Lemma 3.23(1) implies that $\bar{\Delta}(b)$ is of finite projective dimension for each $b \in \mathbf{B}$. Moreover, there is a short exact sequence $0 \rightarrow K \rightarrow \bar{\Delta}(b) \rightarrow L(b) \rightarrow 0$ where all composition factors of K are of the form $L(c)$ for c with $\rho(c) < \rho(b)$. Ascending induction on the partial order on the finite set $\rho(\mathbf{B}) \subseteq \Lambda$ implies that each $L(b)$ has finite projective dimension. \square

²We mean the strata \mathcal{R}_λ for $\lambda \in \Lambda$ such that $\varepsilon(\lambda) = -$.

A special case of Corollary 3.26 recovers the following well-known result, see e.g. [CPS1]. For further detailed remarks about the history of this, and the general notion of highest weight category, we refer to [Don4, §A5] and [DR].

Corollary 3.27. *Finite highest weight categories are of finite global dimension.*

Remark 3.28. In the fully stratified case, Lemma 3.22 can be used to give a precise bound on the global dimension of \mathcal{R} in Corollary 3.26. Assuming Λ is finite, let

$$|\lambda| := \sup \left\{ \frac{\max(\text{gl. dim } \mathcal{R}_{\lambda_1}, \dots, \text{gl. dim } \mathcal{R}_{\lambda_n})}{2} + n - 1 \mid \begin{array}{l} n \geq 1 \text{ and } \lambda_1, \dots, \lambda_n \in \Lambda \\ \text{with } \lambda_1 < \dots < \lambda_n = \lambda \end{array} \right\}.$$

By mimicking the proof of [Don4, Prop. A2.3], one shows that $\text{Ext}_{\mathcal{R}}^i(L(b), L(c)) = 0$ for $b, c \in \mathbf{B}$ and any $i > |\rho(b)| + |\rho(c)|$. Hence, $\text{gl. dim } \mathcal{R} \leq 2 \max\{|\lambda| \mid \lambda \in \Lambda\}$. For finite highest weight categories, this shows that $\text{gl. dim } \mathcal{R} \leq 2(n-1)$ where n is length of the longest chain of weights in the weight poset Λ .

Remark 3.29. Outside of the highest weight case, *finitistic dimension* is used as a replacement for global dimension. In particular, finite fibered highest weight categories have finitistic dimension $\leq 2(n-1)$ where n is length of the longest chain of weights in the weight poset Λ ; this can be proved following the argument of [AHLU, Cor. 2.7]. For finite fully stratified categories, it should be possible to bound the finitistic dimension of \mathcal{R} in terms of the finitistic dimensions of the strata and chains in the poset like in the previous remark.

Remark 3.30. Another remarkable result about global dimension of finite highest weight categories was obtained in [MO], [MP] proving conjectures formulated in [CaeZ], [EP]: if \mathcal{R} is a finite highest weight category with duality, i.e., possessing a contravariant autoequivalence preserving isomorphism classes of irreducible objects, then the global dimension of \mathcal{R} is equal to twice the projective dimension of a tilting generator (see Definition 4.9 below). More generally, Mazorchuk and Ovsienko show that the finitistic dimension is equal to twice the projective dimension of a tilting generator in any finite fibered highest weight category with duality which is also tilting-rigid in the sense of Definition 4.36 below. Recently, Cruz and Marczinik [CM, Th. 2.2] (see also Corollary 4.40 below) have shown that a finite fibered highest weight category \mathcal{R} is tilting-rigid if and only if it is Gorenstein, in which case the finitistic dimension of \mathcal{R} coincides with its Gorenstein dimension (e.g., see [Che, Lem. 2.3.2]).

3.3. Upper finite ε -stratified categories. In this subsection we assume that \mathcal{R} is a Schurian category equipped with an upper finite stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Also $\varepsilon : \Lambda \rightarrow \{\pm\}$ denotes a sign function. Let $I(b)$ and $P(b)$ be an injective hull and a projective cover of $L(b)$ in \mathcal{R} . Recall (1.1)–(1.2), the properties $(P\Delta_\varepsilon)\text{--}(I\nabla_\varepsilon)$ and $(P\Delta)\text{--}(I\nabla)$ from the introduction, and the seemingly stronger properties $(\widehat{P\Delta}_\varepsilon)\text{--}(\widehat{I\nabla}_\varepsilon)$ and $(\widehat{P\Delta})\text{--}(\widehat{I\nabla})$ from the previous subsection.

Before formulating the main definitions in the upper finite setting, we prove an analog of the homological criterion for ∇_ε -flags from Theorem 3.11. The proof depends on the lemmas proved in §3.4 below, which we used already in the previous subsection, together with the following two technical lemmas, which we prove by truncating to finite Abelian quotients.

Lemma 3.31. *Suppose that \mathcal{R} is Schurian with upper finite stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ and sign function ε , and assume that the property $(\widehat{P\Delta}_\varepsilon)$ holds in \mathcal{R} . Let Λ^\dagger be a finite upper set in Λ , $\mathbf{B}^\dagger := \rho^{-1}(\Lambda^\dagger)$, and $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be the corresponding Serre quotient category with the induced stratification. The functor j_* sends short exact sequences of objects in $\nabla_\varepsilon(\mathcal{R}^\dagger)$ to short exact sequences of objects in $\nabla_\varepsilon(\mathcal{R})$.*

Proof (assuming lemmas in §3.4 below). Take a short exact sequence $0 \rightarrow K \rightarrow X \rightarrow Q \rightarrow 0$ in \mathcal{R}^\dagger such that K, X and Q have ∇_ε -flags. We must show that $0 \rightarrow j_*K \rightarrow$

$j_*X \rightarrow j_*Q \rightarrow 0$ is exact with all objects belonging to $\nabla_\varepsilon(\mathcal{R})$. We proceed by induction on the length of the ∇_ε -flag of $j_*(X)$, with the base case (length one) following from Lemma 3.49(1). For the induction step, we may assume that $K, Q \neq 0$ and know by induction that j_*K and j_*Q have ∇_ε -flags. We must show that $0 \rightarrow j_*K \rightarrow j_*X \rightarrow j_*Q \rightarrow 0$ is exact. Since j_* is left exact, this follows if we can show that

$$[j_*X : L(b)] = [j_*K : L(b)] + [j_*Q : L(b)]$$

for all $b \in \mathbf{B}$. To see this, let Λ^\dagger be the finite upper set generated by Λ^\dagger and b . Let $\mathbf{B}^\dagger := \rho^{-1}(\Lambda^\dagger)$ and $k : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be the corresponding Serre quotient. By Lemma 2.27, we have that $[j_*X : L(b)] = [k(j_*X) : kL(b)] = [k(j_*X) : L^\dagger(b)]$, and similarly for K and Q . Since Λ^\dagger is an upper set in Λ^\dagger , we can also view \mathcal{R}^\dagger as a quotient of \mathcal{R}^\dagger , and the quotient functor j factors as $j = \bar{j} \circ k$ for another quotient functor $\bar{j} : \mathcal{R}^\dagger \rightarrow \mathcal{R}^\dagger$. We have that $k_* \circ \bar{j}_* \cong j_*$, hence, applying k , we get that $\bar{j}_* \cong k \circ j_*$. It follows that $[k(j_*X) : L^\dagger(b)] = [\bar{j}_*X : L^\dagger(b)]$, and similarly for K and Q . We have now reduced the proof to showing that

$$[\bar{j}_*X : L^\dagger(b)] = [\bar{j}_*K : L^\dagger(b)] + [\bar{j}_*Q : L^\dagger(b)].$$

To see this, we note that \mathcal{R}^\dagger and \mathcal{R}^\dagger are finite ε -highest weight categories due to Lemma 3.49(2) and Theorem 3.5. So we can apply Theorem 3.18(4) to see that the sequence $0 \rightarrow \bar{j}_*K \rightarrow \bar{j}_*X \rightarrow \bar{j}_*Q \rightarrow 0$ is exact. \square

Lemma 3.32. *Suppose that \mathcal{R} is Schurian with upper finite stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ and sign function ε , and assume that the property $(\widehat{P}\Delta_\varepsilon)$ holds in \mathcal{R} . Let $V \in \mathcal{R}$ be a finitely cogenerated object such that $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), V) = 0$ for all $b \in \mathbf{B}$. Then we have that $V \in \nabla_\varepsilon(b)$, and the multiplicity $(V : \nabla_\varepsilon(b))$ of $\nabla_\varepsilon(b)$ in any ∇_ε -flag is equal to the dimension of $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V)$.*

Proof (assuming lemmas from §3.4 below). Since V is finitely cogenerated, its injective hull is a finite direct sum of the indecomposable injective objects $I(b)$. This means that we can find a finite upper set Λ^\dagger and $\mathbf{B}^\dagger := \rho^{-1}(\Lambda^\dagger)$ so that there is a short exact sequence

$$0 \longrightarrow V \longrightarrow \bigoplus_{b \in \mathbf{B}^\dagger} I(b)^{\oplus n_b} \longrightarrow Q \longrightarrow 0$$

for some $n_b \geq 0$. Let $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be the corresponding Serre quotient. This is a finite ε -stratified category by Lemma 3.49(2) and Theorem 3.5.

Applying j to the above short exact sequence gives us a short exact sequence in \mathcal{R}^\dagger . Then we take $b \in \mathbf{B}^\dagger$ and apply the functor $\text{Hom}_{\mathcal{R}^\dagger}(\Delta_\varepsilon^\dagger(b), ?)$ to this using also Lemma 3.49(1) to obtain the long exact sequence

$$\begin{aligned} 0 \longrightarrow \text{Hom}_{\mathcal{R}^\dagger}(\Delta_\varepsilon^\dagger(b), jV) &\longrightarrow \text{Hom}_{\mathcal{R}^\dagger}(\Delta_\varepsilon^\dagger(b), \bigoplus_{b \in \mathbf{B}^\dagger} I^\dagger(b)^{\oplus n_b}) \\ &\longrightarrow \text{Hom}_{\mathcal{R}^\dagger}(\Delta_\varepsilon^\dagger(b), jQ) \longrightarrow \text{Ext}_{\mathcal{R}^\dagger}^1(\Delta_\varepsilon^\dagger(b), jV) \longrightarrow 0. \end{aligned}$$

From adjunction and Lemma 3.49(1) again, we get a commuting diagram

$$\begin{array}{ccccc} 0 \rightarrow \text{Hom}_{\mathcal{R}^\dagger}(\Delta_\varepsilon^\dagger(b), jV) & \rightarrow & \text{Hom}_{\mathcal{R}^\dagger}(\Delta_\varepsilon^\dagger(b), \bigoplus_{b \in \mathbf{B}^\dagger} I^\dagger(b)^{\oplus n_b}) & \rightarrow & \text{Hom}_{\mathcal{R}^\dagger}(\Delta_\varepsilon^\dagger(b), jQ) \rightarrow 0 \\ \downarrow & & \downarrow & & \downarrow \\ 0 \rightarrow \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) & \rightarrow & \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), \bigoplus_{b \in \mathbf{B}^\dagger} I(b)^{\oplus n_b}) & \rightarrow & \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), Q) \rightarrow 0. \end{array}$$

The vertical maps are isomorphisms and the bottom row is exact since $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), V) = 0$. Hence the top row is exact. Comparing with the previously displayed long exact sequence, it follows that $\text{Ext}_{\mathcal{R}^\dagger}^1(\Delta_\varepsilon^\dagger(b), jV) = 0$. Now we can apply Theorem 3.11 in the finite ε -stratified category \mathcal{R}^\dagger to deduce that jV has a ∇_ε -flag.

From Lemma 3.31, we deduce that j_*jV has a ∇_ε -flag. Moreover the multiplicity of $\nabla_\varepsilon(b)$ in any ∇_ε -flag in j_*jV is $\dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), j_*jV)$ thanks to Lemma 3.48. To

complete the proof, we show that the unit of adjunction $f : V \rightarrow j_*j^*V$ is an isomorphism. We know from Lemma 3.49(1) that the unit of adjunction is an isomorphism $I(b) \rightarrow j_*j^*I(b)$ for each $b \in \mathbf{B}^\dagger$. Since V embeds into a direct sum of such $I(b)$, it follows that f is injective. To show that it is surjective, it suffices to show that

$$[j_*j^*V : L(b)] = [V : L(b)]$$

for all $b \in \mathbf{B}$. To prove this, we fix a choice of $b \in \mathbf{B}$ then define $\Lambda^\dagger, \mathbf{B}^\dagger, k : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ and $\bar{j} : \mathcal{R}^\dagger \rightarrow \mathcal{R}^\dagger$ as in the proof of Lemma 3.31. Since $b \in \mathbf{B}^\dagger$, we have that $[V : L(b)] = [kV : L^\dagger(b)]$ and $[j_*j^*V : L(b)] = [k(j_*j^*V) : L^\dagger(b)]$. As in the proof of Lemma 3.31, $k(j_*j^*V) \cong \bar{j}_*(j^*V) \cong \bar{j}_*(kV)$. Thus, we are reduced to showing that

$$[\bar{j}_*(kV) : L^\dagger(b)] = [kV : L^\dagger(b)].$$

This follows because $kV \cong \bar{j}_*(kV)$. To see this, we repeat the arguments in the previous paragraph to show that $kV \in \mathcal{R}^\dagger$ has a ∇_ε -flag. Since the unit of adjunction is an isomorphism $\nabla_\varepsilon^\dagger(b) \xrightarrow{\sim} \bar{j}_*\bar{j}^\dagger\nabla_\varepsilon^\dagger(b)$ for each $b \in \mathbf{B}^\dagger$, we deduce using the exactness from Theorem 3.18(4) that it gives an isomorphism $kV \xrightarrow{\sim} \bar{j}_*(kV)$ too. \square

Theorem 3.33. *Theorem 3.5 holds in the upper finite setup too.*

Proof. This is almost the same as the proof of Theorem 3.5 given in the previous subsection. One needs to use Lemma 3.32 in place of Theorem 3.11 to see that $I(b)$ has a ∇_ε -flag with the appropriate multiplicities. The exactness of j_*^λ when $\varepsilon(\lambda) = +$ follows from Lemma 3.31 applied to the quotient functor $j^\lambda : \mathcal{R}_{\leq \lambda} \rightarrow \mathcal{R}_\lambda$. Note for this that $\mathcal{R}_{\leq \lambda}$ satisfies $(\widehat{P\Delta}_\varepsilon)$ due to Lemma 3.45(2), and we have that $\nabla_\varepsilon(\mathcal{R}_\lambda) = \mathcal{R}_\lambda$ as $\varepsilon(\lambda) = +$. \square

We are ready to proceed to the main definition.

Definition 3.34. Let $(\mathbf{B}, L, \rho, \Lambda, \leq)$ be an upper finite stratification on \mathcal{R} .

- (ε S) We say that \mathcal{R} is an *upper finite ε -stratified category* if one of the equivalent properties $(\widehat{P\Delta}_\varepsilon)$ – $(\widehat{I\nabla}_\varepsilon)$ holds for a given choice of sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$.
- (FS) We say that \mathcal{R} is an *upper finite fully stratified category* if one of these properties holds for all choices of sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$.
- (ε HW) We say that \mathcal{R} is an *upper finite ε -highest weight category* if the stratification function ρ is a bijection, and one of the equivalent properties $(\widehat{P\Delta}_\varepsilon)$ – $(\widehat{I\nabla}_\varepsilon)$ holds for a given choice of sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$.
- (FHW) We say that \mathcal{R} is an *upper finite fibered highest weight category* if the stratification function is a bijection and one of these properties holds for all choices of sign function.
- (HW) We say that \mathcal{R} is an *upper finite highest weight category* if all of the stata are simple (cf. Lemma 3.4) and one of the equivalent properties $(\widehat{P\Delta})$ – $(\widehat{I\nabla})$ holds.

The Ext^1 -vanishing (3.10) and Theorem 3.9 both still hold in the same way as before.

Next we are going to consider two (in fact dual) notions of ascending Δ_ε - and descending ∇_ε -flags, generalizing the finite flags discussed already. One might be tempted to say that an ascending Δ_ε -flag in V is an ascending chain $0 = V_0 < V_1 < V_2 < \dots$ of subobjects of V with $V = \sum_{n \in \mathbb{N}} V_n$ such that $V_m/V_{m-1} \cong \Delta_\varepsilon(b_m)$, and a descending ∇_ε -flag is a descending chain $V = V_0 > V_1 > V_2 > \dots$ of subobjects of V such that $\bigcap_{n \in \mathbb{N}} V_n = 0$ and $V_{m-1}/V_m \cong \Delta_\varepsilon(b_m)$, for $b_m \in \mathbf{B}$. These would be serviceable definitions when Λ is countable. In order to avoid this unnecessary restriction, we will work instead with the following more general formulations.

Definition 3.35. Suppose that \mathcal{R} is an upper finite ε -stratified category and $V \in \mathcal{R}$.

- (A Δ) An *ascending Δ_ε -flag* in V is the data of a directed set Ω with smallest element 0 and a direct system $(V_\omega)_{\omega \in \Omega}$ of subobjects of V such that $V_0 = 0$, $\sum_{\omega \in \Omega} V_\omega = V$, and $V_v/V_\omega \in \Delta_\varepsilon(\mathcal{R})$ for each $\omega < v$. Let $\Delta_\varepsilon^{\text{asc}}(\mathcal{R})$ be the full subcategory of \mathcal{R} consisting of all objects V possessing such a flag.

(D ∇) A *descending ∇_ε -flag* in V is the data of a directed set Ω with smallest element 0 and an inverse system $(V/V_\omega)_{\omega \in \Omega}$ of quotients of V such that $V_0 = V$, $\bigcap_{\omega \in \Omega} V_\omega = 0$, and $V_\omega/V_\nu \in \nabla_\varepsilon(\mathcal{R})$ for each $\omega < \nu$. Let $\nabla_\varepsilon^{\text{dsc}}(\mathcal{R})$ be the full subcategory of \mathcal{R} consisting of all objects V possessing such a flag.

We stress that $\Delta_\varepsilon^{\text{asc}}(\mathcal{R})$ and $\nabla_\varepsilon^{\text{dsc}}(\mathcal{R})$ are subcategories of \mathcal{R} : we have *not* passed to the completion $\text{Ind}(\mathcal{R}_c)$.

Lemma 3.36. *Suppose that \mathcal{R} is an upper finite ε -stratified category.*

- (1) For $V \in \Delta_\varepsilon^{\text{asc}}(\mathcal{R})$, $W \in \nabla_\varepsilon^{\text{dsc}}(\mathcal{R})$ and $n \geq 1$, we have that $\text{Ext}_{\mathcal{R}}^n(V, W) = 0$.
(2) For $V \in \Delta_\varepsilon^{\text{asc}}(\mathcal{R})$ the multiplicity of $\Delta_\varepsilon(b)$ in a Δ_ε -flag may be defined from

$$(V : \Delta_\varepsilon(b)) := \dim \text{Hom}_{\mathcal{R}}(V, \nabla_\varepsilon(b)) = \sup \{(V_\omega : \Delta_\varepsilon(b)) \mid \omega \in \Omega\} < \infty,$$

where $(V_\omega)_{\omega \in \Omega}$ is any choice of ascending Δ_ε -flag.

- (3) For $V \in \nabla_\varepsilon^{\text{dsc}}(\mathcal{R})$, the multiplicity of $\nabla_\varepsilon(b)$ in a ∇_ε -flag may be defined from

$$(V : \nabla_\varepsilon(b)) := \dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) = \sup \{(V/V_\omega : \nabla_\varepsilon(b)) \mid \omega \in \Omega\} < \infty,$$

where $(V/V_\omega)_{\omega \in \Omega}$ is any choice of descending ∇_ε -flag.

Proof. (1) We first prove this in the special case that $W = \nabla_\varepsilon(b)$. Let $(V_\omega)_{\omega \in \Omega}$ be an ascending Δ_ε -flag in V , so that $V \cong \varprojlim V_\omega$. Since $\text{Ext}_{\mathcal{R}}^n(V_\omega, W) = 0$ by Lemma 3.48, it suffices to show that

$$\text{Ext}_{\mathcal{R}}^n(V, W) \cong \varprojlim \text{Ext}_{\mathcal{R}}^n(V_\omega, W).$$

To see this, like in [Wei, 3.5.10], we need to check a Mittag-Leffler condition. We show that the natural map $\text{Ext}_{\mathcal{R}}^{n-1}(V_\nu, W) \rightarrow \text{Ext}_{\mathcal{R}}^{n-1}(V_\omega, W)$ is surjective for each $\omega < \nu$ in Ω . Applying $\text{Hom}_{\mathcal{R}}(?, W)$ to the short exact sequence $0 \rightarrow V_\omega \rightarrow V_\nu \rightarrow V_\nu/V_\omega \rightarrow 0$ gives an exact sequence

$$\text{Ext}_{\mathcal{R}}^{n-1}(V_\nu, W) \longrightarrow \text{Ext}_{\mathcal{R}}^{n-1}(V_\omega, W) \longrightarrow \text{Ext}_{\mathcal{R}}^n(V_\nu/V_\omega, W).$$

It remains to observe that $\text{Ext}_{\mathcal{R}}^n(V_\nu/V_\omega, W) = 0$ by Lemma 3.48 again, since we know from the definition of ascending Δ_ε -flag that $V_\nu/V_\omega \in \Delta_\varepsilon(\mathcal{R})$.

The dual of the previous paragraph plus Lemma 2.12 gives that $\text{Ext}_{\mathcal{R}}^n(V, W) = 0$ for $n \geq 1$, $V = \Delta_\varepsilon(b)$ and $W \in \nabla_\varepsilon^{\text{dsc}}(\mathcal{R})$. Then we can repeat the argument of the previous paragraph yet again, using this assertion in place of Lemma 3.48, to obtain the result we are after for general $V \in \Delta_\varepsilon^{\text{asc}}(\mathcal{R})$ and $W \in \nabla_\varepsilon^{\text{dsc}}(\mathcal{R})$.

- (2) This follows from (1) and (3.5) because

$$\text{Hom}_{\mathcal{R}}(V, \nabla_\varepsilon(b)) \cong \text{Hom}_{\mathcal{R}}(\varprojlim V_\omega, \nabla_\varepsilon(b)) \cong \varprojlim \text{Hom}_{\mathcal{R}}(V_\omega, \nabla_\varepsilon(b)),$$

which is finite-dimensional as $\nabla_\varepsilon(b)$, hence, each V_ω , is finitely cogenerated.

- (3) Similarly to (2), we have that

$$\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) \cong \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), \varprojlim (V/V_\omega)) \cong \varprojlim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V/V_\omega),$$

which is finite-dimensional as $\Delta_\varepsilon(b)$ is finitely generated. Then we can apply (1) and (3.5) once again. \square

Theorem 3.37 (Homological criterion for ascending Δ_ε -flags). *Assume that \mathcal{R} is an upper finite ε -stratified category. For $V \in \mathcal{R}$, the following are equivalent:*

- (i) $V \in \Delta_\varepsilon^{\text{asc}}(\mathcal{R})$;
(ii) $\text{Ext}_{\mathcal{R}}^1(V, \nabla_\varepsilon(b)) = 0$ for all $b \in \mathbf{B}$;
(iii) $\text{Ext}_{\mathcal{R}}^n(V, \nabla_\varepsilon(b)) = 0$ for all $b \in \mathbf{B}$ and $n \geq 1$.

Assuming these properties, we have that $V \in \Delta_\varepsilon(\mathcal{R})$ if and only if it is finitely generated.

Proof. (iii) \Rightarrow (ii). Trivial.

(i) \Rightarrow (iii). This follows from Lemma 3.36(1).

(ii) \Rightarrow (i). Let Ω be the directed set of finite upper sets in Λ . Take $\omega \in \Omega$; it is some finite upper set Λ^\dagger . Let $\mathbf{B}^\dagger := \rho^{-1}(\Lambda^\dagger)$ and $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be the corresponding Serre quotient. By Lemma 3.49(3), $\text{Ext}_{\mathcal{R}^\dagger}^1(jV, \nabla_\varepsilon(b)) = 0$ for all $b \in \mathbf{B}^\dagger$. Hence, $V_\omega := j_!(jV) \in \Delta_\varepsilon(\mathcal{R})$ thanks to the dual of Lemma 3.31. Let $f_\omega : V_\omega \rightarrow V$ be the morphism induced by the counit of adjunction. We claim for any $b \in \mathbf{B}^\dagger$ that the map

$$f_\omega(b) : \text{Hom}_{\mathcal{R}}(P(b), V_\omega) \rightarrow \text{Hom}_{\mathcal{R}}(P(b), V), \theta \mapsto f_\omega \circ \theta$$

is an isomorphism. To see this, we assume that $\mathcal{R} = A\text{-mod}_{\text{fd}}$ for a pointed locally finite-dimensional locally unital algebra $A = \bigoplus_{a,b \in \mathbf{B}} e_a A e_b$. Then $\mathcal{R}^\dagger = eAe\text{-mod}_{\text{fd}}$ where $e = \sum_{a \in \mathbf{B}^\dagger} e_a$, and $V_\omega = Ae \otimes_{eAe} eV$. In these terms, the map f_ω is the natural multiplication map. For $b \in \mathbf{B}^\dagger$, this multiplication map gives an isomorphism $e_b V_\omega \xrightarrow{\sim} e_b V$ with inverse $e_b v \mapsto e_b \otimes e_b v$. This proves the claim.

Now take $v > \omega$, i.e., another finite upper set $\Lambda^{\dagger\dagger} \supset \Lambda^\dagger$, and let $k : \mathcal{R} \rightarrow \mathcal{R}^{\dagger\dagger}$ be the associated quotient. The quotient functor $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ factors as $j = \bar{j} \circ k$ for another quotient functor $\bar{j} : \mathcal{R}^{\dagger\dagger} \rightarrow \mathcal{R}^\dagger$, and we have that

$$V_\omega = (\bar{j} \circ k)_!(kV) \cong k_!(\bar{j}_!(kV)), \quad V_v = k_!(kV).$$

By Corollary 3.19(2), there is a short exact sequence $0 \rightarrow \bar{j}_!(kV) \rightarrow kV \rightarrow Q \rightarrow 0$ such that both $\bar{j}_!(kV)$ and Q belong to $\Delta_\varepsilon(\mathcal{R}^{\dagger\dagger})$. Applying $k_!$ and using the exactness from the dual of Lemma 3.31, we get an embedding $f_\omega^v : V_\omega \hookrightarrow V_v$ such that $V_v/V_\omega \cong k_!Q \in \Delta_\varepsilon(\mathcal{R})$. Since the morphisms all came from counits of adjunction, we have that $f_v \circ f_\omega^v = f_\omega$.

Now we can show that each f_ω is a monomorphism. It suffices to show that $f_\omega(b) : \text{Hom}_{\mathcal{R}}(P(b), V_\omega) \rightarrow \text{Hom}_{\mathcal{R}}(P(b), V)$ is injective for all $b \in \mathbf{B}$. Choose v in the previous paragraph to be sufficiently large so as to ensure that $b \in \mathbf{B}^{\dagger\dagger}$. We explained already that $f_v(b)$ is an isomorphism. Since $f_\omega = f_v \circ f_\omega^v$ and f_ω^v is a monomorphism, it follows that $f_\omega(b)$ is injective too. Thus, identifying V_ω with its image under f_ω , we have defined a direct system $(V_\omega)_{\omega \in \Omega}$ of subobjects of V such that $V_v/V_\omega \in \Delta_\varepsilon(\mathcal{R})$ for each $\omega < v$. It remains to observe that $V_\emptyset = 0$ for a trivial reason, and $\sum_{\omega \in \Omega} V_\omega = V$ because we know for each $b \in \mathbf{B}$ that $f_\omega(b)$ is surjective for sufficiently large ω .

Final part: If $V \in \Delta_\varepsilon(\mathcal{R})$, it is obvious that it is finitely generated since each $\Delta_\varepsilon(b)$ is finitely generated. Conversely, suppose that V is finitely generated and has an ascending Δ_ε -flag. To see that it is actually a finite flag, it suffices to show that $\text{Hom}_{\mathcal{R}}(V, \nabla_\varepsilon(b)) = 0$ for all but finitely many $b \in \mathbf{B}$. Say $\text{hd } V \cong L(b_1) \oplus \cdots \oplus L(b_n)$. If $V \rightarrow \nabla_\varepsilon(b)$ is a non-zero homomorphism, we must have that $\rho(b_i) \leq \rho(b)$ for some $i = 1, \dots, n$. Hence, there are only finitely many choices for b as the poset is upper finite. \square

Corollary 3.38. *Let $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ be a short exact sequence in \mathcal{R} .*

- (1) *If U and W belong to $\Delta_\varepsilon^{\text{asc}}(\mathcal{R})$ (resp., $\Delta_\varepsilon(\mathcal{R})$) so does V .*
- (2) *If V and W belong to $\Delta_\varepsilon^{\text{asc}}(\mathcal{R})$ (resp., $\Delta_\varepsilon(\mathcal{R})$) so does U .*

Theorem 3.39 (Homological criterion for descending ∇_ε -flags). *Assume that \mathcal{R} is an upper finite ε -stratified category. For $V \in \mathcal{R}$, the following are equivalent:*

- (i) $V \in \nabla_\varepsilon^{\text{dsc}}(\mathcal{R})$;
- (ii) $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), V) = 0$ for all $b \in \mathbf{B}$;
- (iii) $\text{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(b), V) = 0$ for all $b \in \mathbf{B}$ and $n \geq 1$.

Assuming these properties, $V \in \nabla_\varepsilon(\mathcal{R})$ if and only if it is finitely cogenerated.

Proof. This is the equivalent dual statement to Theorem 3.37. \square

Corollary 3.40. *Let $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ be a short exact sequence in \mathcal{R} .*

- (1) *If U and W belong to $\nabla_\varepsilon^{\text{dsc}}(\mathcal{R})$ (resp., $\nabla_\varepsilon(\mathcal{R})$) so does V .*
- (2) *If U and V belong to $\nabla_\varepsilon^{\text{dsc}}(\mathcal{R})$ (resp., $\nabla_\varepsilon(\mathcal{R})$) so does W .*

The following is the upper finite analog of Theorem 3.17; we have dropped part (6) since the proof of that required objects of \mathcal{R}^\downarrow to have finite length.

Theorem 3.41 (Truncation to lower sets). *Assume that \mathcal{R} is an upper finite ε -stratified category. Suppose that Λ^\downarrow is a lower set in Λ . Let $\mathbf{B}^\downarrow := \rho^{-1}(\Lambda^\downarrow)$ and $i : \mathcal{R}^\downarrow \rightarrow \mathcal{R}$ be the corresponding Serre subcategory of \mathcal{R} with the induced stratification. Then \mathcal{R}^\downarrow is an upper finite ε -stratified category. Moreover:*

- (1) *The distinguished objects in \mathcal{R}^\downarrow satisfy $L^\downarrow(b) \cong L(b)$, $P^\downarrow(b) \cong i^*P(b)$, $I^\downarrow(b) \cong i^!I(b)$, $\Delta^\downarrow(b) \cong \Delta(b)$, $\bar{\Delta}^\downarrow(b) \cong \bar{\Delta}(b)$, $\nabla^\downarrow(b) \cong \nabla(b)$ and $\bar{\nabla}^\downarrow(b) \cong \bar{\nabla}(b)$ for $b \in \mathbf{B}^\downarrow$.*
- (2) *i^* sends short exact sequences of objects in $\Delta_\varepsilon(\mathcal{R})$ to short exact sequences, $i^*\Delta(b) \cong \Delta^\downarrow(b)$ and $i^*\bar{\Delta}(b) \cong \bar{\Delta}^\downarrow(b)$ for $b \in \mathbf{B}^\downarrow$, and $i^*\Delta(b) = i^*\bar{\Delta}(b) = 0$ for $b \notin \mathbf{B}^\downarrow$.*
- (3) *$\text{Ext}_{\mathcal{R}}^n(V, iW) \cong \text{Ext}_{\mathcal{R}^\downarrow}^n(i^*V, W)$ for $V \in \Delta_\varepsilon(\mathcal{R})$, $W \in \mathcal{R}^\downarrow$ and all $n \geq 0$.*
- (4) *$i^!$ sends short exact sequences of objects in $\nabla_\varepsilon(\mathcal{R})$ to short exact sequences, $i^!\nabla(b) \cong \nabla^\downarrow(b)$ and $i^!\bar{\nabla}(b) \cong \bar{\nabla}^\downarrow(b)$ for $b \in \mathbf{B}^\downarrow$, and $i^!\nabla(b) = i^!\bar{\nabla}(b) = 0$ for $b \notin \mathbf{B}^\downarrow$.*
- (5) *$\text{Ext}_{\mathcal{R}}^n(iV, W) \cong \text{Ext}_{\mathcal{R}^\downarrow}^n(V, i^!W)$ for $V \in \mathcal{R}^\downarrow$, $W \in \nabla_\varepsilon(\mathcal{R})$ and all $n \geq 0$.*

Proof. This follows from Lemma 3.45 and the dual statement. \square

Next is the upper finite analog of Theorem 3.18.

Theorem 3.42 (Truncation to upper sets). *Assume that \mathcal{R} is an upper finite ε -stratified category. Suppose that Λ^\uparrow is an upper set in Λ . Let $\mathbf{B}^\uparrow := \rho^{-1}(\Lambda^\uparrow)$ and $j : \mathcal{R} \rightarrow \mathcal{R}^\uparrow$ be the corresponding Serre quotient category of \mathcal{R} with the induced stratification. Then \mathcal{R}^\uparrow is itself a finite or upper finite ε -stratified category according to whether Λ^\uparrow is finite or infinite. Moreover:*

- (1) *For $b \in \mathbf{B}^\uparrow$, the distinguished objects $L^\uparrow(b)$, $P^\uparrow(b)$, $I^\uparrow(b)$, $\Delta^\uparrow(b)$, $\bar{\Delta}^\uparrow(b)$, $\nabla^\uparrow(b)$ and $\bar{\nabla}^\uparrow(b)$ in \mathcal{R}^\uparrow are isomorphic to the images under j of the corresponding objects of \mathcal{R} .*
- (2) *We have that $jL(b) = j\Delta(b) = j\bar{\Delta}(b) = j\nabla(b) = j\bar{\nabla}(b) = 0$ if $b \notin \mathbf{B}^\uparrow$.*
- (3) *$\text{Ext}_{\mathcal{R}}^n(V, j_*W) \cong \text{Ext}_{\mathcal{R}^\uparrow}^n(jV, W)$ for $V \in \mathcal{R}$, $W \in \nabla_\varepsilon^{\text{dsc}}(\mathcal{R}^\uparrow)$ and all $n \geq 0$.*
- (4) *j_* sends short exact sequences of objects in $\nabla_\varepsilon(\mathcal{R}^\uparrow)$ to short exact sequences, $j_*\nabla^\uparrow(b) \cong \nabla(b)$, $j_*\bar{\nabla}^\uparrow(b) \cong \bar{\nabla}(b)$ and $j_*I^\uparrow(b) \cong I(b)$ for $b \in \mathbf{B}^\uparrow$.*
- (5) *$\text{Ext}_{\mathcal{R}^\uparrow}^n(jV, W) \cong \text{Ext}_{\mathcal{R}}^n(V, jW)$ for $V \in \Delta_\varepsilon^{\text{asc}}(\mathcal{R}^\uparrow)$, $W \in \mathcal{R}$ and all $n \geq 0$.*
- (6) *$j_!$ sends short exact sequences of objects in $\Delta_\varepsilon(\mathcal{R}^\uparrow)$ to short exact sequences, $j_!\Delta^\uparrow(b) \cong \Delta(b)$, $j_!\bar{\Delta}^\uparrow(b) \cong \bar{\Delta}(b)$ and $j_!P^\uparrow(b) = P(b)$ for $b \in \mathbf{B}^\uparrow$.*

Proof. If Λ^\uparrow is finite, this is proved in just the same way as Theorem 3.18. Assume instead that Λ^\uparrow is infinite. Then the same arguments prove (1) and (2), but the proofs of the remaining parts need some slight modifications. It suffices to prove (3) and (4), since (5) and (6) are the same results for \mathcal{R}^{op} .

For (3), the argument from the proof of Lemma 3.49(3) reduces to checking that j sends projectives to objects that are acyclic for $\text{Hom}_{\mathcal{R}^\uparrow}(?, W)$. To see this, it suffices to show that $\text{Ext}_{\mathcal{R}^\uparrow}^n(jP(b), W) = 0$ for $n \geq 1$ and $b \in \mathbf{B}$, which follows from Lemma 3.36(1).

Finally, for (4), the argument from the proof of Theorem 3.18(4) cannot be used since it depends on \mathcal{R}^\uparrow being essentially finite Abelian. So we provide an alternate argument. Take a short exact sequence $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ in $\nabla_\varepsilon(\mathcal{R}^\uparrow)$. Applying j_* , we get $0 \rightarrow j_*U \rightarrow j_*V \rightarrow j_*W$, and just need to show that the final morphism here is an epimorphism. This follows because, by (3) and Theorem 3.39, j_*U , j_*V and j_*W all have ∇_ε -flags such that $(j_*V : \nabla_\varepsilon(b)) = (j_*U : \nabla_\varepsilon(b)) + (j_*W : \nabla_\varepsilon(b))$ for all $b \in \mathbf{B}$. \square

The reader should have no difficulty in transporting Lemma 3.20 and Corollary 3.21 to the upper finite setting. Also, Lemma 3.23 remains valid when “finite ε -stratified category” is replaced by “upper finite ε -stratified category”. To see this, we just note that the argument by downwards induction on the partial order explained in the proof

works just as well when Λ is upper finite rather than finite. The following is the upper finite analog of Corollary 3.24.

Lemma 3.43. *If \mathcal{R} is an upper finite $+$ -stratified (resp., $-$ -stratified) category then all $V \in \Delta(\mathcal{R})$ (resp., $V \in \nabla(\mathcal{R})$) are of finite projective (resp., injective) dimension.*

Proof. This follows from the upper finite analog of Lemma 3.23. \square

3.4. Shared lemmas for §§3.2–3.3. In this subsection, we prove a series of lemmas needed in both §3.2 and in §3.3. Let \mathcal{R} be an Abelian category equipped with a stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ which is *either* essentially finite (§3.2) *or* upper finite (§3.3). Also let $\varepsilon : \Lambda \rightarrow \{\pm\}$ be a sign function. We assume throughout the subsection that the property $(\widehat{P\Delta}_\varepsilon)$ from §3.2 holds.

Lemma 3.44. *We have that $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), \Delta_\varepsilon(c)) = 0$ for $b, c \in \mathbf{B}$ such that $\rho(b) \not\leq \rho(c)$.*

Proof. Using the projective objects P_b given by the assumed property $(\widehat{P\Delta}_\varepsilon)$, we can construct the first terms of a projective resolution of $\Delta_\varepsilon(b)$ in the form

$$Q \longrightarrow \bigoplus_{\substack{a \in \mathbf{B} \\ \rho(a) \geq \rho(b)}} P_a^{\oplus n_a} \longrightarrow P_b \longrightarrow \Delta_\varepsilon(b) \longrightarrow 0 \quad (3.11)$$

for some $n_a \geq 0$. Now apply $\text{Hom}_{\mathcal{R}}(?, \Delta_\varepsilon(c))$ to get that $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), \Delta_\varepsilon(c))$ is the homology of the complex

$$\text{Hom}_{\mathcal{R}}(P_b, \Delta_\varepsilon(c)) \longrightarrow \text{Hom}_{\mathcal{R}}\left(\bigoplus_{\substack{a \in \mathbf{B} \\ \rho(a) \geq \rho(b)}} P_a^{\oplus n_a}, \Delta_\varepsilon(c)\right) \longrightarrow \text{Hom}_{\mathcal{R}}(Q, \Delta_\varepsilon(c)).$$

The middle term of this already vanishes as $[\Delta_\varepsilon(c) : L(a)] \neq 0 \Rightarrow \rho(a) \leq \rho(c)$. \square

Lemma 3.45. *Let Λ^\downarrow be a lower set in Λ and $\mathbf{B}^\downarrow := \rho^{-1}(\Lambda^\downarrow)$. Let $i : \mathcal{R}^\downarrow \rightarrow \mathcal{R}$ be the corresponding Serre subcategory of \mathcal{R} equipped with the induced stratification.*

- (1) *The standard, proper standard and indecomposable projective objects of \mathcal{R}^\downarrow are the objects $\Delta(b)$, $\widehat{\Delta}(b)$ and $i^*P(b)$ for $b \in \mathbf{B}^\downarrow$.*
- (2) *The object i^*P_b is zero unless $b \in \mathbf{B}^\downarrow$, in which case it is a projective object admitting a Δ_ε -flag with top section $\Delta_\varepsilon(b)$ and other sections of the form $\Delta_\varepsilon(c)$ for $c \in \mathbf{B}^\downarrow$ with $\rho(c) \geq \rho(b)$. In particular, this shows that $(\widehat{P\Delta}_\varepsilon)$ holds in \mathcal{R}^\downarrow .*
- (3) *$(\mathbb{L}_n i^*)V = 0$ for $V \in \Delta_\varepsilon(\mathcal{R})$ and $n \geq 1$.*
- (4) *$\text{Ext}_{\mathcal{R}}^n(V, iW) \cong \text{Ext}_{\mathcal{R}^\downarrow}^n(i^*V, W)$ for $V \in \Delta_\varepsilon(\mathcal{R})$, $W \in \mathcal{R}^\downarrow$ and $n \geq 0$.*

Proof. (1) For projectives, this follows from the usual adjunction properties. This also shows that i^*P_b is projective, as needed for (2). For standard and proper standard objects, just note that the standardization functors for \mathcal{R}^\downarrow are some of the ones for \mathcal{R} .

(2) Consider a Δ_ε -flag of P_b . Using Lemma 3.44, we can rearrange this filtration if necessary so that all of the sections $\Delta_\varepsilon(c)$ with $c \in \mathbf{B}^\downarrow$ appear above the sections $\Delta_\varepsilon(d)$ with $d \in \mathbf{B} \setminus \mathbf{B}^\downarrow$. So there exists a short exact sequence $0 \rightarrow K \rightarrow P_b \rightarrow Q \rightarrow 0$ in which Q has a finite filtration with sections $\Delta_\varepsilon(c)$ for $c \in \mathbf{B}^\downarrow$ with $\rho(c) \geq \rho(b)$, and K has a finite filtration with sections $\Delta_\varepsilon(c)$ for $c \in \mathbf{B} \setminus \mathbf{B}^\downarrow$. It follows easily that i^*P_b is isomorphic to Q , so it has the appropriate filtration.

(3) It suffices to show that $(\mathbb{L}_n i^*)\Delta_\varepsilon(b) = 0$ for all $b \in \mathbf{B}$ and $n > 0$. Take a short exact sequence $0 \rightarrow K \rightarrow P_b \rightarrow \Delta_\varepsilon(b) \rightarrow 0$ such that K has a Δ_ε -flag with sections $\Delta_\varepsilon(c)$ for c with $\rho(c) \geq \rho(b)$. Applying i^* , we obtain the long exact sequence

$$0 \longrightarrow (\mathbb{L}_1 i^*)\Delta_\varepsilon(b) \longrightarrow i^*K \longrightarrow i^*P_b \longrightarrow i^*\Delta_\varepsilon(b) \longrightarrow 0$$

and isomorphisms $(\mathbb{L}_{n+1} i^*)\Delta_\varepsilon(b) \cong (\mathbb{L}_n i^*)K$ for $n > 0$. We claim that $(\mathbb{L}_1 i^*)\Delta_\varepsilon(b) = 0$. We use Lemma 3.44 to order the Δ_ε -flag of K so that it yields a short exact sequence $0 \rightarrow L \rightarrow K \rightarrow Q \rightarrow 0$ in which Q has a Δ_ε -flag with sections $\Delta_\varepsilon(c)$ for $c \in \mathbf{B}^\downarrow$, and L has a Δ_ε -flag with sections $\Delta_\varepsilon(c)$ for $c \in \mathbf{B} \setminus \mathbf{B}^\downarrow$. It follows that $i^*K = Q$ and there

is a short exact sequence $0 \rightarrow i^*K \rightarrow i^*P_b \rightarrow \Delta_\varepsilon(b) \rightarrow 0$. Comparing with the long exact sequence, we deduce that $(\mathbb{L}_1 i^*)\Delta_\varepsilon(b) = 0$. Finally some degree shifting using the isomorphisms $(\mathbb{L}_{n+1} i^*)\Delta_\varepsilon(b) \cong (\mathbb{L}_n i^*)K$ gives that $(\mathbb{L}_n i^*)\Delta_\varepsilon(b) = 0$ for $n > 1$ too.

(4) By the adjunction, we have that $\mathrm{Hom}_{\mathcal{R}}(?, iW) \cong \mathrm{Hom}_{\mathcal{R}^\perp}(?, W) \circ i^*$, i.e., the result holds when $n = 0$. Also i^* sends projectives to projectives as it is left adjoint to an exact functor. Now the result for $n > 0$ follows by a standard Grothendieck spectral sequence argument; the spectral sequence degenerates due to (3). \square

Lemma 3.46. *Suppose that $\lambda \in \Lambda$ is maximal and $b \in \mathbf{B}_\lambda$. Then $P(b) \cong \Delta(b)$ and $I(b) \cong \nabla(b)$.*

Proof. Lemma 3.1 shows that $\Delta(b) \cong i_{\leq \lambda}^* P(b)$ and $\nabla(b) \cong i_{\leq \lambda}^! I(b)$.

To complete the proof for $P(b)$, it remains to observe that $P(b)$ belongs to $\mathcal{R}_{\leq \lambda}$, so $i_{\leq \lambda}^* P(b) = P(b)$. This follows from $\widehat{P\Delta_\varepsilon}$: the object P_b belongs to $\mathcal{R}_{\leq \lambda}$ due to the maximality of λ and $P(b)$ is a summand of it.

The proof for $I(b)$ needs a different approach. From $\nabla(b) \cong i_{\leq \lambda}^! I(b)$, we deduce that there is a short exact sequence $0 \rightarrow \nabla(b) \rightarrow I(b) \rightarrow Q \rightarrow 0$ with $i_{\leq \lambda}^! Q = 0$, and we must show that $Q = 0$. Take $a \in \mathbf{B}$ and apply $\mathrm{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), ?)$ to this short exact sequence to get an exact sequence

$$\mathrm{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), I(b)) \longrightarrow \mathrm{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), Q) \longrightarrow 0 \quad (3.12)$$

and isomorphisms

$$\mathrm{Ext}_{\mathcal{R}}^{n+1}(\Delta_\varepsilon(a), \nabla(b)) \cong \mathrm{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(a), Q) \quad (3.13)$$

for $n \geq 1$. If $\rho(a) = \lambda$ then $\mathrm{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), Q) = 0$ because $i_{\leq \lambda}^! Q = 0$. If $\rho(a) \neq \lambda$, then in fact we have that $\rho(a) \not\geq \lambda$ by the assumed maximality of λ , so $[\Delta_\varepsilon(a) : L(b)] = 0$. Hence, $\mathrm{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), I(b)) = 0$, implying in view of (3.12) that $\mathrm{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), Q) = 0$ again. Thus, we have shown that $\mathrm{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), Q) = 0$ for all $a \in \mathbf{B}$. This implies that $\mathrm{soc} Q = 0$. In the essentially finite Abelian case, this is all that is needed to deduce that $Q = 0$, completing the proof. In the Schurian case, we need to argue a little further because Q need not be finitely cogenerated, so can have zero socle even when it is itself non-zero. We have for any $a \in \mathbf{B}$ that $\mathrm{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(a), \nabla(b)) = 0$ for $n > 0$. This follows using Lemma 3.45(4): it shows that $\mathrm{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(a), \nabla(b)) \cong \mathrm{Ext}_{\mathcal{R}_{\leq \lambda}}^n(i_{\leq \lambda}^* \Delta_\varepsilon(a), \nabla(b))$ which is zero as $\nabla(b)$ is injective in $\mathcal{R}_{\leq \lambda}$. Combining this with (3.13), we get that $\mathrm{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(a), Q) = 0$. Now we observe that the properties $\mathrm{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), Q) = 0 = \mathrm{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(a), Q)$ for all $a \in \mathbf{B}$ do imply that Q is zero. Indeed, we have that $\mathrm{Hom}_{\mathcal{R}}(P, Q) = \mathrm{Ext}_{\mathcal{R}}^1(P, Q) = 0$ for any $P \in \mathcal{R}$ with a Δ_ε -flag. This follows using induction on the length of the flag plus the long exact sequence. Since P_b has a Δ_ε -flag by the hypothesis ($\widehat{P\Delta_\varepsilon}$) and $P(b)$ is a summand of it, we deduce that $\mathrm{Hom}_{\mathcal{R}}(P(b), Q) = 0$ for all $b \in \mathbf{B}$, which certainly implies that $Q = 0$. \square

Lemma 3.47. *Assume that $\lambda \in \Lambda$ is maximal and $\varepsilon(\lambda) = +$. For any $V \in \mathcal{R}_\lambda$ and $b \in \mathbf{B}$, we have that $\mathrm{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), j_*^\lambda V) = 0$.*

Proof. If $b \in \mathbf{B}_\lambda$ then $\Delta_\varepsilon(b)$ is projective in $\mathcal{R}_{\leq \lambda}$ by Lemma 3.46, so we get the Ext^1 -vanishing in this case. For the remainder of the proof, suppose that $b \notin \mathbf{B}_\lambda$. Let I be an injective hull of V in \mathcal{R}_λ . Applying j_*^λ to a short exact sequence $0 \rightarrow V \rightarrow I \rightarrow Q \rightarrow 0$, we get an exact sequence $0 \rightarrow j_*^\lambda V \rightarrow j_*^\lambda I \rightarrow j_*^\lambda Q$. By properties of adjunctions, $j_*^\lambda Q$ is finitely cogenerated and all constituents of its socle are of the form $L(c)$ for $c \in \mathbf{B}_\lambda$. The same is true for $j_*^\lambda I / j_*^\lambda V$ since it embeds into $j_*^\lambda Q$. We deduce that $\mathrm{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), j_*^\lambda I / j_*^\lambda V) = 0$.

Now take an extension $0 \rightarrow j_*^\lambda V \rightarrow E \rightarrow \Delta_\varepsilon(b) \rightarrow 0$. Since $j_*^\lambda I$ is injective, we can find morphisms f and g making the following diagram with exact rows commute:

$$\begin{array}{ccccccc} 0 & \longrightarrow & j_*^\lambda V & \xrightarrow{s} & E & \longrightarrow & \Delta_\varepsilon(b) & \longrightarrow & 0 \\ & & \parallel & & f \downarrow & & \downarrow g & & \\ 0 & \longrightarrow & j_*^\lambda V & \xrightarrow{t} & j_*^\lambda I & \longrightarrow & j_*^\lambda I / j_*^\lambda V & \longrightarrow & 0. \end{array}$$

The previous paragraph implies that $g = 0$. Hence, $\text{im } f \subseteq \text{im } t$. Thus, $f = t \circ \bar{f}$ for some $\bar{f}: E \rightarrow j_*^\lambda V$. We deduce that $\bar{f} \circ s = \text{id}$, i.e., the top short exact sequence splits, proving that $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), j_*^\lambda V) = 0$. \square

Lemma 3.48. *For $b, c \in \mathbf{B}$ and $n \geq 0$, we have that $\dim \text{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(b), \nabla_\varepsilon(c)) = \delta_{b,c} \delta_{n,0}$.*

Proof. The case $n = 0$ follows from (3.5), so assume that $n > 0$. Suppose that $b \in \mathbf{B}_\lambda$ and $c \in \mathbf{B}_\mu$. By Lemma 3.45(4), we have that

$$\text{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(b), \nabla_\varepsilon(c)) \cong \text{Ext}_{\mathcal{R}_{\leq \mu}}^n(i_{\leq \mu}^* \Delta_\varepsilon(b), \nabla_\varepsilon(c)).$$

If $\lambda \not\leq \mu$ then $i_{\leq \mu}^* \Delta_\varepsilon(b) = 0$ and we get the desired vanishing. Now assume that $\lambda \leq \mu$, when we may identify $i_{\leq \mu}^* \Delta_\varepsilon(b) = \Delta_\varepsilon(b)$. If $\varepsilon(\mu) = -$ then $\nabla_\varepsilon(c) = \nabla(c)$, and the result follows since $\nabla(c)$ is injective in $\mathcal{R}_{\leq \mu}$ by Lemma 3.1(2). So we may assume also that $\varepsilon(\mu) = +$. If $\lambda = \mu$ then $\Delta(b)$ is projective in $\mathcal{R}_{\leq \mu}$ by the same lemma, so again we are done. Finally, we are reduced to $\lambda < \mu$ and $\varepsilon(\mu) = +$, and need to show that $\text{Ext}_{\mathcal{R}_{\leq \mu}}^n(\Delta_\varepsilon(b), \bar{\nabla}(c)) = 0$ for $n > 0$. If $n = 1$, we get the desired conclusion from Lemma 3.47 applied in the subcategory $\mathcal{R}_{\leq \mu}$ (allowed due to Lemma 3.45(2)). Then for $n \geq 2$ we use a degree shifting argument: let $P := i_{\leq \mu}^* P_b$. By Lemma 3.45(2), P is projective in $\mathcal{R}_{\leq \mu}$, and there is a short exact sequence $0 \rightarrow K \rightarrow P \rightarrow \Delta_\varepsilon(b) \rightarrow 0$ such that K has a Δ_ε -flag with sections $\Delta_\varepsilon(a)$ for $a \in \mathbf{B}_{\leq \mu}$. Applying $\text{Hom}_{\mathcal{R}_{\leq \mu}}(?, \bar{\nabla}(c))$ we obtain $\text{Ext}_{\mathcal{R}_{\leq \mu}}^n(\Delta_\varepsilon(b), \bar{\nabla}(c)) \cong \text{Ext}_{\mathcal{R}_{\leq \mu}}^{n-1}(K, \bar{\nabla}(c))$, which is zero by induction. \square

Lemma 3.49. *Let Λ^\dagger be an upper set in Λ and $\mathbf{B}^\dagger := \rho^{-1}(\Lambda^\dagger)$. Let $j: \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be the corresponding Serre quotient category of \mathcal{R} equipped with the induced stratification.*

- (1) *For $b \in \mathbf{B}^\dagger$, the objects $P^\dagger(b)$, $I^\dagger(b)$, $\Delta^\dagger(b)$, $\bar{\Delta}^\dagger(b)$, $\nabla^\dagger(b)$ and $\bar{\nabla}^\dagger(b)$ in \mathcal{R}^\dagger are the images under j of the corresponding objects of \mathcal{R} . Moreover, we have that $j_! \Delta^\dagger(b) \cong \Delta(b)$, $j_! \bar{\Delta}^\dagger(b) \cong \bar{\Delta}(b)$, $j_! P^\dagger(b) \cong P(b)$ and $j_* \nabla^\dagger(b) \cong \nabla(b)$, $j_* \bar{\nabla}^\dagger(b) \cong \bar{\nabla}(b)$, $j_* I^\dagger(b) \cong I(b)$.*
- (2) *For any $b \in \mathbf{B}$, the object jP_b has a Δ_ε -flag with top section $\Delta_\varepsilon^\dagger(b)$ and other sections of the form $\Delta_\varepsilon^\dagger(c)$ for $c \in \mathbf{B}^\dagger$ with $\rho(c) \geq \rho(b)$. In particular, this shows that $(P\widehat{\Delta}_\varepsilon)$ holds in \mathcal{R}^\dagger .*
- (3) *$\text{Ext}_{\mathcal{R}}^n(V, j_* W) \cong \text{Ext}_{\mathcal{R}^\dagger}^n(jV, W)$ for $V \in \mathcal{R}$, $W \in \nabla_\varepsilon(\mathcal{R}^\dagger)$ and $n \geq 0$.*

Proof. (1) By Lemma 2.27, $P^\dagger(b) = jP(b)$ for each $b \in \mathbf{B}^\dagger$. Now take $b \in \mathbf{B}_\lambda$ for $\lambda \in \Lambda^\dagger$. Let $j^\lambda: \mathcal{R}_{\leq \lambda} \rightarrow \mathcal{R}_\lambda$ be the quotient functor as usual, and denote the analogous functor for \mathcal{R}^\dagger by $k^\lambda: \mathcal{R}_{\leq \lambda}^\dagger \rightarrow \mathcal{R}_\lambda^\dagger$. The universal property of quotient category gives us an exact functor $\bar{j}: \mathcal{R}_\lambda \rightarrow \mathcal{R}_\lambda^\dagger$ making the diagram

$$\begin{array}{ccc} \mathcal{R}_{\leq \lambda} & \xrightarrow{j} & \mathcal{R}_{\leq \lambda}^\dagger \\ j^\lambda \downarrow & & \downarrow k^\lambda \\ \mathcal{R}_\lambda & \xrightarrow{\bar{j}} & \mathcal{R}_\lambda^\dagger \end{array}$$

commute. In fact, \bar{j} is an equivalence of categories because it sends the indecomposable projective $j^\lambda P(b)$ in \mathcal{R}_λ to the indecomposable projective $k^\lambda P^\dagger(b)$ in $\mathcal{R}_\lambda^\dagger$ for each $b \in \mathbf{B}_\lambda$. We deduce that there is an isomorphism of functors $j_! \circ k_!^\lambda \circ \bar{j} \cong j_!^\lambda$. Applying this to $P_\lambda(b)$ and to $L_\lambda(b)$ gives that $j_! \Delta^\dagger(b) \cong \Delta(b)$ and $j_! \bar{\Delta}^\dagger(b) \cong \bar{\Delta}(b)$. Also by adjunction properties we have that $j_! P^\dagger(b) \cong P(b)$. Similarly, applying the isomorphism

$j_* \circ k_*^\lambda \circ \bar{j} \cong j_*^\lambda$ to $I_\lambda(b)$ and $L_\lambda(b)$ gives that $j_* \nabla^\dagger(b) \cong \nabla(b)$ and $j_* \bar{\nabla}^\dagger(b) \cong \bar{\nabla}(b)$. Also by adjunction properties we have that $j_* I^\dagger(b) \cong I(b)$. It just remains to apply j to the isomorphisms constructed thus far and use $j \circ j_* \cong \text{Id}_{\mathcal{R}^\dagger} \cong j \circ j_!$.

(2) This follows from (1) and the exactness of j , using also that $j \Delta_\varepsilon(b) = 0$ if $b \notin \mathbf{B}^\dagger$.

(3) The adjunction gives an isomorphism $\text{Hom}_{\mathcal{R}}(?, j_* W) \cong \text{Hom}_{\mathcal{R}^\dagger}(?, W) \circ j$. This proves the result when $n = 0$. For $n > 0$, the functor j is exact. In order to invoke the usual degenerate Grothendieck spectral sequence argument, all that remains is to check that j sends projectives to objects that are acyclic for $\text{Hom}_{\mathcal{R}^\dagger}(?, W)$. By (2), the functor j sends projectives in \mathcal{R} to objects with a Δ_ε -flag. It remains to note that $\text{Ext}_{\mathcal{R}^\dagger}^1(X, W) = 0$ for $X \in \Delta_\varepsilon(\mathcal{R}^\dagger)$, $W \in \nabla_\varepsilon(\mathcal{R}^\dagger)$. This follows from the analog of Lemma 3.48 for \mathcal{R}^\dagger , which is valid due to (2). \square

3.5. Lower finite ε -stratified categories. In this subsection, \mathcal{R} is a locally finite Abelian category equipped a lower finite stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ and $\varepsilon : \Lambda \rightarrow \{\pm\}$ denotes a sign function. For $b \in \mathbf{B}$, we use the notation $I(b)$ to denote an injective hull of $L(b)$ in $\text{Ind}(\mathcal{R})$.

Definition 3.50. Let $(\mathbf{B}, L, \rho, \Lambda, \leq)$ be a lower finite stratification of the locally finite Abelian category \mathcal{R} . For a finite lower set Λ^\downarrow in Λ , let $\mathbf{B}^\downarrow := \rho^{-1}(\Lambda^\downarrow)$ and \mathcal{R}^\downarrow be corresponding Serre subcategory of \mathcal{R} . We say that \mathcal{R} is a *lower finite ε -stratified category* (resp., *lower finite fully stratified category*, resp., *lower finite ε -highest weight category*, resp., *lower finite fibered highest weight category*, resp., *lower finite highest weight category*) if \mathcal{R}^\downarrow with its naturally induced stratification is a finite ε -stratified category (resp., finite fully stratified category, resp., finite ε -highest weight category, resp., finite fibered highest weight category, resp., finite highest weight category) for every finite lower set $\Lambda^\downarrow \subseteq \Lambda$.

Remark 3.51. For a simple example, let Q be any quiver. The category \mathcal{R} of finite length nilpotent representations of Q can be realized naturally as the category of finite-dimensional comodules over the path coalgebra of Q as in [Sim, (8.3)]. In order for this to be a lower finite highest weight category, one must assume that there are only finitely many different paths between any two vertices. In that case, the path algebra $\mathbb{k}Q$ is locally finite-dimensional, and we have that $\mathcal{R} \cong \mathbb{k}Q\text{-mod}_{\text{fd}}$ with irreducible objects labelled by the set Λ of vertices of Q in the usual way. We claim now that \mathcal{R} is a lower finite highest weight category with weight poset (Λ, \leq) for any lower finite partial ordering \leq on Λ . To see this, the Serre subcategory \mathcal{R}^\downarrow corresponding to a finite lower set $\Lambda^\downarrow \subset \Lambda$ is $\mathbb{k}Q^\downarrow\text{-mod}_{\text{fd}}$ where Q^\downarrow is the full subquiver Q^\downarrow of Q generated by Λ^\downarrow . It is well known that this is a hereditary category, hence, it is a finite highest weight category (e.g., see [Mad, Th. 4.1]).

Let \mathcal{R} be a lower finite ε -stratified category. Since $\mathcal{R}_{\leq \lambda}$ is a finite Abelian category, the admissibility axiom (A) from §3.1 holds, so we can introduce the objects $\Delta(b)$, $\bar{\Delta}(b)$, $\bar{\nabla}(b)$ and $\nabla(b)$ as explained there, also adopting the shorthands $\Delta_\varepsilon(b)$ and $\nabla_\varepsilon(b)$. These objects are of finite length. Note also that Theorem 3.9, Lemma 3.20 and Corollary 3.21 carry over immediately to the lower finite setting.

Now we are going to consider another sort of infinite good filtration in objects of $\text{Ind}(\mathcal{R})$. Usually (e.g., if Λ is countable), it is sufficient to restrict attention to filtrations given by an ascending chain of subobjects $0 = V_0 < V_1 < V_2 < \dots$ such that $V = \sum_{n \in \mathbb{N}} V_n$ and $V_m/V_{m-1} \cong \nabla_\varepsilon(b_m)$ for some $b_m \in \mathbf{B}$. Here is the general definition which avoids this restriction.

Definition 3.52. An *ascending ∇_ε -flag* in an object $V \in \text{Ind}(\mathcal{R})$ is the data of a direct system $(V_\omega)_{\omega \in \Omega}$ of subobjects of V such that the following properties hold:

$$(A\nabla 1) \quad V = \sum_{\omega \in \Omega} V_\omega;$$

$$(A\nabla 2) \quad \text{each } V_\omega \text{ has a } \nabla_\varepsilon\text{-flag with } \nabla_\varepsilon(b) \text{ appearing with multiplicity } (V_\omega : \nabla_\varepsilon(b)) \in \mathbb{N};$$

(A ∇ 3) $(V : \nabla_\varepsilon(b)) := \sup\{(V_\omega, \nabla_\varepsilon(b)) \mid \omega \in \Omega\} < \infty$ for each $b \in \mathbf{B}$.

Let $\nabla_\varepsilon^{\text{asc}}(\mathcal{R})$ be the full subcategory of \mathcal{R} consisting of all objects V that possess an ascending ∇_ε -flag. In the special case $\varepsilon = +$ (resp., $\varepsilon = -$), we call it an ascending $\bar{\nabla}$ -flag (resp., ∇ -flag), denoting the category $\nabla_\varepsilon(\mathcal{R})$ by $\bar{\nabla}(\mathcal{R})$ (resp., $\nabla(\mathcal{R})$).

The multiplicities $(V_\omega : \nabla_\varepsilon(b))$ and $(V : \nabla_\varepsilon(b))$ appearing in this definition depend *a priori* on the choice of flag. In fact, they do not, so that the notation is unambiguous:

Lemma 3.53. *Assume \mathcal{R} is a lower finite ε -stratified category. For $V \in \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$, the multiplicity $(V : \nabla_\varepsilon(b))$ of $\nabla_\varepsilon(b)$ in the ascending ∇_ε -flag appearing in Definition 3.52 is equal to $\dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V)$. Hence, it is well-defined independent of the particular choice for this flag.*

Proof. By Theorem 3.14 applied in the Serre subcategory \mathcal{R}^\downarrow associated to a finite lower set Λ^\downarrow of Λ chosen so that $V_\omega \in \mathcal{R}^\downarrow$, we have that $(V_\omega : \nabla_\varepsilon(b)) = \dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V_\omega)$. Also $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) = \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), \varinjlim V_\omega) \cong \varinjlim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V_\omega)$. We deduce that

$$\dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) = \max\{(V_\omega : \Delta_\varepsilon(b)) \mid \omega \in \Omega\},$$

which is the definition of the multiplicity $(V : \nabla_\varepsilon(b))$ from Definition 3.52. \square

Lemma 3.54. *Assume that \mathcal{R} is a lower finite ε -stratified category. For $V \in \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$ and $b \in \mathbf{B}$, we have that $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), V) = 0$.*

Proof. If V is of finite length then it belongs to the finite Abelian category \mathcal{R}^\downarrow associated to some finite lower set Λ^\downarrow of Λ , and the lemma follows from Theorem 3.11. Now suppose that V is not of finite length. Let $(V_\omega)_{\omega \in \Omega}$ be an ascending ∇_ε -flag in V . Take an extension $V \hookrightarrow E \twoheadrightarrow \Delta_\varepsilon(b)$. We can find a subobject E_1 of E of finite length such that $V + E_1 = V + E$; this follows easily by induction on the length of $\Delta_\varepsilon(b)$ as explained at the start of the proof of [CPS1, Lem. 3.8(a)]. Since $V \cap E_1$ is of finite length, there exists $\omega \in \Omega$ with $V \cap E_1 \subseteq V_\omega$. Then we have that $V \cap E_1 = V_\omega \cap E_1$ and

$$(V_\omega + E_1)/V_\omega \cong E_1/V_\omega \cap E_1 = E_1/V \cap E_1 \cong (V + E_1)/V = (V + E)/V \cong \Delta_\varepsilon(b).$$

Thus, there is a short exact sequence $0 \rightarrow V_\omega \rightarrow V_\omega + E_1 \rightarrow \Delta_\varepsilon(b) \rightarrow 0$. The first sentence of the proof implies that $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), V_\omega) = 0$, hence, this splits. Thus, we can find a subobject $E_2 \cong \Delta_\varepsilon(b)$ of $V_\omega + E_1$ such that $V_\omega + E_1 = V_\omega \oplus E_2$. Then $V + E = V + E_1 = V + V_\omega + E_1 = V + V_\omega + E_2 = V + E_2 = V \oplus E_2$, and our original short exact sequence splits too. \square

Corollary 3.55. *Let $i : \mathcal{R}^\downarrow \rightarrow \mathcal{R}$ be the inclusion of the Serre subcategory of \mathcal{R} associated to a finite lower set Λ^\downarrow of Λ and i^\downarrow be its right adjoint. For $V \in \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$, we have that $i^\downarrow V \in \nabla_\varepsilon(\mathcal{R}^\downarrow)$.*

Proof. Take a short exact sequence $0 \rightarrow i^\downarrow V \rightarrow V \rightarrow Q \rightarrow 0$. Note that

$$\text{Hom}_{\mathcal{R}^\downarrow}(\Delta_\varepsilon(b), i^\downarrow V) \cong \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V)$$

is finite-dimensional for each $b \in \mathbf{B}^\downarrow$. Since \mathcal{R}^\downarrow is finite Abelian, it follows that $i^\downarrow V \in \mathcal{R}^\downarrow$ (rather than $\text{Ind}(\mathcal{R}^\downarrow)$). Moreover, $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), Q) = 0$ for $b \in \mathbf{B}^\downarrow$. So, on applying $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), ?)$ and considering the long exact sequence using Lemma 3.54, we get that $\text{Ext}_{\mathcal{R}^\downarrow}^1(\Delta_\varepsilon(b), i^\downarrow V) = \text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), i^\downarrow V) = 0$ for all $b \in \mathbf{B}^\downarrow$. Thus, by Theorem 3.11, we have that $i^\downarrow V \in \nabla_\varepsilon(\mathcal{R}^\downarrow)$. \square

The following homological criterion for ascending ∇_ε -flags generalizes Theorem 3.11.

Theorem 3.56 (Homological criterion for ascending ∇_ε -flags). *Assume that \mathcal{R} is a lower finite ε -stratified category. For $V \in \text{Ind}(\mathcal{R})$, the following are equivalent:*

- (i) $V \in \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$;
- (ii) $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(b), V) = 0$ and $\dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) < \infty$ for all $b \in \mathbf{B}$;
- (iii) $\text{Ext}_{\mathcal{R}}^n(\Delta_\varepsilon(b), V) = 0$ and $\dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) < \infty$ for all $b \in \mathbf{B}$ and $n \geq 1$.

Assuming these properties, we have that $V \in \nabla_\varepsilon(\mathcal{R})$ if and only if $V \in \mathcal{R}$.

Proof. (ii) \Rightarrow (i): Let Ω be the directed set consisting of all finite lower sets in Λ . Take $\omega \in \Omega$. It is a finite lower set $\Lambda^\downarrow \subseteq \Lambda$, so we have associated the corresponding finite ε -stratified subcategory \mathcal{R}^\downarrow . Letting $i : \mathcal{R}^\downarrow \rightarrow \mathcal{R}$ be the inclusion, we set $V_\omega := i^\downarrow V$. By Corollary 3.55, we have that $V_\omega \in \nabla_\varepsilon(\mathcal{R})$. So the subobject $V' := \sum_{\omega \in \Omega} V_\omega$ of V has an ascending ∇_ε -flag.

Now we complete the proof by showing that $V = V'$. Applying $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), ?)$ to the short exact sequence $0 \rightarrow V' \rightarrow V \rightarrow V/V' \rightarrow 0$ using Lemma 3.54, we get a short exact sequence

$$0 \longrightarrow \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V') \longrightarrow \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V) \longrightarrow \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V/V') \longrightarrow 0$$

for every $b \in \mathbf{B}$. But any homomorphism $\Delta_\varepsilon(b) \rightarrow V$ has image contained in V_ω for sufficiently large ω , hence, also in V' . Thus, the first morphism in this short exact sequence is an isomorphism, and $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), V/V') = 0$ for all $b \in \mathbf{B}$. This implies that $V/V' = 0$ as required.

(i) \Rightarrow (ii): This follows by Lemmas 3.53 and 3.54.

(iii) \Rightarrow (ii): Trivial.

(i) \Rightarrow (iii): This follows from Lemma 3.53 and Theorem 3.59(4). Since this is a forward reference, we should note that the proof of Theorem 3.59 only depends on (i) \Leftrightarrow (ii) from the present theorem. \square

Corollary 3.57. *In a lower finite ε -stratified category, each indecomposable injective object $I(b)$ belongs to $\nabla_\varepsilon^{\text{asc}}(\mathcal{R})$ and $(I(b) : \nabla_\varepsilon(c)) = [\Delta_\varepsilon(c) : L(b)]$ for each $b, c \in \mathbf{B}$.*

Proof. The first part follows from the implication (ii) \Rightarrow (i) in the theorem. For the second part, we get from Lemma 3.53 that $(I(b) : \nabla_\varepsilon(c)) = \dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(c) : L(b))$. \square

Corollary 3.58. *Let $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ be a short exact sequence in a lower finite ε -stratified category. If $U, V \in \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$ then $W \in \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$ too. Moreover*

$$(V : \nabla_\varepsilon(b)) = (U : \nabla_\varepsilon(b)) + (W : \nabla_\varepsilon(b)).$$

The following is the lower finite counterpart of Theorem 3.17.

Theorem 3.59 (Truncation to lower sets). *Suppose \mathcal{R} is a lower finite ε -stratified category. Let Λ^\downarrow be a lower set, $\mathbf{B}^\downarrow := \rho^{-1}(\Lambda^\downarrow)$, and $i : \mathcal{R}^\downarrow \rightarrow \mathcal{R}$ be the corresponding Serre subcategory of \mathcal{R} with the induced stratification. Then \mathcal{R}^\downarrow is a finite or lower finite ε -stratified category according to whether Λ^\downarrow is finite or infinite. Moreover:*

- (1) *The distinguished objects of \mathcal{R}^\downarrow are $L^\downarrow(b) \cong L(b)$, $I^\downarrow(b) \cong i^\downarrow I(b)$, $\Delta^\downarrow(b) \cong \Delta(b)$, $\bar{\Delta}^\downarrow(b) \cong \bar{\Delta}(b)$, $\nabla^\downarrow(b) \cong \nabla(b)$ and $\bar{\nabla}^\downarrow(b) \cong \bar{\nabla}(b)$ for $b \in \mathbf{B}^\downarrow$.*
- (2) *$(\mathbb{R}^n i^\downarrow)V = 0$ for $n \geq 1$ assuming either that $V \in \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$ or that $V \in \mathcal{R}^\downarrow$.*
- (3) *i^\downarrow takes short exact sequences of objects in $\nabla_\varepsilon^{\text{asc}}(\mathcal{R})$ to short exact sequences of objects in $\nabla_\varepsilon^{\text{asc}}(\mathcal{R}^\downarrow)$, with $i^\downarrow \nabla(b) \cong \nabla^\downarrow(b)$ and $i^\downarrow \bar{\nabla}(b) \cong \bar{\nabla}^\downarrow(b)$ for $b \in \mathbf{B}^\downarrow$ and $i^\downarrow \nabla(b) = i^\downarrow \bar{\nabla}(b) = 0$ for $b \notin \mathbf{B}^\downarrow$.*
- (4) *$\text{Ext}_{\mathcal{R}}^n(iV, W) \cong \text{Ext}_{\mathcal{R}^\downarrow}^n(V, i^\downarrow W)$ for $V \in \mathcal{R}^\downarrow, W \in \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$ and all $n \geq 0$.*
- (5) *$\text{Ext}_{\mathcal{R}}^n(iV, iW) \cong \text{Ext}_{\mathcal{R}^\downarrow}^n(V, W)$ for $V, W \in \mathcal{R}^\downarrow$ and all $n \geq 0$.*

Proof. The fact that \mathcal{R}^\downarrow is itself a lower finite ε -stratified follows immediately from Definition 3.50. It is finite if and only if \mathbf{B}^\downarrow is finite. The identification of objects as in (1) is straightforward. In particular, the objects $\nabla_\varepsilon(b)$ in \mathcal{R}^\downarrow are just the same as the ones in \mathcal{R} indexed by $b \in \mathbf{B}^\downarrow$, while the indecomposable injectives in $\text{Ind}(\mathcal{R}^\downarrow)$ are the objects $i^\downarrow I(b)$ for $b \in \mathbf{B}^\downarrow$.

To prove (2), assume first that $V \in \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$. Let I be an injective hull of $\text{soc } V$ in $\text{Ind}(\mathcal{R})$. Note that I is of the form $\bigoplus_{a \in \mathbf{B}} I(a)^{\oplus n_a}$ for

$$0 \leq n_a \leq \dim \text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), V) = (V : \nabla_\varepsilon(a)) < \infty.$$

Hence, for $b \in \mathbf{B}_{\leq \lambda}$, we have that

$$\dim \operatorname{Hom}_{\mathcal{R}}(\Delta_{\varepsilon}(b), I) = \sum_{a \in \mathbf{B}_{\leq \lambda}} n_a[\Delta_{\varepsilon}(b) : L(a)] < \infty$$

too. We deduce that $I \in \nabla_{\varepsilon}^{\operatorname{asc}}(\mathcal{R})$ using the implication (ii) \Rightarrow (i) of Theorem 3.56. Now consider the short exact sequence $0 \rightarrow V \rightarrow I \rightarrow Q \rightarrow 0$. By Corollary 3.58, we have that $Q \in \nabla_{\varepsilon}^{\operatorname{asc}}(\mathcal{R})$ too. Applying the left exact functor $i^!$ and considering the long exact sequence, we see that to prove that $(\mathbb{R}^1 i^!)V = 0$ it suffices to show that the canonical map $i^!I \rightarrow i^!Q$ is an epimorphism. Once that has been proved we can use degree shifting to establish the desired vanishing for all higher n ; it is important for the induction step that we have already established that $Q \in \nabla_{\varepsilon}^{\operatorname{asc}}(\mathcal{R})$ just like V .

To prove the surjectivity, look at $0 \rightarrow i^!I/i^!V \rightarrow i^!Q \rightarrow C \rightarrow 0$. Both $i^!I$ and $i^!V$ have ∇_{ε} -flags by Lemma 3.55. Hence, so does $i^!I/i^!V$, and on applying $\operatorname{Hom}_{\mathcal{R}^{\downarrow}}(\Delta_{\varepsilon}(b), ?)$ for $b \in \mathbf{B}^{\downarrow}$, we get a short exact sequence

$$0 \longrightarrow \operatorname{Hom}_{\mathcal{R}^{\downarrow}}(\Delta_{\varepsilon}(b), i^!I/i^!V) \longrightarrow \operatorname{Hom}_{\mathcal{R}^{\downarrow}}(\Delta_{\varepsilon}(b), i^!Q) \longrightarrow \operatorname{Hom}_{\mathcal{R}^{\downarrow}}(\Delta_{\varepsilon}(b), C) \longrightarrow 0.$$

The first space here has dimension

$$(i^!I : \nabla_{\varepsilon}(b)) - (i^!V : \nabla_{\varepsilon}(b)) = (I : \nabla_{\varepsilon}(b)) - (V : \nabla_{\varepsilon}(b)) = (Q : \nabla_{\varepsilon}(b)) = (i^!Q : \nabla_{\varepsilon}(b)),$$

which is the dimension of the second space. This shows that the first morphism is an isomorphism. Hence, $\operatorname{Hom}_{\mathcal{R}^{\downarrow}}(\Delta_{\varepsilon}(b), C) = 0$. This implies that $C = 0$ as required.

Finally let $V \in \mathcal{R}^{\downarrow}$. Then V is of finite length, so it suffices just to consider the case that $V = L(b)$ for $b \in \mathbf{B}^{\downarrow}$. Then we consider the short exact sequence $0 \rightarrow L(b) \rightarrow \nabla_{\varepsilon}(b) \rightarrow Q \rightarrow 0$. Applying $i^!$ and using the vanishing established so far gives $0 \rightarrow i^!L(b) \rightarrow i^!\nabla_{\varepsilon}(b) \rightarrow i^!Q \rightarrow (\mathbb{R}^1 i^!)L(b) \rightarrow 0$ and isomorphisms $(\mathbb{R}^n i^!)Q \cong (\mathbb{R}^{n+1} i^!)L(b)$ for $n \geq 1$. But $i^!$ is the identity on $L(b), \nabla_{\varepsilon}(b)$ and Q , so this immediately yields $(\mathbb{R}^1 i^!)L(b) = 0$, and then $(\mathbb{R}^n i^!)L(b) = 0$ for higher n by degree shifting.

Having proved (2), property (3) follows easily. Finally (4)–(5) follow by the usual Grothendieck spectral sequence argument starting from the adjunction isomorphism $\operatorname{Hom}_{\mathcal{R}^{\downarrow}}(iV, ?) \cong \operatorname{Hom}_{\mathcal{R}}(V, ?) \circ i^!$. One just needs (2) and the observation that $i^!$ sends injectives to injectives. \square

Our next result gives an alternative characterization of lower finite ε -stratified categories. Note for this that if \mathcal{R} is a lower finite ε -stratified category then the hypotheses of the theorem are automatically satisfied taking $I_b^{\operatorname{asc}} := I(b)$; cf. Corollary 3.57.

Theorem 3.60 (Global characterization of lower finite ε -stratified categories). *Let \mathcal{R} be a locally finite Abelian category equipped with a lower finite stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ and $\varepsilon : \Lambda \rightarrow \{\pm\}$ be a sign function. Assume for each $b \in \mathbf{B}$ that $L(b)$ has an injective hull in $\mathcal{R}_{\leq \rho(b)}$ so that we can introduce the objects $\nabla_{\varepsilon}(b)$ in the usual way³. Suppose that the following property holds:*

($\widehat{I\nabla}_{\varepsilon}^{\operatorname{asc}}$) For every $b \in \mathbf{B}$, there exists an injective object $I_b \in \operatorname{Ind}(\mathcal{R})$ with an ascending ∇_{ε} -flag $(V_{\omega})_{\omega \in \Omega}$ in the sense of Definition 3.52 such that for each $\omega \in \Omega$ the given ∇_{ε} -flag of V_{ω} has $\nabla_{\varepsilon}(b)$ at the bottom and all other sections are of the form $\nabla_{\varepsilon}(c)$ for $c \in \mathbf{B}$ with $\rho(c) \geq \rho(b)$.

Then \mathcal{R} is a lower finite ε -stratified category.

Proof. We must verify the condition from Definition 3.50. Let Λ^{\downarrow} be a finite lower set, $\mathbf{B}^{\downarrow} := \rho^{-1}(\Lambda^{\downarrow})$, and \mathcal{R}^{\downarrow} be the corresponding Serre subcategory of \mathcal{R} . This is a locally finite Abelian category with irreducible objects labelled by the finite set \mathbf{B}^{\downarrow} . We need to show it is a finite ε -stratified category with respect to the induced stratification.

Step 1: $\operatorname{Ext}_{\mathcal{R}}^1(\nabla_{\varepsilon}(a), \nabla_{\varepsilon}(b)) = 0$ for $\rho(a) \not\geq \rho(b)$. Let $(V_{\omega})_{\omega \in \Omega}$ be the given ascending ∇_{ε} -flag of I_b . We have that $\nabla_{\varepsilon}(b) \hookrightarrow I_b$ and $I_b/\nabla_{\varepsilon}(b) = \sum_{\omega \in \Omega} (V_{\omega}/\nabla_{\varepsilon}(b))$. The socle of

³We do not insist that $L(b)$ has a projective cover in $\mathcal{R}_{\leq \rho(b)}$ and do not need the objects $\Delta_{\varepsilon}(b)$.

the latter object only involves constituents $L(c)$ with $\rho(c) \geq \rho(b)$. We deduce that there is an injective resolution $0 \rightarrow \nabla_\varepsilon(b) \rightarrow I_b \rightarrow J \rightarrow \cdots$ in $\text{Ind}(\mathcal{R})$ in which J is a direct sum of I_c with $\rho(c) \geq \rho(b)$. The Ext^1 -vanishing now follows on applying $\text{Hom}_{\mathcal{R}}(\nabla_\varepsilon(a), ?)$ to this resolution and taking homology.

Step 2: For $b \in \mathbf{B}^\downarrow$, the object $I_b^\downarrow := i^! I_b \in \text{Ind}(\mathcal{R}^\downarrow)$ has a ∇_ε -flag with $\nabla_\varepsilon(b)$ at the bottom and other sections of the form $\nabla_\varepsilon(c)$ for $c \in \mathbf{B}^\downarrow$ with $\rho(c) \geq \rho(b)$. In particular, I_b^\downarrow is of finite length. Take $b \in \mathbf{B}^\downarrow$ and let $(V_\omega)_{\omega \in \Omega}$ be the given ascending ∇_ε -flag in I_b . Since \mathbf{B}^\downarrow is finite, we can choose some sufficiently large $\omega \in \Omega$ so that $(V : \nabla_\varepsilon(c)) = (V_\omega : \nabla_\varepsilon(c))$ for all $c \in \mathbf{B}^\downarrow$; these multiplicities are the given ones from Definition 3.52. Then we see that $i^! V_v = i^! V_\omega$ for all larger v , hence, $i^! V = i^! V_\omega$. In view of Step 1, we can rearrange the ∇_ε -flag of V_ω so that the sections $\nabla_\varepsilon(c)$ with $c \in \mathbf{B}^\downarrow$ appear below the other sections, with bottom section $\nabla_\varepsilon(b)$. So there is a short exact sequence $0 \rightarrow U_\omega \rightarrow V_\omega \rightarrow W_\omega \rightarrow 0$ such that $U_\omega \in \nabla_\varepsilon(\mathcal{R}^\downarrow)$ and $i^! W_\omega = 0$. Then we get that $i^! V = i^! V_\omega = U_\omega$, which has the desired ∇_ε -flag.

Step 3: \mathcal{R}^\downarrow is a finite ε -stratified category with respect to the induced stratification. By adjunction properties, the object $I_b^\downarrow \in \mathcal{R}^\downarrow$ from Step 2 is injective and it has $L(b)$ in its socle. This shows that the locally finite Abelian category \mathcal{R}^\downarrow has enough injectives, hence, it is a finite Abelian category by Lemma 2.21. Moreover, the objects I_b^\downarrow ($b \in \mathbf{B}^\downarrow$) satisfy the condition $(\widehat{I\nabla}_\varepsilon)$ from §3.2, so \mathcal{R}^\downarrow is a finite ε -stratified category according to Definition 3.7. \square

Corollary 3.61. *Let \mathcal{R} be a locally finite Abelian category, (Λ, \leq) be a lower finite poset, and $L : \Lambda \rightarrow \mathcal{R}$ be a function labelling a complete set of pairwise inequivalent irreducible objects. Assume for all $\lambda \in \Lambda$ that $L(\lambda)$ has an injective hull $\nabla(\lambda) \in \mathcal{R}_{\leq \lambda}$ such that $[\nabla(\lambda) : L(\lambda)] = 1$. Suppose that the following property holds:*

$(\widehat{I\nabla}^{\text{asc}})$ For every $\lambda \in \Lambda$ there exists an injective object $I_\lambda \in \text{Ind}(\mathcal{R})$ with an ascending ∇ -flag $(V_\omega)_{\omega \in \Omega}$ such that for each $\omega \in \Omega$ the given ∇ -flag of V_ω has $\nabla(\lambda)$ at the bottom and all other sections are of the form $\nabla(\mu)$ for $\mu \in \Lambda$ with $\mu \geq \lambda$

Then \mathcal{R} is a lower finite highest weight category.

Proof. Apply the theorem taking $\mathbf{B} = \Lambda$ and ρ to be the identity function, using also Lemma 3.4. \square

Remark 3.62. Using Corollary 3.61, it follows that \mathcal{R} is a lower finite highest weight category with all intervals $(\lambda, \infty]$ in the weight poset being countable if and only if $\text{Ind}(\mathcal{R})$ is a highest weight category in the original sense of [CPS1, Def. 3.1] with a weight poset that is lower finite. This is also mentioned in [Cou3].

The following theorem gives a related characterization for lower finite fully stratified categories. The proof is based on the well-known proof of the homological criterion for good filtrations in the context of reductive algebra groups from [Jan1, Prop. II.4.16]. The Ext^2 -vanishing property needed for this is used as one of the defining properties in [RW, Def. 2.1]; see also [Cou3, Def. 3.1.2(†)]. We know already that lower finite fully stratified categories automatically satisfy the conditions of this theorem.

Theorem 3.63 (Homological characterization of lower finite fully stratified categories). *Suppose that \mathcal{R} is a locally finite Abelian category equipped with a lower finite stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Suppose that every $L(b)$ has a projective cover and an injective hull in $\mathcal{R}_{\leq \rho(b)}$ so that we can introduce standard and costandard objects. Consider the following properties:*

- (1) $\text{Ext}_{\mathcal{R}}^1(\bar{\Delta}(b), \nabla(c)) = \text{Ext}_{\mathcal{R}}^2(\bar{\Delta}(b), \nabla(c)) = 0$ for all $b, c \in \mathbf{B}$.
- (2) $\text{Ext}_{\mathcal{R}}^1(\Delta(b), \bar{\nabla}(c)) = \text{Ext}_{\mathcal{R}}^2(\Delta(b), \bar{\nabla}(c)) = 0$ for all $b, c \in \mathbf{B}$.

If (1) holds then \mathcal{R} is a lower finite $-$ -stratified category, and if (2) holds then \mathcal{R} is a lower finite $+$ -stratified category. Hence, if both (1) and (2) hold then \mathcal{R} is a lower finite fully stratified category.

Proof. We will prove that (1) implies that \mathcal{R} is a lower finite $-$ -stratified category. The fact that (2) implies that \mathcal{R} is $+$ -stratified then follows from this assertion with \mathcal{R} replaced by \mathcal{R}^{op} . Hence, if both hold then \mathcal{R} is fully stratified thanks to Lemma 3.20(iii).

So now we just assume (1). Define ascending ∇ -flags and the corresponding full subcategory $\nabla^{\text{asc}}(\mathcal{R})$ by repeating the $\varepsilon = -$ case of Definition 3.52. We first establish two claims.

Claim 1: For $V \in \nabla^{\text{asc}}(\mathcal{R})$, we have that $\text{Ext}_{\mathcal{R}}^1(\bar{\Delta}(b), V) = 0$ for all $b \in \mathbf{B}$. Moreover, the multiplicity $(V : \nabla(b))$ defined from a specific choice of ascending ∇ -flag in V is equal to $\dim \text{Hom}_{\mathcal{R}}(\bar{\Delta}(b), V)$. For any $c \in \mathbf{B}$, we have as always that $\dim \text{Hom}_{\mathcal{R}}(\bar{\Delta}(b), \nabla(c)) = \delta_{b,c}$, and moreover $\text{Ext}_{\mathcal{R}}^1(\bar{\Delta}(b), \nabla(c)) = 0$ by property (1). Hence, Claim 1 holds when the ∇ -flag is of finite length. Then it follows for arbitrary $V \in \nabla^{\text{asc}}(\mathcal{R})$ by the same arguments as used to prove Lemmas 3.53 and 3.54 above, using the special case just established in place of the references to Theorems 3.11 and 3.14 made in those proofs.

Claim 2: If $V \in \text{Ind}(\mathcal{R})$ satisfies $\dim \text{Hom}_{\mathcal{R}}(\bar{\Delta}(b), V) < \infty$ and $\text{Ext}_{\mathcal{R}}^1(\bar{\Delta}(b), V) = 0$ for all $b \in \mathbf{B}$ then V has an ascending ∇ -flag $(V_{\omega})_{\omega \in \Omega}$. Let Ω be the poset of finite lower sets in Λ ordered by containment. For $\omega = \Lambda^{\downarrow} \in \Omega$, define V_{ω} to be the subobject $i^{\downarrow}V$ where $i : \mathcal{R}^{\downarrow} \rightarrow \mathcal{R}$ is the inclusion of the Serre subcategory of \mathcal{R} associated to $\mathbf{B}^{\downarrow} := \rho^{-1}(\Lambda^{\downarrow})$. This defines a direct system $(V_{\omega})_{\omega \in \Omega}$ of subobjects of V . We prove the claim by establishing the following:

- (a) Each V_{ω} ($\omega \in \Omega$) has a finite ∇ -flag.
- (b) $V = \sum_{\omega \in \Omega} V_{\omega}$.

To check (a), take $\omega = \Lambda^{\downarrow} \in \Omega$ setting $\mathbf{B}^{\downarrow} := \rho^{-1}(\Lambda^{\downarrow})$ once again. We show that V_{ω} has a finite ∇ -flag by induction on $n(V) := \sum_{b \in \mathbf{B}^{\downarrow}} \dim \text{Hom}_{\mathcal{R}}(\bar{\Delta}(b), V)$. If $n(V) = 0$ then $V_{\omega} = 0$ and there is nothing to do. If $n(V) > 0$, let λ be minimal such that $\dim \text{Hom}_{\mathcal{R}}(\bar{\Delta}(b), V) \neq 0$ for some $b \in \mathbf{B}_{\lambda}$. Then $\text{Hom}_{\mathcal{R}}(L(c), V) = 0$ for $c \in \mathbf{B}_{<\lambda}$ and $\text{Hom}_{\mathcal{R}}(L(b), V) \neq 0$. By applying $\text{Hom}_{\mathcal{R}}(?, V)$ to the short exact sequence $0 \rightarrow K \rightarrow \bar{\Delta}(c) \rightarrow L(c) \rightarrow 0$, it follows that $\text{Ext}_{\mathcal{R}}^1(L(c), V) = 0$ for all $c \in \mathbf{B}_{\leq \lambda}$. Then by applying $\text{Hom}_{\mathcal{R}}(?, V)$ to the short exact sequence $0 \rightarrow L(b) \rightarrow \nabla(b) \rightarrow Q \rightarrow 0$, it follows that the natural map $\text{Hom}_{\mathcal{R}}(\nabla(b), V) \rightarrow \text{Hom}_{\mathcal{R}}(L(b), V)$ is surjective. Since the right hand space is non-zero and $\text{soc } \nabla(b) = L(b)$, it follows that there is an injective homomorphism $f : \nabla(b) \rightarrow V$. Let $U := \text{im } f$ and $W := V/U$. Thus, $U \cong \nabla(b)$ and there is a short exact sequence $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$. Applying $\text{Hom}_{\mathcal{R}}(\bar{\Delta}(a), ?)$ and using the hypotheses $\text{Ext}_{\mathcal{R}}^1(\bar{\Delta}(a), U) = \text{Ext}_{\mathcal{R}}^1(\bar{\Delta}(a), V) = \text{Ext}_{\mathcal{R}}^2(\bar{\Delta}(a), U) = 0$, we deduce that $n(W) < n(V)$ and $\text{Ext}_{\mathcal{R}}^1(\bar{\Delta}(a), W) = 0$ for all $a \in \mathbf{B}$. Thus we can apply induction to prove that W_{ω} has a finite ∇ -flag. Since $V_{\omega} = W_{\omega}/U$ it follows that V_{ω} does too, and (a) is proved. To check (b), we let $V' := \sum_{\omega \in \Omega} V_{\omega}$ and show that $V = V'$ by repeating the argument from the proof of (ii) \Rightarrow (i) in Theorem 3.56 with $\Delta_{\varepsilon}(b)$ replaced by $\bar{\Delta}(b)$, using Claim 1 to get that $\text{Ext}_{\mathcal{R}}^1(\bar{\Delta}(b), V') = 0$. Thus, we have proved Claim 2.

Now we complete the proof of the theorem. For $b \in \mathbf{B}$, let $I_b := I(b)$. Like in the proof of Corollary 3.57, Claims 1 and 2 imply that I_b has an ascending ∇ -flag $(V_{\omega})_{\omega \in \Omega}$ with $(I_b : \nabla(c)) = [\bar{\Delta}(c) : L(b)]$. By passing to a subset of Ω if necessary, we may assume that all V_{ω} are non-zero. It follows that the condition $(\widehat{I\nabla}_{-}^{\text{asc}})$ from Theorem 3.60 is satisfied, and \mathcal{R} is a lower finite $-$ -stratified category. \square

Corollary 3.64. *Suppose that \mathcal{R} is a locally finite Abelian category, (Λ, \leq) is a lower finite poset, and $L : \Lambda \rightarrow \mathcal{R}$ is a function labelling a complete set of pairwise inequivalent irreducible objects. Assume $L(\lambda)$ has both an injective hull $\nabla(\lambda)$ and a projective cover $\Delta(\lambda)$ in $\mathcal{R}_{\leq \lambda}$. Suppose that the following properties hold for all $\lambda, \mu \in \Lambda$:*

- (i) $\mathrm{Hom}_{\mathcal{R}}(\Delta(\lambda), \nabla(\lambda))$ is one-dimensional;
- (ii) $\mathrm{Ext}_{\mathcal{R}}^1(\Delta(\lambda), \nabla(\mu)) = \mathrm{Ext}_{\mathcal{R}}^2(\Delta(\lambda), \nabla(\mu)) = 0$.

Then \mathcal{R} is a lower finite highest weight category.

Proof. Property (i) implies that all strata are simple; cf. Lemma 3.4. Now apply the theorem. \square

Corollary 3.64 applies in particular to the category $\mathcal{R} = \mathrm{Rep}(G)$ for a reductive algebraic group G ; see §6.4. The Ext-vanishing properties in the corollary are consequences of Kempf's vanishing theorem; see [Jan1, Prop. II.4.13].

3.6. Refining stratifications in fully stratified categories. We end the section by formulating a basic lemma about refinement of stratifications in fully stratified categories in any of the settings (finite, essentially finite, upper finite or lower finite).

Definition 3.65. Let $(\mathbf{B}, L, \rho, \Lambda, \leq)$ be a stratification of an Abelian category \mathcal{R} . A *refinement* of it means a stratification $(\mathbf{B}, L, \sigma, \Gamma, \leq)$ of \mathcal{R} with the same underlying labelling function together with a surjective function $q : \Gamma \rightarrow \Lambda$ such that the following properties hold:

- (R1) $\Gamma \cap \Lambda = \emptyset$.
- (R2) $\rho = q \circ \sigma$.
- (R3) For $\beta, \gamma \in \Gamma$, we have that $\beta \leq \gamma \Rightarrow q(\beta) \leq q(\gamma)$ and $q(\beta) < q(\gamma) \Rightarrow \beta < \gamma$.

In the setup of Definition 3.65, if one of the stratifications is admissible of one of our four types then the other one is automatically admissible of the same type. Assuming this is the case, take $\gamma \in \Gamma$ and set $\lambda := q(\gamma)$. We have the stratum $\mathcal{R}_\lambda := \mathcal{R}_{\leq \lambda} / \mathcal{R}_{< \lambda}$ with quotient functor j^λ coming from the original stratification, and the stratum $\mathcal{R}_\gamma := \mathcal{R}_{\leq \gamma} / \mathcal{R}_{< \gamma}$ with quotient functor j^γ coming from the refined stratification⁴. There is also an induced finite stratification $(\rho_\lambda, \mathbf{B}_\lambda, \Gamma, \leq, L_\lambda)$ on \mathcal{R}_λ defined by setting $\rho_\lambda := \rho|_{\mathbf{B}_\lambda}$ and $L_\lambda(b) := j^\lambda L(b)$ for each $b \in \mathbf{B}_\lambda$. We denote the stratum of this labelled by γ by $\mathcal{R}_{\lambda, \gamma}$ with quotient functor $(j_\lambda)^\gamma : \mathcal{R}_{\lambda, \leq \gamma} \rightarrow \mathcal{R}_{\lambda, \gamma}$. In fact, $\mathcal{R}_{\lambda, \gamma}$ may naturally be identified with \mathcal{R}_γ so that $j^\gamma = (j_\lambda)^\gamma \circ j^\lambda|_{\mathcal{R}_{\leq \gamma}}$. Now one can denote the standard and proper objects of \mathcal{R} for the original stratification by

$$\{\rho\Delta(b) := j_!^\lambda P_\lambda(b) \mid \lambda \in \Lambda, b \in \mathbf{B}_\lambda\}, \quad \{\rho\bar{\Delta}(b) := j_!^\lambda L_\lambda(b) \mid \lambda \in \Lambda, b \in \mathbf{B}_\lambda\},$$

and the standard and proper standard objects of \mathcal{R} for the refined stratification by

$$\{\sigma\Delta(b) := j_!^\gamma P_\gamma(b) \mid \gamma \in \Gamma, b \in \mathbf{B}_\gamma\}, \quad \{\sigma\bar{\Delta}(b) := j_!^\gamma L_\gamma(b) \mid \gamma \in \Gamma, b \in \mathbf{B}_\gamma\}.$$

The standard and proper standard objects of \mathcal{R}_λ for its induced stratification are

$$\{\Delta_\lambda(b) := (j_\lambda)^\gamma P_\gamma(b) \mid b \in \bigcup_{\gamma \in q^{-1}(\lambda)} \mathbf{B}_\gamma\}, \quad \{\bar{\Delta}_\lambda(b) := (j_\lambda)^\gamma L_\gamma(b) \mid b \in \bigcup_{\gamma \in q^{-1}(\lambda)} \mathbf{B}_\gamma\},$$

and for such b we have that $\sigma\Delta(b) = j_!^\lambda \Delta_\lambda(b)$, $\sigma\bar{\Delta}(b) = j_!^\lambda \bar{\Delta}_\lambda(b)$ since $j_!^\gamma = j_!^\lambda \circ (j_\lambda)^\gamma$. We deduce for all $b \in \mathbf{B}$ that

$$\rho\Delta(b) \twoheadrightarrow \sigma\Delta(b), \quad \sigma\bar{\Delta}(b) \twoheadrightarrow \rho\bar{\Delta}(b), \quad (3.14)$$

Similar notation can be introduced for the costandard objects, and one sees that

$$\rho\bar{\nabla}(b) \hookrightarrow \sigma\bar{\nabla}(b), \quad \sigma\nabla(b) \hookrightarrow \rho\nabla(b) \quad (3.15)$$

since $j_*^\gamma = j_*^\lambda \circ (j_\lambda)_*^\gamma$.

Lemma 3.66. *Let \mathcal{R} be an Abelian category equipped with an admissible stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Let $(\mathbf{B}, L, \sigma, \Gamma, \leq)$ be a refinement of it in the sense of Definition 3.65.*

- (1) *If \mathcal{R} is fully stratified with respect to the original stratification, and the strata \mathcal{R}_λ are fully stratified with respect to their induced stratifications for all $\lambda \in \Lambda$, then \mathcal{R} is fully stratified with respect to the refined stratification.*

⁴The axiom (R1) is needed so that this notation is unambiguous.

- (2) If \mathcal{R} is fully stratified with respect to the refined stratification, and the functors $j_1^\lambda, j_*^\lambda : \mathcal{R}_\lambda \rightarrow \mathcal{R}_{\leq \lambda}$ are exact for all $\lambda \in \Lambda$, then \mathcal{R} is fully stratified with respect to the original stratification.

Proof. Due to the local nature of the definition of “fully stratified” in the lower finite case, the proof reduces just to the finite, essentially finite and upper finite cases. We assume we are in one of these three situations for the remainder of the argument.

(1) Note that the functors j_1^γ and j_*^γ are exact since they are compositions of exact functors. In view of Lemma 3.20(iv), it remains to show that $P(b)$ has a $\sigma\Delta$ -flag with $\sigma\Delta(b)$ at the top and other sections of the form $\sigma\Delta(c)$ for $c \in \mathbf{B}$ with $\sigma(c) > \sigma(b)$. To see this, let $\lambda := \rho(b)$. As \mathcal{R} is fully stratified with respect to the original stratification, $P(b)$ has a $\rho\Delta$ -flag with $\rho\Delta(b)$ at the top and other sections of the form $\rho\Delta(c)$ for $c \in \mathbf{B}$ with $\rho(c) > \rho(b)$. Moreover each $\rho\Delta(b)$ has a $\sigma\Delta$ -flag with $\sigma\Delta(b)$ at the top and other sections of the form $\sigma\Delta(c)$ for $c \in \mathbf{B}_\lambda$ with $\sigma(c) > \sigma(b)$; this follows by applying the exact functor j_1^λ to a Δ_λ -flag in $P_\lambda(b)$.

(2) To show that \mathcal{R} is fully stratified with respect to the original stratification, both j_1^λ and j_*^λ are exact by assumption, so it suffices to show that each $P(b)$ has a $\rho\bar{\Delta}$ -flag. This follows because $P(b)$ has a $\sigma\bar{\Delta}$ -flag and each $\sigma\bar{\Delta}(b)$ has a $\rho\bar{\Delta}(b)$ -flag; to see the latter assertion apply the exact functor j_1^λ to a composition series of $\bar{\Delta}_\lambda(b)$. \square

Corollary 3.67. *Let \mathcal{R} be fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Assume that each stratum \mathcal{R}_λ ($\lambda \in \Lambda$) is a highest weight category with weight poset $(\Gamma_\lambda, \leq_\lambda)$ and labelling function L_λ . Let $\Gamma := \bigsqcup_{\lambda \in \Lambda} \Gamma_\lambda$, $\sigma : \mathbf{B} \xrightarrow{\sim} \Gamma$ be the bijection such that $j^\lambda L(b) \cong L_\lambda(\sigma(b))$ for $b \in \mathbf{B}_\lambda$, and \leq be the partial order on Γ defined by $\sigma(b) \leq \sigma(c)$ if and only if either $\rho(b) < \rho(c)$, or $\lambda := \rho(b) = \rho(c)$ and $\sigma(b) \leq_\lambda \sigma(c)$. Then $(\mathbf{B}, L, \sigma, \Gamma, \leq)$ is a refinement of the original stratification which makes \mathcal{R} into a highest weight category.*

Remark 3.68. It is also interesting to consider changing the underlying partial order on the set Λ . For a fully stratified category \mathcal{R} with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$, one can always replace the given order \leq by the *minimal order* \leq , that is, the partial order generated by the relation that $\lambda < \mu$ if $[\nabla(b) : L(c)] + [\bar{\Delta}(b) : L(c)] \neq 0$ for some $b \in \mathbf{B}_\lambda, c \in \mathbf{B}_\mu$. Then \mathcal{R} is also fully stratified with respect to $(\mathbf{B}, L, \rho, \Lambda, \leq)$ with all the same strata, standard objects, etc.. For highest weight categories, Coulembier [Cou2], [Cou3] has made the following elegant observation: if \mathcal{R} is a finite Abelian, locally finite Abelian or Schurian category, $\{L(\lambda) \mid \lambda \in \Lambda\}$ is a full set of pairwise inequivalent irreducible objects, and \mathcal{R} possesses a contravariant autoequivalence preserving isomorphism classes of irreducible objects, then all partial orders on Λ making \mathcal{R} into a highest weight category give rise to the same minimal order. There are examples showing that this statement is false for essentially finite highest weight categories.

4. TILTING MODULES AND SEMI-INFINITE RINGEL DUALITY

We now develop the theory of tilting objects and Ringel duality. Even in the finite case, we are not aware of a complete exposition of these results in the existing literature in the general ε -stratified setting.

4.1. Tilting objects in the finite and lower finite cases. In this subsection, \mathcal{R} is a finite or locally finite Abelian category with a finite or lower finite stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$, and $\varepsilon : \Lambda \rightarrow \{\pm\}$ is a fixed sign function with respect to which \mathcal{R} is a finite or lower finite ε -stratified category, respectively; see Definitions 3.7 and 3.50. By an ε -tilting object, we mean an object of the following full subcategory of \mathcal{R} :

$$\mathcal{Tilt}_\varepsilon(\mathcal{R}) := \Delta_\varepsilon(\mathcal{R}) \cap \nabla_\varepsilon(\mathcal{R}). \quad (4.1)$$

The following shows that the additive subcategory $\mathcal{Tilt}_\varepsilon(\mathcal{R})$ of \mathcal{R} is Karoubian.

Lemma 4.1. *Direct summands of ε -tilting objects are ε -tilting objects.*

Proof. This follows easily from the homological criteria from Theorems 3.11 and 3.14. In the lower finite case, one needs to pass first to a finite ε -stratified subcategory containing the object in question using Theorem 3.59. \square

The next goal is to construct and classify ε -tilting objects. Our exposition of this is based roughly on [Don4, Appendix], which in turn goes back to the work of Ringel [Rin]. There are some additional complications in the ε -stratified setting.

Theorem 4.2 (Classification of ε -tilting objects). *Assume that \mathcal{R} is a finite or lower finite ε -stratified category. For $b \in \mathbf{B}_\lambda$ there is an indecomposable object $T_\varepsilon(b) \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$ satisfying the following properties:*

- (i) $T_\varepsilon(b)$ has a Δ_ε -flag with bottom section isomorphic to $\Delta_\varepsilon(b)$;
- (ii) $T_\varepsilon(b)$ has a ∇_ε -flag with top section isomorphic to $\nabla_\varepsilon(b)$;
- (iii) $T_\varepsilon(b) \in \mathcal{R}_{\leq \lambda}$ and $j^\lambda T_\varepsilon(b) \cong \begin{cases} P_\lambda(b) & \text{if } \varepsilon(\lambda) = + \\ I_\lambda(b) & \text{if } \varepsilon(\lambda) = - \end{cases}$.

These properties determine $T_\varepsilon(b)$ uniquely up to isomorphism: if U is any indecomposable object of $\mathcal{Tilt}_\varepsilon(\mathcal{R})$ satisfying any one of the properties (i)–(iii) then $U \cong T_\varepsilon(b)$; hence, it satisfies the other two properties as well.

Proof. By replacing \mathcal{R} by the Serre subcategory associated to a sufficiently large but finite lower set Λ^\dagger in Λ , chosen so as to contain λ and (for the uniqueness statement) all $\rho(b)$ for b such that $[T : L(b)] \neq 0$, one reduces to the case that \mathcal{R} is a finite ε -stratified category. This reduction depends only on Theorem 3.59. Thus, we may assume henceforth that Λ is finite.

Existence: The main step is to construct an indecomposable object $T_\varepsilon(b) \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$ such that (iii) holds. The argument for this proceeds by induction on $|\Lambda|$. If $\lambda \in \Lambda$ is minimal, we set $T_\varepsilon(b) := \Delta(b)$ if $\varepsilon(\lambda) = +$ or $\nabla(b)$ if $\varepsilon(\lambda) = -$. Since $\bar{\Delta}(b) = L(b) = \bar{\nabla}(b)$ by the minimality of λ , this has both a Δ_ε - and a ∇_ε -flag. It is indecomposable, and we get (iii) from Lemma 2.27.

For the induction step, suppose that λ is not minimal and pick $\mu < \lambda$ that is minimal. Let $\Lambda^\dagger := \Lambda \setminus \{\mu\}$, $\mathbf{B}^\dagger := \rho^{-1}(\Lambda^\dagger)$, and $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be the corresponding Serre quotient. By induction, there is an indecomposable object $T_\varepsilon^\dagger(b) \in \mathcal{Tilt}_\varepsilon(\mathcal{R}^\dagger)$ satisfying the analog of (iii). Now there are two cases according to whether $\varepsilon(\mu) = +$ or $-$.

Case $\varepsilon(\mu) = +$: For any $V \in \mathcal{R}$, let $d_+(V) := \sum_{c \in \mathbf{B}_\mu} \dim \text{Ext}_{\mathcal{R}}^1(\Delta(c), V)$. We recursively construct $n \geq 0$ and T_0, T_1, \dots, T_n so that $d_+(T_0) > d_+(T_1) > \dots > d_+(T_n) = 0$ and the following properties hold for all m :

- (1) $T_m \in \Delta_\varepsilon(\mathcal{R})$.
- (2) $j^\lambda T_m \cong P_\lambda(b)$ if $\varepsilon(\lambda) = +$ or $I_\lambda(b)$ if $\varepsilon(\lambda) = -$.
- (3) $\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(a), T_m) = 0$ for all $a \in \mathbf{B} \setminus \mathbf{B}_\mu$.

To start with, set $T_0 := j_! T_\varepsilon^\dagger(b)$. This satisfies all of the above properties: (1) follows from Theorem 3.18(6); (2) follows because j^λ factors through j and we know that $T_\varepsilon^\dagger(b)$ satisfies the analogous property; (3) follows by Theorem 3.18(5). For the recursive step, assume that we are given T_m satisfying (1), (2) and (3) and $d_+(T_m) > 0$. We can find $c \in \mathbf{B}_\mu$ and a non-split extension

$$0 \longrightarrow T_m \longrightarrow T_{m+1} \longrightarrow \Delta(c) \longrightarrow 0. \quad (4.2)$$

This constructs T_{m+1} . We claim that $d_+(T_{m+1}) < d_+(T_m)$ and that T_{m+1} satisfies (1), (2) and (3) too. Part (1) is clear from the definition. For (2), we just apply the exact functor j^λ to the exact sequence (4.2), noting that $j^\lambda \Delta(c) = 0$. For (3), take $a \in \mathbf{B} \setminus \mathbf{B}_\mu$ and apply the functor $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(a), ?)$ to the short exact sequence (4.2) to get

$$\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(a), T_m) \longrightarrow \text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(a), T_{m+1}) \longrightarrow \text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(a), \Delta(c)).$$

The first and last term are zero by hypothesis and (3.10), implying $\text{Ext}_{\mathcal{R}}^1(T_{m+1}, \nabla_{\varepsilon}(a)) = 0$. It remains to show $d_+(T_{m+1}) < d_+(T_m)$. For $a \in \mathbf{B}_{\mu}$, we have $\text{Ext}_{\mathcal{R}}^1(\Delta(a), \Delta(c)) = 0$ by (3.10), so again we have an exact sequence

$$\text{Hom}_{\mathcal{R}}(\Delta(a), \Delta(c)) \xrightarrow{f} \text{Ext}_{\mathcal{R}}^1(\Delta(a), T_m) \longrightarrow \text{Ext}_{\mathcal{R}}^1(\Delta(a), T_{m+1}) \longrightarrow 0.$$

This shows that $\dim \text{Ext}_{\mathcal{R}}^1(\Delta(a), T_{m+1}) \leq \dim \text{Ext}_{\mathcal{R}}^1(\Delta(a), T_m)$, and we just need to observe that the inequality is actually a strict one in the case $a = c$. To see this, note that the first morphism f is non-zero in the case $a = c$ as $f(\text{id}_{\Delta(c)}) \neq 0$ due to the fact that the original short exact sequence was not split. This completes the proof of the claim. We have now defined an object $T_n \in \Delta_{\varepsilon}(\mathcal{R})$ such that $j^{\lambda}T_n \cong P_{\lambda}(b)$ if $\varepsilon(\lambda) = +$ or $I_{\lambda}(b)$ if $\varepsilon(\lambda) = -$, and moreover $\text{Ext}_{\mathcal{R}}^1(\Delta_{\varepsilon}(a), T_n) = 0$ for all $a \in \mathbf{B}$. By Theorem 3.11, we deduce that $T_n \in \nabla_{\varepsilon}(\mathcal{R}_{\leq \lambda})$ too, hence, it is an ε -tilting object. Decompose T_n into indecomposables $T_n = T_{n,1} \oplus \cdots \oplus T_{n,r}$. Then each $T_{n,i}$ is also an ε -tilting object by Lemma 4.1. Since $j^{\lambda}T_n$ is indecomposable, we must have that $j^{\lambda}T_n = j^{\lambda}T_{n,i}$ for some unique i . Then we set $T_{\varepsilon}(b) := T_{n,i}$ for this i . This gives us the desired indecomposable ε -tilting object.

Case $\varepsilon(\mu) = -$: Let $d_-(V) := \sum_{c \in \mathbf{B}_{\mu}} \dim \text{Ext}_{\mathcal{R}}^1(V, \nabla(c))$. This time, one recursively constructs $T_0 := j_*T_{\varepsilon}^{\dagger}(b), T_1, \dots, T_n$ so that $d_-(T_0) > \cdots > d_-(T_n) = 0$ and

- (1') $T_m \in \nabla_{\varepsilon}(\mathcal{R})$.
- (2') $j^{\lambda}T_m \cong P_{\lambda}(b)$ if $\varepsilon(\lambda) = +$ or $I_{\lambda}(b)$ if $\varepsilon(\lambda) = -$.
- (3') $\text{Ext}_{\mathcal{R}}^1(T_m, \nabla_{\varepsilon}(a)) = 0$ for all $a \in \mathbf{B} \setminus \mathbf{B}_{\mu}$.

Since this is just the dual construction to the case $\varepsilon(\mu) = +$ already treated, i.e., it is the same construction in the opposite category, we omit the details. Then, at the end, one decomposes T_n into indecomposables $T_n = T_{n,1} \oplus \cdots \oplus T_{n,r}$. By Theorem 3.14 each $T_{n,i}$ is an ε -tilting object. Since $j^{\lambda}T_n$ is indecomposable, we must have that $j^{\lambda}T_n = j^{\lambda}T_{n,i}$ for some unique i , and set $T_{\varepsilon}(b) := T_{n,i}$ for this i .

This completes the construction of $T_{\varepsilon}(b)$ in general. We have shown it satisfies (iii). Let us show that it also satisfies (i) and (ii). For (i), we know by (iii) that $T_{\varepsilon}(b)$ belongs to $\mathcal{R}_{\leq \lambda}$, and it has a Δ_{ε} -flag. By (3.10), we may order this flag so that the sections $\Delta_{\varepsilon}(c)$ for $c \in \mathbf{B}_{\lambda}$ appear at the bottom. Thus, there is a short exact sequence $0 \rightarrow K \rightarrow T_{\varepsilon}(b) \rightarrow Q \rightarrow 0$ such that K has a Δ_{ε} -flag with sections $\Delta_{\varepsilon}(c)$ for $c \in \mathbf{B}_{\lambda}$ and $j^{\lambda}Q = 0$. Then $j^{\lambda}K \cong j^{\lambda}T_{\varepsilon}(b)$. If $\varepsilon(\lambda) = +$, this is $P_{\lambda}(b)$. Since j^{λ} is exact and $j^{\lambda}\Delta(c) = P_{\lambda}(c)$ for each $c \in \mathbf{B}_{\lambda}$, we must have that $K \cong \Delta(b)$, and (1) follows. Instead, if $\varepsilon(\lambda) = -$, the bottom section of the $\bar{\nabla}$ -flag of K must be $\bar{\nabla}(b)$ since $j^{\lambda}K \cong I_{\lambda}(b)$ has irreducible socle $L_{\lambda}(b)$, giving (i) in this case too. The proof of (ii) is similar.

Uniqueness: Let $T := T_{\varepsilon}(b)$ and U be some other indecomposable object of $\mathcal{Tilt}_{\varepsilon}(\mathcal{R})$ satisfying one of the properties (i)–(iii). We must prove that $T \cong U$. By the argument from the previous paragraph, we may assume actually that U satisfies either (i) or (ii). We just explain how to see this in the case that U satisfies (i); the dual argument treats the case that U satisfies (ii). So there are short exact sequences $0 \rightarrow \Delta_{\varepsilon}(b) \xrightarrow{f} U \rightarrow Q_1 \rightarrow 0$ and $0 \rightarrow \Delta_{\varepsilon}(b) \xrightarrow{g} T \rightarrow Q_2 \rightarrow 0$ such that Q_1, Q_2 have Δ_{ε} -flags. Applying $\text{Hom}_{\mathcal{R}}(?, T)$ to the first and using $\text{Ext}_{\mathcal{R}}^1(Q_1, T) = 0$, we get that $\text{Hom}_{\mathcal{R}}(U, T) \rightarrow \text{Hom}_{\mathcal{R}}(\Delta_{\varepsilon}(b), T)$. Hence, g extends to a homomorphism $\bar{g} : U \rightarrow T$. Similarly, f extends to $\bar{f} : T \rightarrow U$. We have constructed morphisms making the triangles in the following diagram commute:

$$\begin{array}{ccc} & & U \\ & \nearrow f & \uparrow \bar{f} \\ \Delta_{\varepsilon}(b) & & \\ & \searrow g & \downarrow \bar{g} \\ & & T \end{array}$$

Since $\bar{f} \circ \bar{g} \circ f = f$, we deduce that $\bar{f} \circ \bar{g}$ is not nilpotent. Since U is indecomposable, Fitting's Lemma implies $\bar{f} \circ \bar{g}$ is an isomorphism. Similarly, so is $\bar{g} \circ \bar{f}$. Hence, $U \cong T$. \square

Remark 4.3. Let $b \in \mathbf{B}_\lambda$. When $\varepsilon(\lambda) = +$, Theorem 4.2 implies that $(T_\varepsilon(b) : \Delta_\varepsilon(b)) = 1$ and $(T_\varepsilon(b) : \Delta_\varepsilon(c)) = 0$ for all other $c \in \mathbf{B}_\lambda$. Similarly, when $\varepsilon(\lambda) = -$, we have that $(T_\varepsilon(b) : \nabla_\varepsilon(b)) = 1$ and $(T_\varepsilon(b) : \nabla_\varepsilon(c)) = 0$ for all other $c \in \mathbf{B}_\lambda$.

The following corollaries show that ε -tilting objects behave well with respect to passage to lower and upper sets, extending Theorems 3.17, 3.59 and 3.18. This follows easily from those theorems plus the characterization of tilting objects in Theorem 4.2; the situation is just like [Don4, Lem. A4.5].

Corollary 4.4. *Let \mathcal{R} be a finite or lower finite ε -stratified category and \mathcal{R}^\downarrow be the finite ε -stratified subcategory associated to a finite lower set Λ^\downarrow of Λ . For $b \in \mathbf{B}^\downarrow := \rho^{-1}(\Lambda^\downarrow)$, the corresponding indecomposable ε -tilting object of \mathcal{R}^\downarrow is $T_\varepsilon(b)$ (the same as in \mathcal{R}).*

Corollary 4.5. *Assume \mathcal{R} is a finite ε -stratified category and let Λ^\uparrow be an upper set in Λ with associated quotient $j : \mathcal{R} \rightarrow \mathcal{R}^\uparrow$. Let $b \in \mathbf{B}^\uparrow := \rho^{-1}(\Lambda^\uparrow)$. The corresponding indecomposable ε -tilting object $T_\varepsilon^\uparrow(b)$ of \mathcal{R}^\uparrow satisfies $T_\varepsilon^\uparrow(b) \cong jT_\varepsilon(b)$. Also $jT_\varepsilon(b) = 0$ if $b \notin \mathbf{B}^\uparrow$.*

The next result is concerned with tilting resolutions.

Definition 4.6. Assume that \mathcal{R} is a finite or lower finite ε -stratified category. An ε -tilting resolution $d : T_\bullet \rightarrow V$ of $V \in \mathcal{R}$ is the data of an exact sequence

$$\cdots \xrightarrow{d_2} T_1 \xrightarrow{d_1} T_0 \xrightarrow{d_0} V \longrightarrow 0$$

such that

- (TR1) $T_m \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$ for each $m = 0, 1, \dots$;
- (TR2) $\text{im } d_m \in \nabla_\varepsilon(\mathcal{R})$ for $m \gg 0$.

Similarly, an ε -tilting coresolution $d : V \rightarrow T^\bullet$ of $V \in \mathcal{R}$ is the data of an exact sequence

$$0 \longrightarrow V \xrightarrow{d^0} T^0 \xrightarrow{d^1} T^1 \xrightarrow{d^2} \cdots$$

such that

- (TC1) $T^m \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$ for $m = 0, 1, \dots$;
- (TC2) $\text{coim } d^m \in \Delta_\varepsilon(\mathcal{R})$ for $m \gg 0$.

We say it is a *finite* resolution (resp., coresolution) if there is some n such that $T_m = 0$ (resp., $T^m = 0$) for $m > n$. Note in the finite case that axioms (TR2) and (TC2) are redundant since the zero object belongs to both $\nabla_\varepsilon(\mathcal{R})$ and $\Delta_\varepsilon(\mathcal{R})$.

Lemma 4.7. *Assume that \mathcal{R} is a finite or lower finite ε -stratified category.*

- (1) *If $d : T_\bullet \rightarrow V$ is an ε -tilting resolution of $V \in \mathcal{R}$ then $\text{im } d_m \in \nabla_\varepsilon(\mathcal{R})$ for all $m \geq 0$. In particular, $V \in \nabla_\varepsilon(\mathcal{R})$.*
- (2) *If $d : V \rightarrow T^\bullet$ is an ε -tilting coresolution of $V \in \mathcal{R}$ then $\text{coim } d^m \in \Delta_\varepsilon(\mathcal{R})$ for all $m \geq 0$. In particular, $V \in \Delta_\varepsilon(\mathcal{R})$.*

Proof. (1) It suffices to show for any exact sequence $A \xrightarrow{f} B \xrightarrow{g} C$ in a finite or lower finite ε -stratified category that $B \in \nabla_\varepsilon(\mathcal{R})$ and $\text{im } f \in \nabla_\varepsilon(\mathcal{R})$ implies $\text{im } g \in \nabla_\varepsilon(\mathcal{R})$. Since $\text{im } f = \ker g$, there is a short exact sequence $0 \rightarrow \text{im } f \rightarrow B \rightarrow \text{im } g \rightarrow 0$. Now apply Corollary 3.13 (or Corollary 3.58).

(2) An ε -tilting coresolution of V in \mathcal{R} is the same thing as a $(-\varepsilon)$ -tilting resolution of V in \mathcal{R}^{op} . Hence, this follows as it is the dual statement to (1). \square

Theorem 4.8 (Tilting resolutions and coresolutions). *Let \mathcal{R} be a finite or lower finite ε -stratified category and take $V \in \mathcal{R}$.*

- (1) *V has an ε -tilting resolution if and only if $V \in \nabla_\varepsilon(\mathcal{R})$.*
- (2) *V has an ε -tilting coresolution if and only if $V \in \Delta_\varepsilon(\mathcal{R})$.*

Proof. We just prove (1), since (2) is the equivalent dual statement. If V has an ε -tilting resolution, then we must have that $V \in \nabla_\varepsilon(\mathcal{R})$ thanks to Lemma 4.7(1). For the converse, we claim for $V \in \nabla_\varepsilon(\mathcal{R})$ that there is a short exact sequence $0 \rightarrow S_V \rightarrow T_V \rightarrow V \rightarrow 0$ with $S_V \in \nabla_\varepsilon(\mathcal{R})$ and $T_V \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$. Given the claim, one can construct an ε -tilting resolution of V by ‘‘Splicing’’ (e.g., see [Wei, Fig. 2.1]), to complete the proof.

To prove the claim, we argue by induction on the length $\sum_{b \in \mathbf{B}} (V : \nabla_\varepsilon(b))$ of a ∇_ε -flag of V . If this number is one, then $V \cong \nabla_\varepsilon(b)$ for some $b \in \mathbf{B}$, and there is a short exact sequence $0 \rightarrow S_V \rightarrow T_V \rightarrow V \rightarrow 0$ with $S_V \in \nabla_\varepsilon(b)$ and $T_V := T_\varepsilon(b)$ due to Theorem 4.2(ii). If it is greater than one, then there is a short exact sequence $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$ such that U and W have strictly shorter ∇_ε -flags. By induction, there are short exact sequences $0 \rightarrow S_U \rightarrow T_U \rightarrow U \rightarrow 0$ and $0 \rightarrow S_W \rightarrow T_W \rightarrow W \rightarrow 0$ with $S_U, S_W \in \nabla_\varepsilon(\mathcal{R})$ and $T_U, T_W \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$. It remains to show that these short exact sequences can be assembled to produce the desired short exact sequence for V . The argument is like in the proof of the Horseshoe Lemma in [Wei, Lem. 2.2.8].

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & S_U & \longrightarrow & T_U & \xrightarrow{i} & U \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow f \\
0 & \longrightarrow & S_V & \longrightarrow & T_V & \xrightarrow{j} & V \longrightarrow 0 \\
& & \downarrow & & \downarrow & \nearrow \hat{k} & \downarrow g \\
0 & \longrightarrow & S_W & \longrightarrow & T_W & \xrightarrow{k} & W \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array} \tag{4.3}$$

Since $\text{Ext}_{\mathcal{R}}^1(T_W, U) = 0$, we can lift $k : T_W \rightarrow W$ to $\hat{k} : T_W \rightarrow V$ so that $k = g \circ \hat{k}$. Let $T_V := T_U \oplus T_W$ and $j : T_V \rightarrow V$ be $\text{diag}(fi, \hat{k})$. This gives us a split short exact sequence in the middle column in (4.3), such that the right hand squares commute. Then we let $S_V := \ker j$, and see that there are induced maps making the left hand column and middle row into short exact sequences such that the left hand squares commute too. \square

4.2. Finite Ringel duality. In this subsection, we review the theory of Ringel duality for finite ε -stratified categories. Our exposition is based in part on [Don4, Appendix], which gives a self-contained treatment in the highest weight setting, and [AHLU], where the $+$ -highest weight case is considered assuming $\Lambda = \{1 < \dots < n\}$; the survey in [Rei, Ch. 3] is also helpful. Throughout, we assume that \mathcal{R} is a finite ε -stratified category with the usual stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$.

Definition 4.9. Let \mathcal{R} be a finite ε -stratified category. By an ε -tilting generator T for \mathcal{R} , we mean an object $T \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$ such that T has a summand isomorphic to $T_\varepsilon(b)$ for each $b \in \mathbf{B}$. Given such an object, we define the *Ringel dual* of \mathcal{R} relative to T to be the finite Abelian category $\mathcal{R}' := B\text{-mod}_{\text{fd}}$ where $B := \text{End}_{\mathcal{R}}(T)^{\text{op}}$. We also define the two (covariant) *Ringel duality functors*

$$F := \text{Hom}_{\mathcal{R}}(T, ?) : \mathcal{R} \rightarrow \mathcal{R}', \tag{4.4}$$

$$G := \text{Cohom}_{\mathcal{R}}(T, ?) = \text{Hom}_{\mathcal{R}}(?, T)^* : \mathcal{R} \rightarrow \mathcal{R}'. \tag{4.5}$$

Note for the second of these that $\text{Hom}_{\mathcal{R}}(V, T)$ is naturally a finite-dimensional right B -module for $V \in \mathcal{R}$, hence, its dual is a left B -module.

Theorem 4.10 (Finite Ringel duality). *In the setup of Definition 4.9, the Ringel dual \mathcal{R}' of \mathcal{R} relative to T is a finite $(-\varepsilon)$ -stratified category with stratification $(\mathbf{B}, L', \rho, \Lambda, \geq)$*

and distinguished objects

$$\begin{aligned} P'(b) &= FT_\varepsilon(b), & I'(b) &= GT_\varepsilon(b), & L'(b) &= \text{hd } P'(b) \cong \text{soc } I'(b), \\ \Delta'_{-\varepsilon}(b) &= F\nabla_\varepsilon(b), & \nabla'_{-\varepsilon}(b) &= G\Delta_\varepsilon(b), & T'_{-\varepsilon}(b) &= FI(b) \cong GP(b). \end{aligned}$$

The restrictions $F : \nabla_\varepsilon(\mathcal{R}) \rightarrow \Delta_{-\varepsilon}(\mathcal{R}')$ and $G : \Delta_\varepsilon(\mathcal{R}) \rightarrow \nabla_{-\varepsilon}(\mathcal{R}')$ are equivalences; in fact, they induce isomorphisms

$$\text{Ext}_{\mathcal{R}}^n(V_1, V_2) \cong \text{Ext}_{\mathcal{R}'}^n(FV_1, FV_2), \quad \text{Ext}_{\mathcal{R}}^n(W_1, W_2) \cong \text{Ext}_{\mathcal{R}'}^n(GW_1, GW_2), \quad (4.6)$$

for all $V_i \in \nabla_\varepsilon(\mathcal{R})$, $W_i \in \Delta_\varepsilon(\mathcal{R})$ and $n \geq 0$.

Before the proof, we give some applications.

Corollary 4.11 (Double centralizer property). *Suppose that the finite ε -stratified category \mathcal{R} in Theorem 4.10 is the category $A\text{-mod}_{\text{fd}}$ for a finite-dimensional algebra A , so that T is an (A, B) -bimodule. Let $T' := T^*$ be the dual (B, A) -bimodule. Then the following holds.*

- (1) T' is a $(-\varepsilon)$ -tilting generator for $\mathcal{R}' = B\text{-mod}_{\text{fd}}$ and there is an algebra isomorphism

$$\mu : A \xrightarrow{\sim} \text{End}_{\mathcal{R}'}(T')^{\text{op}} \quad (4.7)$$

sending $x \in A$ to $\mu(x) : T' \rightarrow T', v \mapsto vx$. So the Ringel dual of \mathcal{R}' relative to T' is equivalent to the original category \mathcal{R} .

- (2) Denote the Ringel duality functors for \mathcal{R}' relative to T' now by

$$G_* := \text{Hom}_{\mathcal{R}'}(T', ?) : \mathcal{R}' \rightarrow \mathcal{R}, \quad (4.8)$$

$$F^* := \text{Cohom}_{\mathcal{R}'}(T', ?) = \text{Hom}_{\mathcal{R}'}(?, T')^* : \mathcal{R}' \rightarrow \mathcal{R}. \quad (4.9)$$

We have that $F^* \cong T \otimes_B ?$ and $G \cong T' \otimes_A ?$, hence, (F^*, F) and (G, G_*) are adjoint pairs.

Proof. (1) Note that GA is a $(-\varepsilon)$ -tilting generator since $GP(b) \cong T'_{-\varepsilon}(b)$ for $b \in \mathbf{B}$. Actually, $GA = \text{Hom}_A(A, T)^* \cong T^* = T'$. Thus, T' is a $(-\varepsilon)$ -tilting generator for \mathcal{R}' . Its opposite endomorphism algebra is isomorphic to A as stated since G defines an algebra isomorphism

$$A \cong \text{End}_A(A)^{\text{op}} \xrightarrow{\sim} \text{End}_B(GA)^{\text{op}} \cong \text{End}_B(T')^{\text{op}}.$$

(2) As F^* is right exact and commutes with direct sums, a standard argument using the Five Lemma shows that it is isomorphic to $(F^*B) \otimes_B ? \cong T \otimes_B ?$. Thus, F^* is left adjoint to F . Similarly, $G \cong T' \otimes_A ?$ is left adjoint to G_* . \square

The next corollary describes the strata \mathcal{R}'_λ of the Ringel dual category; see also Lemma 4.41 below. For $\lambda \in \Lambda$, denote the quotient functor $\mathcal{R}'_{\geq \lambda} \rightarrow \mathcal{R}'_\lambda$ by $(j')^\lambda$, and denote its left and right adjoints by $(j')^\lambda_! : \mathcal{R}'_\lambda \rightarrow \mathcal{R}'_{\geq \lambda}$ and $(j')^\lambda_* : \mathcal{R}'_\lambda \rightarrow \mathcal{R}'_{\geq \lambda}$. We also have the inclusion $(i')_{\geq \lambda} : \mathcal{R}'_{\geq \lambda} \rightarrow \mathcal{R}'$ with left and right adjoints $(i')^*_{\geq \lambda}$ and $(i')^!_{\geq \lambda}$.

Corollary 4.12. *For $\lambda \in \Lambda$, the strata \mathcal{R}_λ and \mathcal{R}'_λ are equivalent. More precisely:*

- (1) If $\varepsilon(\lambda) = +$ the functor $F_\lambda := (j')^\lambda \circ (i')^!_{\geq \lambda} \circ F \circ i_{\leq \lambda} \circ j^*_\lambda : \mathcal{R}_\lambda \rightarrow \mathcal{R}'_\lambda$ is an equivalence of categories taking $L_\lambda(b) = j^\lambda L(b)$ to $L'_\lambda(b) = (j')^\lambda L'(b)$.
(2) If $\varepsilon(\lambda) = -$ the functor $G_\lambda := (j')^\lambda \circ (i')^*_{\geq \lambda} \circ G \circ i_{\leq \lambda} \circ j^\lambda_! : \mathcal{R}_\lambda \rightarrow \mathcal{R}'_\lambda$ is an equivalence of categories taking $L_\lambda(b) = j^\lambda L(b)$ to $L'_\lambda(b) = (j')^\lambda L'(b)$.

Proof. We just prove (1), since (2) is similar. So assume that $\varepsilon(\lambda) = +$. We first note that F_λ is exact. Indeed, j^*_λ is exact by Theorem 3.5, so it sends objects of \mathcal{R}_λ to objects of $\mathcal{R}_{\leq \lambda}$ which have filtrations with sections $\nabla_\varepsilon(b)$ for $b \in \mathbf{B}_\lambda$. Then we apply the exact functor $i_{\leq \lambda}$ followed by F , which takes short exact sequences in $\nabla_\varepsilon(\mathcal{R})$ to short exact sequences in $\Delta_\varepsilon(\mathcal{R})$, to obtain an object of $\Delta_{-\varepsilon}(\mathcal{R}'_{\geq \lambda})$. The functor $(i')^!_{\geq \lambda}$ is the identity

on this subcategory, and finally $(j')^\lambda$ is exact. Adopting the setup of Corollary 4.11, we can also define

$$F_\lambda^* := j^\lambda \circ i_{\leq \lambda}^* \circ F^* \circ (i')_{\geq \lambda} \circ (j')_!^\lambda : \mathcal{R}'_\lambda \rightarrow \mathcal{R}_\lambda.$$

A similar argument to before gives that this is exact too. We complete the proof by showing that F_λ and F_λ^* are quasi-inverse equivalences. Note that F_λ^* is left adjoint to F_λ . The counit of adjunction gives us a natural transformation $F_\lambda^* \circ F_\lambda \rightarrow \text{Id}_{\mathcal{R}_\lambda}$. We claim this is an isomorphism. Since both functors are exact, it suffices to prove this on irreducible objects: we have $F_\lambda^*(F_\lambda L_\lambda(b)) \cong F_\lambda^* L'_\lambda(b) \cong L_\lambda(b)$. Similar argument shows that the unit of adjunction is an isomorphism in the other direction. \square

Corollary 4.13. *Let \mathcal{R} be a finite ε -stratified category.*

- (1) *All $V \in \nabla_\varepsilon(\mathcal{R})$ have finite ε -tilting resolutions if and only if all positive strata are of finite global dimension.*
- (2) *All $V \in \Delta_\varepsilon(\mathcal{R})$ have finite ε -tilting coresolutions if and only if all negative strata are of finite global dimension.*

Proof. We just explain the proof of (1). By Theorem 4.10, all $V \in \nabla_\varepsilon(\mathcal{R})$ have finite ε -tilting resolutions if and only if all $V' \in \Delta_{-\varepsilon}(\mathcal{R}')$ have finite projective resolutions. By Lemma 3.23(1), this is equivalent to all negative strata of the $(-\varepsilon)$ -stratified category \mathcal{R}' are of finite global dimension. Equivalently, by Corollary 4.12, all positive strata of the ε -stratified category \mathcal{R} are of finite global dimension. \square

Corollary 4.14. *If \mathcal{R} is a finite $+$ -stratified (resp., $--$ -stratified) category then all $V \in \Delta(\mathcal{R})$ (resp., $V \in \nabla(\mathcal{R})$) have finite $+$ -tilting coresolutions (resp., finite $--$ -tilting resolutions).*

The next theorem is a consequence of Happel's tilting theory for finite-dimensional algebras. To prepare for this, we explain the connection between ε -tilting objects in our setting and the general notions of tilting and cotilting modules from that theory; e.g., see [Hap], [Rei]. Suppose that $\mathcal{R} = A\text{-mod}_{\text{fd}}$ is a finite ε -stratified algebra for a finite-dimensional algebra A , and let T be an ε -tilting generator for \mathcal{R} . If all negative strata are of finite global dimension (this assumption being vacuous in the case $\varepsilon = +$) then T is a *tilting module* in the sense of tilting theory; if all positive strata are of finite global dimension (this assumption being vacuous in the case $\varepsilon = -$) then T is a *cotilting module*. These assertions follow using Theorem 3.11 to see that $\text{Ext}_{\mathcal{R}}^1(T, T) = 0$, Lemma 3.23 to see that $\text{pd } T < \infty$ or $\text{id } T < \infty$, and Corollary 4.13. Without assumptions on the global dimensions of strata, T need not be tilting or cotilting, but Theorem 4.8 implies that it is still an example of a *Wakamatsu tilting module*⁵ as defined in [BR, Ch. 3]; see also [Rei, §4.1]. The *WT-conjecture* formulated in [BR, Ch. 3] is the assertion that any Wakamatsu tilting module of finite projective (resp., injective) dimension is tilting (resp., cotilting). This motivates the following conjecture in our special situation; we will prove this assuming a mild additional hypothesis on strata in Lemma 4.38 below.

Conjecture 4.15 (ε T-conjecture). *Suppose that \mathcal{R} is a finite fully stratified category and ε is a given sign function. For $b \in \mathbf{B}$, the ε -tilting module $T_\varepsilon(b)$ is of finite projective (resp., injective) dimension if and only if $T_\varepsilon(b)$ belongs to $\mathcal{Tilt}_+(\mathcal{R})$ (resp., $\mathcal{Tilt}_-(\mathcal{R})$).*

Let $\mathbb{R}F$ and $\mathbb{L}G$ be the total derived functors of the Ringel duality functors. These are triangulated functors between the bounded derived categories $D^b(\mathcal{R})$ and $D^b(\mathcal{R}')$.

Theorem 4.16 (Derived equivalences). *Let \mathcal{R}' be the Ringel dual of a finite ε -stratified category \mathcal{R} . Assume that all negative strata (resp., all positive strata) of \mathcal{R} are of finite global dimension. Then $\mathbb{R}F : D^b(\mathcal{R}) \rightarrow D^b(\mathcal{R}')$ (resp., $\mathbb{L}G : D^b(\mathcal{R}) \rightarrow D^b(\mathcal{R}')$) is an*

⁵With this in mind, the fact that the map (4.7) is an isomorphism could also be deduced from [Wak, Cor. 2].

equivalence of triangulated categories. Moreover, if \mathcal{R} is of finite global dimension, then so is \mathcal{R}' .

Proof. Assuming \mathcal{R} has finite global dimension, this all follows by [Hap, Lem. 2.9, Th. 2.10]; the hypotheses there hold thanks to Corollary 4.13. To get the derived equivalence without assuming \mathcal{R} has finite global dimension, we cite instead Keller's exposition of Happel's result in [Kel, Th. 4.1], since it assumes slightly less; the hypotheses (a) and (c) there hold due to Corollary 4.13(2) and Lemma 3.23(1). \square

Corollary 4.17. *If \mathcal{R} is $+$ -highest weight (resp., $-$ -highest weight) and \mathcal{R}' is the Ringel dual relative to a $+$ -tilting generator (resp., $-$ -tilting generator), then $\mathbb{R}F : D^b(\mathcal{R}) \rightarrow D^b(\mathcal{R}')$ (resp., $\mathbb{L}G : D^b(\mathcal{R}) \rightarrow D^b(\mathcal{R}')$) is an equivalence.*

Proof of Theorem 4.10. This follows the same steps as in [Don4, pp.158–160]. Assume without loss of generality that $\mathcal{R} = A\text{-mod}_{\text{fd}}$ for a finite-dimensional algebra A . For each $b \in \mathbf{B}$, let $f_b \in B = \text{End}_B(T)^{\text{op}}$ be an idempotent such that $Tf_b \cong T_\varepsilon(b)$. Then $P'(b) := Bf_b$ is an indecomposable projective B -module and the modules

$$\{L'(b) := \text{hd } P'(b) \mid b \in \mathbf{B}\}$$

give a full set of pairwise inequivalent irreducible left B -modules. Since \mathcal{R}' is a finite Abelian category, it is immediate that $(\mathbf{B}, L', \rho, \Lambda, \geq)$ is a stratification of it. Let $\Delta'_{-\varepsilon}(b)$ and $\nabla'_{-\varepsilon}(b)$ be the $(-\varepsilon)$ -standard and $(-\varepsilon)$ -costandard objects of \mathcal{R}' defined from this stratification. Set $V(b) := F\nabla_\varepsilon(b)$.

Step 1: For $b \in \mathbf{B}$ we have that $P'(b) \cong FT_\varepsilon(b)$. This follows immediately from the equality $\text{Hom}_A(T, T)f_b = \text{Hom}_A(T, Tf_b)$.

Step 2: The functor F sends short exact sequences of objects in $\nabla_\varepsilon(\mathcal{R})$ to short exact sequences in \mathcal{R}' . This follows because $\text{Ext}_{\mathcal{R}}^1(T, V) = 0$ for $V \in \nabla_\varepsilon(\mathcal{R})$ by the usual Ext^1 -vanishing between Δ_ε - and ∇_ε -filtered objects.

Step 3: For $a, b \in \mathbf{B}$, we have that $[V(b) : L'(a)] = (T_\varepsilon(a) : \Delta_\varepsilon(b))$. The left hand side is $\dim f_a V(b) = \dim f_a \text{Hom}_A(T, \nabla_\varepsilon(b)) \cong \dim \text{Hom}_A(T_\varepsilon(a), \nabla_\varepsilon(b))$, which equals the right hand side.

Step 4: $V(b)$ is a non-zero quotient of $P'(b)$, thus, $\text{hd } V(b) = L'(b)$. By Theorem 4.2(i), there is a short exact sequence $0 \rightarrow K \rightarrow T_\varepsilon(b) \rightarrow \nabla_\varepsilon(b) \rightarrow 0$ with $K \in \nabla_\varepsilon(\mathcal{R})$. Hence, Step 2 implies that $V(b)$ is quotient of $P'(b)$. It is non-zero by Step 3.

Step 5: We have that $V(b) \cong \Delta'_{-\varepsilon}(b)$. Let $\lambda := \rho(b)$. We treat the cases $\varepsilon(\lambda) = +$ and $\varepsilon(\lambda) = -$ separately. If $\varepsilon(\lambda) = +$ we must show that $V(b)$ is the largest quotient of $P'(b)$ with the property that $[V(b) : L'(a)] \neq 0 \Rightarrow \rho(a) \geq \rho(b)$. We have already observed in Step 4 that $V(b)$ is a quotient of $P'(b)$. Also $(T_\varepsilon(a) : \Delta_\varepsilon(b)) \neq 0 \Rightarrow \rho(b) \leq \rho(a)$ by Theorem 4.2(iii). Using Step 3, this implies that $V(b)$ has the property $[V(b) : L'(a)] \neq 0 \Rightarrow \rho(a) \geq \rho(b)$. It remains to show that any strictly larger quotient of $P'(b)$ fails this condition. To see this, since $\varepsilon(\lambda) = +$, a ∇_ε -flag in $T_\varepsilon(b)$ has $\nabla_\varepsilon(b)$ at the top and other sections $\nabla_\varepsilon(c)$ for c with $\rho(c) < \rho(b)$. In view of Step 4, any strictly larger quotient of $P'(b)$ than $V(b)$ therefore has an additional composition factor $L'(c)$ arising from the head of $V(c)$ for some c with $\rho(c) < \rho(b)$.

Instead, if $\varepsilon(\lambda) = -$, we use the characterization of $\Delta'_{-\varepsilon}(b)$ from Lemma 3.1(1): we must show that $V(b)$ is the largest quotient of $P'(b)$ with the property that $[\text{rad } V(b) : L'(a)] \neq 0 \Rightarrow \rho(a) > \rho(b)$. Since $\varepsilon(\lambda) = -$, we have that $(T_\varepsilon(b) : \nabla_\varepsilon(b)) = 1$ and $(T_\varepsilon(b) : \nabla_\varepsilon(a)) \neq 0 \Rightarrow \rho(a) < \rho(b)$ for $a \neq b$. Hence, using Step 3 again, the quotient $V(b)$ of $P'(b)$ has the required properties. A ∇_ε -flag in $T_\varepsilon(b)$ has $\nabla_\varepsilon(b)$ at the top and other sections $\nabla_\varepsilon(c)$ for c with $\rho(c) \leq \rho(b)$. So any strictly larger quotient of $P'(b)$ than $V(b)$ has a composition factor $L'(c)$ arising from the head of $V(c)$ for c with $\rho(c) \leq \rho(b)$. In case $c = b$, this violates the requirement that the quotient has $L'(b)$ appearing with multiplicity one; otherwise, it violates the requirement that all other composition factors of the quotient are of the form $L'(a)$ with $\rho(a) > \rho(b)$.

Step 6: \mathcal{R}' is a finite $(-\varepsilon)$ -stratified category. In view of Step 5, it suffices to show that $P'(b)$ has a filtration with sections $V(c)$ for c with $\rho(c) \leq \rho(b)$. Since $T_\varepsilon(b)$ has a ∇_ε -flag with sections $\nabla_\varepsilon(c)$ for c with $\rho(c) \leq \rho(b)$, this follows using Steps 1 and 2.

Step 7: For any $U \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$ and $V \in \mathcal{R}$, the map $f : \text{Hom}_A(U, V) \rightarrow \text{Hom}_B(FU, FV)$ induced by F is an isomorphism. It suffices to prove this when $U = T$, so that the right hand space is $\text{Hom}_B(B, FV)$ and $FV = \text{Hom}_A(T, V)$. This special case follows because f is the inverse of the isomorphism $\text{Hom}_B(B, FV) \rightarrow FV, \theta \mapsto \theta(1)$.

Step 8: For any $V, W \in \nabla_\varepsilon(\mathcal{R})$ and $n \geq 0$, the functor F induces a linear isomorphism $\text{Ext}_{\mathcal{R}}^n(V, W) \xrightarrow{\sim} \text{Ext}_{\mathcal{R}'}^n(FV, FW)$. Take an ε -tilting resolution $d : T_\bullet \rightarrow V$ in the sense of Definition 4.6, which exists thanks to Theorem 4.8. The functor F takes this resolution to a complex

$$\cdots \longrightarrow FT_1 \longrightarrow FT_0 \longrightarrow FV \longrightarrow 0.$$

In fact, this complex is exact. To see this, take $m \geq 0$ and consider the short exact sequence $0 \rightarrow \ker d_m \rightarrow T_m \rightarrow \text{im } d_m \rightarrow 0$. All of $\ker d_m$, T_m and $\text{im } d_m$ have ∇_ε -flags due to Lemma 4.7(1). Hence, thanks to Step 2, we get a short exact sequence

$$0 \longrightarrow F(\ker d_m) \xrightarrow{i} FT_m \xrightarrow{p} F(\text{im } d_m) \longrightarrow 0$$

on applying F . Since F is left exact, the canonical map $F(\text{im } d_m) \rightarrow FT_{m-1}$ is a monomorphism. Its image is all $\theta : T \rightarrow T_{m-1}$ with image contained in $\text{im } d_m$. As p is an epimorphism, any such θ can be written as $d_m \circ \phi$ for $\phi : T \rightarrow T_m$, i.e., $\theta \in \text{im}(Fd_m)$. Thus, $F(\text{im } d_m) \cong \text{im}(Fd_m)$, and $0 \rightarrow \ker(Fd_m) \rightarrow FT_m \rightarrow \text{im}(Fd_m) \rightarrow 0$ is exact, as required. In view of Step 1, we have constructed a projective resolution of FV in \mathcal{R}' :

$$\cdots \longrightarrow FT_1 \longrightarrow FT_0 \longrightarrow FV \longrightarrow 0.$$

Next, we use the projective resolution just constructed to compute $\text{Ext}_{\mathcal{R}'}^n(FV, FI)$ for any injective $I \in \mathcal{R}$. We have a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}_{\mathcal{R}}(V, I) & \longrightarrow & \text{Hom}_{\mathcal{R}}(T_0, I) & \longrightarrow & \text{Hom}_{\mathcal{R}}(T_1, I) \longrightarrow \cdots \\ & & \downarrow f & & \downarrow f_0 & & \downarrow f_1 \\ 0 & \longrightarrow & \text{Hom}_{\mathcal{R}'}(FV, FI) & \longrightarrow & \text{Hom}_{\mathcal{R}'}(FT_0, FI) & \longrightarrow & \text{Hom}_{\mathcal{R}'}(FT_1, FI) \longrightarrow \cdots \end{array}$$

with vertical maps induced by F . The maps f_0, f_1, \dots are isomorphisms due to Step 7. Also the top row is exact as I is injective. We deduce that the bottom row is exact at the positions $\text{Hom}_{\mathcal{R}'}(FT_m, FI)$ for all $m \geq 1$. It is exact at positions $\text{Hom}_{\mathcal{R}'}(FV, FI)$ and $\text{Hom}_{\mathcal{R}'}(FT_0, FI)$ as $\text{Hom}_{\mathcal{R}'}(?, FI)$ is left exact. Thus, the bottom row is exact everywhere. So the map f is an isomorphism too and $\text{Ext}_{\mathcal{R}'}^n(FV, FI) = 0$ for $n > 0$.

Finally, take a short exact sequence $0 \rightarrow W \rightarrow I \rightarrow Q \rightarrow 0$ in \mathcal{R} with I injective. We have that $Q \in \nabla_\varepsilon(\mathcal{R})$ by Corollary 3.13. Hence, using Step 2 and the previous paragraph, there is a commutative diagram

$$\begin{array}{ccccccc} \text{Hom}_{\mathcal{R}}(V, W) & \longleftarrow & \text{Hom}_{\mathcal{R}}(V, I) & \longrightarrow & \text{Hom}_{\mathcal{R}}(V, Q) & \longrightarrow & \text{Ext}_{\mathcal{R}}^1(V, W) \\ \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 & & \downarrow f_4 \\ \text{Hom}_{\mathcal{R}'}(FV, FW) & \longleftarrow & \text{Hom}_{\mathcal{R}'}(FV, FI) & \longrightarrow & \text{Hom}_{\mathcal{R}'}(FV, FQ) & \longrightarrow & \text{Ext}_{\mathcal{R}'}^1(FV, FW) \end{array}$$

with exact rows. As f_2 is an isomorphism, we get that f_1 is injective. Since this is proved for all W , this means that f_3 is injective too. Then a diagram chase gives that f_1 is surjective, hence, f_3 is surjective and f_4 is an isomorphism. Degree shifting now gives the isomorphisms $\text{Ext}_{\mathcal{R}}^n(V, W) \xrightarrow{\sim} \text{Ext}_{\mathcal{R}'}^n(FV, FW)$ for $n \geq 2$ as well.

Step 9: We have that $T'_\varepsilon(b) \cong FI(b)$. By Steps 5 and 8, we get that

$$\text{Ext}_{\mathcal{R}'}^1(\Delta'_{-\varepsilon}(a), FI(b)) \cong \text{Ext}_{\mathcal{R}}^1(\nabla_\varepsilon(a), I(b)) = 0$$

for all $a \in \mathbf{B}$. Hence, by the homological criterion for $\nabla_{-\varepsilon}$ -flags in the $(-\varepsilon)$ -stratified category \mathcal{R}' , the A -module $FI(b)$ has a $\nabla_{-\varepsilon}$ -flag. It also has a $\Delta_{-\varepsilon}$ -flag with bottom section isomorphism to $\Delta'_{-\varepsilon}(b)$ due to Steps 2 and 5. So $FI(b) \in \mathcal{Tilt}_{-\varepsilon}(\mathcal{R}')$. It is indecomposable as $\text{End}_{\mathcal{R}'}(FI(b)) \cong \text{End}_{\mathcal{R}}(I(b))$ by Step 8, which is local. Therefore $FI(b) \cong T'_{-\varepsilon}(b)$ due to Theorem 4.2(i).

Step 10: *The restriction $F : \nabla_{\varepsilon}(\mathcal{R}) \rightarrow \Delta_{-\varepsilon}(\mathcal{R}')$ is an equivalence of categories.* It is full and faithful by Step 8. It remains to show that it is dense, i.e., for any $V' \in \Delta_{-\varepsilon}(\mathcal{R}')$ there exists $V \in \nabla_{\varepsilon}(\mathcal{R})$ with $FV \cong V'$. The proof of this goes by induction on the length of a $\Delta_{-\varepsilon}$ -flag of V' . If this length is one, we are done by Step 5. For the induction step, consider V' fitting into a short exact sequence $0 \rightarrow U' \rightarrow V' \rightarrow W' \rightarrow 0$ for shorter $U', W' \in \Delta_{-\varepsilon}(\mathcal{R}')$. By induction there are $U, W \in \nabla_{\varepsilon}(\mathcal{R})$ such that $FU \cong U'$ and $FW \cong W'$. Then we use the isomorphism $\text{Ext}_{\mathcal{R}'}^1(FW, FU) \cong \text{Ext}_{\mathcal{R}}^1(W, U)$ from Step 8 to see that there is an extension V of U and W in \mathcal{R} such that $FV \cong V'$.

Step 11: *The dual right A -module T^* to T is a $(-\varepsilon)$ -tilting generator for $\mathcal{R}^{\text{op}} = \text{mod}_{\text{fd}} A$ such that $\text{End}_A(T^*)^{\text{op}} = B^{\text{op}}$. Moreover, letting $F^{\text{op}} := \text{Hom}_A(T^*, ?) : \text{mod}_{\text{fd}} A \rightarrow \text{mod}_{\text{fd}} B$ be the corresponding Ringel duality functor, we have that $G \cong ?^* \circ F^{\text{op}} \circ ?^*$. The first statement is clear from Theorem 3.9, observing that $\text{End}_A(T^*)^{\text{op}} \cong \text{End}_A(T)$ since $* : A\text{-mod}_{\text{fd}} \rightarrow \text{mod}_{\text{fd}} A$ is a contravariant equivalence. It remains to observe that $* \circ F^{\text{op}} \circ * \cong \text{Hom}_A(T^*, ?^*)^* \cong \text{Hom}_A(?, T)^* = G$.*

Step 12: *The restriction $G : \Delta_{\varepsilon}(\mathcal{R}) \rightarrow \nabla_{-\varepsilon}(\mathcal{R}')$ is an equivalence of categories inducing isomorphisms as in (4.6), such that $GT_{\varepsilon}(b) \cong I'(b)$, $G\Delta_{\varepsilon}(b) \cong \nabla'_{-\varepsilon}(b)$ and $GP(b) \cong T'_{-\varepsilon}(b)$.* This follows using Step 11 and the analogs for F^{op} of the statements about F established thus far. \square

4.3. Tilting objects in the upper finite and essentially finite cases. Throughout the subsection, \mathcal{R} will be either be an upper finite or an essentially finite ε -stratified category with the usual stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. It is still possible to make sense of ε -tilting objects but now the iterative procedure used to construct the indecomposable ones in the proof of Theorem 4.2 does not terminate after finitely many steps. Consequently, we must allow for tilting objects which have infinite Δ_{ε} - and ∇_{ε} -flags; see (6.6) below for a baby example of this phenomenon.

Suppose to start with that \mathcal{R} is an upper finite ε -stratified category. Using the notions of ascending Δ_{ε} -flags and descending ∇_{ε} -flags introduced in Definition 3.35, we set

$$\mathcal{Tilt}_{\varepsilon}(\mathcal{R}) := \Delta_{\varepsilon}^{\text{asc}}(\mathcal{R}) \cap \nabla_{\varepsilon}^{\text{dsc}}(\mathcal{R}). \quad (4.10)$$

We emphasize that objects of $\mathcal{Tilt}_{\varepsilon}(\mathcal{R})$ are in particular objects of \mathcal{R} , so all of their composition multiplicities are finite. Like in Lemma 4.1, $\mathcal{Tilt}_{\varepsilon}(\mathcal{R})$ is an additive Karoubian subcategory of \mathcal{R} .

Theorem 4.18 (Classification of tilting objects in the upper finite case). *Assume that \mathcal{R} is an upper finite ε -stratified category. For $b \in \mathbf{B}_{\lambda}$, there is an indecomposable object $T_{\varepsilon}(b) \in \mathcal{Tilt}_{\varepsilon}(\mathcal{R})$ satisfying the following properties:*

- (i) $T_{\varepsilon}(b)$ has an ascending Δ_{ε} -flag with bottom section⁶ isomorphic to $\Delta_{\varepsilon}(b)$;
- (ii) $T_{\varepsilon}(b)$ has a descending ∇_{ε} -flag with top section⁷ isomorphic to $\nabla_{\varepsilon}(b)$;
- (iii) $T_{\varepsilon}(b) \in \mathcal{R}_{\leq \lambda}$ and $j^{\lambda}T_{\varepsilon}(b) \cong \begin{cases} P_{\lambda}(b) & \text{if } \varepsilon(\lambda) = + \\ I_{\lambda}(b) & \text{if } \varepsilon(\lambda) = - \end{cases}$.

These properties determine $T_{\varepsilon}(b)$ uniquely up to isomorphism: if T is any indecomposable object of $\mathcal{Tilt}_{\varepsilon}(\mathcal{R})$ satisfying any one of the properties (i)–(iii) then $T \cong T_{\varepsilon}(b)$; hence, it satisfies the other two properties as well.

⁶We mean that there is an ascending Δ_{ε} -flag $(V_{\omega})_{\omega \in \Omega}$ in which Ω has a smallest non-zero element 1 such that $V_1 \cong \Delta_{\varepsilon}(b)$.

⁷Similarly, we mean that $V/V_1 \cong \nabla_{\varepsilon}(b)$.

Proof. Existence: Replacing \mathcal{R} by $\mathcal{R}_{\leq \lambda}$ if necessary and using Theorem 3.41, we reduce to the special case that λ is the largest element of the poset Λ . Assuming this, the first step in the construction of $T_\varepsilon(b)$ is to define a direct system $(V_\omega)_{\omega \in \Omega}$ of objects of \mathcal{R} . This is indexed by the directed set Ω of all finite upper sets in Λ . Let $V_\emptyset := 0$. Then take $\emptyset \neq \omega \in \Omega$ and denote it instead by Λ^\dagger . Letting $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be the corresponding finite ε -stratified quotient of \mathcal{R} , we set $V_\omega := j_! T_\varepsilon^\dagger(b)$. By Theorem 3.42(6), this has a Δ_ε -flag. Given also $\omega < \nu \in \Omega$, i.e., another upper set $\Lambda^{\dagger\dagger}$ containing Λ^\dagger , let $k : \mathcal{R} \rightarrow \mathcal{R}^{\dagger\dagger}$ be the corresponding quotient. Then j factors as $j = \bar{j} \circ k$ for an induced quotient functor $\bar{j} : \mathcal{R}^\dagger \rightarrow \mathcal{R}^{\dagger\dagger}$. Since $\bar{j} T_\varepsilon^{\dagger\dagger}(b) \cong T_\varepsilon^\dagger(b)$ by Corollary 4.5, we deduce from Corollary 3.19(2) that there is a short exact sequence

$$0 \longrightarrow \bar{j}_! T_\varepsilon^\dagger(b) \longrightarrow T_\varepsilon^{\dagger\dagger}(b) \longrightarrow Q \longrightarrow 0$$

such that Q has a Δ_ε -flag with sections $\Delta_\varepsilon^{\dagger\dagger}(c)$ for c with $\rho(c) \in \Lambda^{\dagger\dagger} \setminus \Lambda^\dagger$. Applying $k_!$ and using the exactness from Theorem 3.42(6) again, we deduce that there is an embedding $f_\omega^v : V_\omega \hookrightarrow V_\nu$ with $\text{coker } f_\omega^v \in \Delta_\varepsilon(\mathcal{R})$. Thus, we have a direct system $(V_\omega)_{\omega \in \Omega}$. Now let $T_\varepsilon(b) := \varinjlim V_\omega \in \text{Ind}(\mathcal{R}_c)$. Using the induced embeddings $f_\omega : V_\omega \hookrightarrow T_\varepsilon(b)$, we identify each V_ω with a subobject of $T_\varepsilon(b)$. We have shown for $\omega < \nu$ that $V_\nu/V_\omega \in \Delta_\varepsilon(\mathcal{R})$ and, moreover, $jV_\nu = jV_\omega$ where $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ is the quotient associated to ω .

In this paragraph, we show that $T_\varepsilon(b)$ actually lies in \mathcal{R} rather than $\text{Ind}(\mathcal{R}_c)$, i.e., all of the composition multiplicities $[T_\varepsilon(b) : L(c)]$ are finite. To see this, take $c \in \mathbf{B}$. Let $\omega = \Lambda^\dagger \in \Omega$ be some fixed finite upper set such that $\rho(c) \in \Lambda^\dagger$, and $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be the quotient functor as usual. Then for any $\nu \geq \omega$ we have that

$$[V_\nu : L(c)] = [jV_\nu : L^\dagger(c)] = [jV_\omega : L^\dagger(c)] = [V_\omega : L(c)].$$

Hence, $[T_\varepsilon(b) : L(c)] = [V_\omega : L(c)] < \infty$.

So now we have defined $T_\varepsilon(b) \in \mathcal{R}$ together with an ascending Δ_ε -flag $(V_\omega)_{\omega \in \Omega}$. The smallest non-empty element of Ω is $\omega := \{\lambda\}$, and $V_\omega = j_!^\lambda P_\lambda(b) = \Delta_\varepsilon(b)$ if $\varepsilon(\lambda) = +$, or $j_!^\lambda I_\lambda(b)$ if $\varepsilon(\lambda) = -$. Since $j^\lambda T_\varepsilon(b) = j^\lambda V_\omega$, we deduce that (iii) holds. Also by construction $T_\varepsilon(b)$ has an ascending Δ_ε -flag. To see that it has a descending ∇_ε -flag, take any $a \in \mathbf{B}$. Let $\omega = \Lambda^\dagger \in \Omega$ be such that $\rho(a) \in \Lambda^\dagger$. Then $\Delta_\varepsilon(a) = j_! \Delta_\varepsilon^\dagger(a)$ and $jT_\varepsilon(b) = jV_\omega = T_\varepsilon^\dagger(b)$, so by Theorem 3.42(5) we get that

$$\text{Ext}_{\mathcal{R}}^1(\Delta_\varepsilon(a), T_\varepsilon(b)) \cong \text{Ext}_{\mathcal{R}^\dagger}^1(\Delta_\varepsilon^\dagger(a), T_\varepsilon^\dagger(b)) = 0.$$

By Theorem 3.39, this shows that $T_\varepsilon(b) \in \nabla_\varepsilon^{\text{dsc}}(\mathcal{R})$.

Note finally that $T_\varepsilon(b)$ is indecomposable. This follows because $jT_\varepsilon(b)$ is indecomposable for every $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ (adopting the usual notation). Indeed, by the construction we have that $jT_\varepsilon(b) \cong T_\varepsilon^\dagger(b)$. This completes the construction of the indecomposable object $T_\varepsilon(b) \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$. We have shown that it satisfies (iii), and it follows easily that it also satisfies (i) and (ii).

Uniqueness: Since (iii) implies (i) and (ii), it suffices to show that any indecomposable $U \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$ satisfying either (i) or (ii) is isomorphic to the object $T := T_\varepsilon(b)$ just constructed. We explain this just in the case of (i), since the argument for (ii) is similar. We take a short exact sequence $0 \rightarrow \Delta_\varepsilon(b) \rightarrow T \rightarrow Q \rightarrow 0$ with $Q \in \Delta_\varepsilon^{\text{asc}}(\mathcal{R})$. Using the Ext-vanishing from Lemma 3.36, we deduce like in the proof of Theorem 4.2 that the inclusion $f : \Delta_\varepsilon(b) \hookrightarrow T$ extends to $\bar{f} : U \rightarrow T$. In fact, \bar{f} is an isomorphism. To see this, take a finite upper set Λ^\dagger containing λ and consider the quotient $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ as usual. Both jU and jT are isomorphic to $T_\varepsilon^\dagger(b)$ by the uniqueness in Theorem 4.2. The proof there implies that any homomorphism $jT \rightarrow jU$ which restricts to an isomorphism on the subobject $\Delta_\varepsilon^\dagger(b)$ is an isomorphism. We deduce that $j\bar{f}$ is an isomorphism. Since this holds for all choices of Λ^\dagger , it follows that \bar{f} itself is an isomorphism. \square

Corollary 4.19. *Any object of $\mathcal{Tilt}_\varepsilon(\mathcal{R})$ is isomorphic to $\bigoplus_{b \in \mathbf{B}} T_\varepsilon(b)^{\oplus n_b}$ for unique multiplicities $n_b \in \mathbb{N}$. Conversely, any such direct sum belongs to $\mathcal{Tilt}_\varepsilon(\mathcal{R})$.*

Proof. Let us first show that any direct sum $U := \bigoplus_{b \in \mathbf{B}} T_\varepsilon(b)^{\oplus n_b}$ belongs to $\mathcal{Tilt}_\varepsilon(\mathcal{R})$. The only issue is to see that U actually belongs to \mathcal{R} rather than $\text{Ind}(\mathcal{R}_c)$, i.e., it has finite composition multiplicities. But for a given $c \in \mathbf{B}$, the multiplicity $[T_\varepsilon(b) : L(c)]$ is zero unless $\rho(c) \leq \rho(b)$. There are only finitely many such $b \in \mathbf{B}$, so $[U : L(c)] = \sum_{b \in \mathbf{B}} n_b [T_\varepsilon(b) : L(c)] < \infty$.

Now take any $U \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$. Let Ω be the directed set of all finite upper sets in Λ . Take $\omega = \Lambda^\dagger \in \Omega$. Let $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be the quotient functor as usual. Then we have that $jU \in \mathcal{Tilt}_\varepsilon(\mathcal{R}^\dagger)$, so it decomposes as a finite direct sum as $jU \cong \bigoplus_{b \in \mathbf{B}^\dagger} T_\varepsilon^\dagger(b)^{\oplus n_b(\omega)}$ for $n_b(\omega) \in \mathbb{N}$. There is a corresponding direct summand $T_\omega \cong \bigoplus_{b \in \mathbf{B}^\dagger} T_\varepsilon(b)^{\oplus n_b(\omega)}$ of U . Then $T = \varinjlim T_\omega$. Moreover, for $b \in \mathbf{B}^\dagger$, the multiplicities $n_b(\omega)$ are stable in the sense that $n_b(v) = n_b(\omega)$ for all $v > \omega$. We deduce that $U \cong \bigoplus_{b \in \mathbf{B}} T_\varepsilon(b)^{\oplus n_b}$ where $n_b := n_b(\omega)$ for any sufficiently large ω . \square

It remains to discuss tilting objects in the essentially finite case. So now we assume that \mathcal{R} is an essentially finite ε -stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Since Λ is interval finite, finite unions of lower sets of the form $(-\infty, \lambda]$ are upper finite. If \mathcal{R}^\dagger is the Serre subcategory of \mathcal{R} associated to such an upper finite lower set then its Schurian envelope $\text{Env}(\mathcal{R}^\dagger)$ in the sense of Lemma 2.22 is a Cartan-bounded upper finite ε -stratified category which is naturally embedded into $\text{Env}(\mathcal{R})$. This follows from Theorem 3.17. For $b \in \mathbf{B}$, we define the corresponding ε -tilting object $T_\varepsilon(b) \in \text{Env}(\mathcal{R})$ as follows: pick any upper finite lower set Λ^\dagger such that $\rho(b) \in \Lambda^\dagger$, let \mathcal{R}^\dagger be the corresponding Serre subcategory of \mathcal{R} , then let $T_\varepsilon(b)$ be the ε -tilting object in $\text{Env}(\mathcal{R}^\dagger)$ from Theorem 4.18. This is well-defined independent of the choice of Λ^\dagger by the uniqueness part of Theorem 4.18. Thus, we have defined the indecomposable ε -tilting objects $\{T_\varepsilon(b) \mid b \in \mathbf{B}\}$ in the essentially finite case too, although these may be of infinite length, i.e., in general they belong to $\text{Env}(\mathcal{R})$ rather than to \mathcal{R} itself.

Definition 4.20. Suppose that \mathcal{R} is a lower finite, upper finite or essentially finite ε -stratified category with the usual stratification. We say that it is *tilting-bounded* if the matrix

$$(\dim \text{Hom}_{\mathcal{R}}(T_\varepsilon(a), T_\varepsilon(b)))_{a, b \in \mathbf{B}} \quad (4.11)$$

has finitely many non-zero entries in each row and each column.

The matrix (4.11) is analogous to the Cartan matrix (2.23) with projectives/injectives replaced by ε -tilting objects. In the lower finite case, all entries of this matrix are obviously $< \infty$, but in the upper finite or essentially finite cases it is possible that some of these dimensions are ∞ . However they are all finite in the tilting-bounded case:

Lemma 4.21. *If \mathcal{R} is tilting-bounded then the spaces $\text{Hom}_{\mathcal{R}}(T_\varepsilon(a), T_\varepsilon(b))$ are finite-dimensional for all $a, b \in \mathbf{B}$.*

Proof. In the lower finite case, the indecomposable tilting objects are of finite length, so these spaces are finite-dimensional even without the assumption that \mathcal{R} is tilting-bounded. In the remaining upper finite or essentially finite cases, we have that

$$\dim \text{Hom}_{\mathcal{R}}(T_\varepsilon(a), T_\varepsilon(b)) = \sum_{c \in \mathbf{B}} (T_\varepsilon(a) : \nabla_\varepsilon(c)) (T_\varepsilon(b) : \Delta_\varepsilon(c)) \in \mathbb{N} \cup \{\infty\}. \quad (4.12)$$

All of the multiplicities $(T_\varepsilon(a) : \nabla_\varepsilon(c))$ and $(T_\varepsilon(b) : \Delta_\varepsilon(c))$ are finite. Moreover, if $(T_\varepsilon(a) : \Delta_\varepsilon(c)) \neq 0$ then $\text{Hom}_{\mathcal{R}}(T_\varepsilon(a), T_\varepsilon(c)) \neq 0$. Hence, assuming the tilting-bounded hypothesis, only finitely many of the terms in the sum on the right hand side are non-zero. \square

Assuming \mathcal{R} is an essentially finite ε -stratified category once again, assume that \mathcal{R} is also tilting-bounded. Then the ε -tilting objects $T_\varepsilon(b)$ actually belong to

$$\mathcal{Tilt}_\varepsilon(\mathcal{R}) := \Delta_\varepsilon(\mathcal{R}) \cap \nabla_\varepsilon(\mathcal{R}), \quad (4.13)$$

i.e., they belong to \mathcal{R} rather than to $\text{Env}(\mathcal{R})$ of \mathcal{R} . Thus, we are in a similar situation to (4.1). Theorem 4.2 carries over easily, to show that $\{T_\varepsilon(b) \mid b \in \mathbf{B}\}$ gives a full set of the indecomposable objects in the additive Karoubian category $\mathcal{Tilt}_\varepsilon(\mathcal{R})$. The construction of Theorem 4.8 also carries over unchanged. So all objects of $\nabla_\varepsilon(\mathcal{R})$ have ε -tilting resolutions and all objects of $\Delta_\varepsilon(\mathcal{R})$ have ε -cotilting resolutions.

Remark 4.22. Most of the interesting examples of essentially finite highest weight categories which arise “in nature” seem to satisfy the tilting-bounded hypothesis, although there is no reason for this to be the case from the recursive construction of Theorem 4.18. We refer the reader to Remark 6.2 for an example which is not tilting-bounded.

Remark 4.23. The tilting-bounded hypothesis is also interesting in the lower finite case; see Corollary 4.28 below. Using (4.12), it is easy to see in the lower finite case that \mathcal{R} is tilting-bounded if and only if for each $b \in \mathbf{B}$ the multiplicities $(T_\varepsilon(a) : \Delta_\varepsilon(b))$ and $(T_\varepsilon(a) : \nabla_\varepsilon(b))$ are zero for all but finitely many $a \in \mathbf{B}$. Natural examples of lower finite highest weight categories which are definitely *not* tilting-bounded include the categories $\mathcal{R}ep(G)$ for reductive groups G (unless this is actually a semisimple category), as follows from the results in [Cou1, §5]. In situations involving quantum groups at roots of unity, tilting-boundedness can be checked combinatorially by considering properties of Kazhdan-Lusztig polynomials; e.g., see [Soe], [Str].

4.4. Semi-infinite Ringel duality. Now we extend Ringel duality to lower finite and upper finite ε -stratified categories. The situation is not as symmetric as in the finite case and demands different constructions when going from lower finite to upper finite or from upper finite to lower finite. If we start with a lower finite ε -stratified category, the Ringel dual is an upper finite $(-\varepsilon)$ -stratified category:

Definition 4.24. Let \mathcal{R} be a lower finite ε -stratified category with the usual stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. An ε -tilting generator for \mathcal{R} is an object $T = \bigoplus_{i \in I} T_i \in \text{Ind}(\mathcal{R})$ with a given decomposition as a direct sum of objects $T_i \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$ such that each $T_\varepsilon(b)$ is isomorphic to a summand of T . Define the *Ringel dual* of \mathcal{R} relative to $T = \bigoplus_{i \in I} T_i$ to be the Schurian category $\mathcal{R}' := A\text{-mod}_{\text{fd}}$ where

$$A := \left(\bigoplus_{i, j \in I} \text{Hom}_{\mathcal{R}}(T_i, T_j) \right)^{\text{op}}.$$

Identifying $\text{Ind}(\mathcal{R}'_c)$ with $A\text{-mod}$ as explained in (2.3), we have the *Ringel duality functor*

$$F := \bigoplus_{i \in I} \text{Hom}_{\mathcal{R}}(T_i, ?) : \text{Ind}(\mathcal{R}) \rightarrow \text{Ind}(\mathcal{R}'_c). \quad (4.14)$$

This functor takes objects of \mathcal{R} to objects of \mathcal{R}' .

Theorem 4.25 (Lower to upper semi-infinite Ringel duality). *In the setup of Definition 4.24, \mathcal{R}' is an upper finite $(-\varepsilon)$ -stratified category with stratification $(\mathbf{B}, L', \rho, \Lambda, \geq)$ and distinguished objects*

$$\begin{aligned} P'(b) &\cong FT_\varepsilon(b), & L'(b) &\cong \text{hd } P'(b), \\ \Delta'_{-\varepsilon}(b) &\cong F\nabla_\varepsilon(b), & T'_{-\varepsilon}(b) &\cong FI(b). \end{aligned}$$

The restriction $F : \nabla_\varepsilon^{\text{asc}}(\mathcal{R}) \rightarrow \Delta'_{-\varepsilon}(\mathcal{R}')$ is an equivalence of categories.

The proof will be explained later in the subsection.

In the other direction, if we start from an upper finite ε -stratified category, the Ringel dual is a lower finite $(-\varepsilon)$ -stratified category:

Definition 4.26. Let \mathcal{R} be an upper finite ε -stratified category with the usual stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. An ε -tilting generator is an object $T \in \mathcal{Tilt}_\varepsilon(\mathcal{R})$ such that $T_\varepsilon(b)$ is isomorphic to a summand of T for every $b \in \mathbf{B}$. Let $C := \text{Coend}_{\mathcal{R}}(T)$ be the coalgebra that is the continuous dual of the pseudo-compact topological algebra

$B := \text{End}_{\mathcal{R}}(T)^{\text{op}}$; see Lemma 2.10. Then the *Ringel dual* of \mathcal{R} relative to T is the category $\mathcal{R}' := \text{comod}_{\text{fd}}\text{-}C = B\text{-mod}_{\text{fd}}$. Recalling Lemma 2.11, the *Ringel duality functor* is

$$G := \text{Cohom}_{\mathcal{R}}(T, ?) = \text{Hom}_{\mathcal{R}}(?, T)^* : \text{Ind}(\mathcal{R}_c) \rightarrow \text{Ind}(\mathcal{R}'), \quad (4.15)$$

which sends finitely generated objects of \mathcal{R} to objects of \mathcal{R}' .

Theorem 4.27 (Upper to lower semi-infinite Ringel duality). *In the setup of Definition 4.26, \mathcal{R}' is a lower finite $(-\varepsilon)$ -stratified category with stratification $(\mathbf{B}, L', \rho, \Lambda, \geq)$ and distinguished objects*

$$\begin{aligned} I'(b) &= GT_{\varepsilon}(b), & L'(b) &= \text{soc } I'(b), \\ \nabla'_{-\varepsilon}(b) &= G\Delta_{\varepsilon}(b), & T'_{-\varepsilon}(b) &= GP(b). \end{aligned}$$

The restriction $G : \Delta_{\varepsilon}^{\text{asc}}(\mathcal{R}) \rightarrow \nabla_{-\varepsilon}^{\text{asc}}(\mathcal{R}')$ is an equivalence of categories.

Again the proof will be explained later.

We proceed to formulate several consequences of Theorems 4.25 and 4.27. The first is concerned with a special case. Recall the definition of Cartan-bounded from just before Lemma 2.22, and the definition of tilting-bounded from Definition 4.20.

Corollary 4.28. *The Ringel dual of a tilting-bounded lower finite ε -stratified category is a Cartan-bounded upper finite $(-\varepsilon)$ -stratified category. Conversely, the Ringel dual of a Cartan-bounded upper finite ε -stratified category is a tilting-bounded lower finite $(-\varepsilon)$ -stratified category.*

Proof. From either Theorem 4.25 or Theorem 4.27, it follows that the Cartan matrix (2.23) for the upper finite category is equal to the matrix (4.11) for the lower finite category. \square

The next two corollaries give the analogs of the double centralizer property from Corollary 4.11 in the semi-infinite setting.

Corollary 4.29 (Lower to upper double centralizer property). *Let notation be as in Definition 4.24. Assume in addition that $\mathcal{R} = \text{comod}_{\text{fd}}\text{-}C$ for a coalgebra C . Let $B := C^*$ be the dual algebra, so that T is a (B, A) -bimodule. Let $T' := T^{\otimes}$ be the dual (A, B) -bimodule.*

(1) *T' is a $(-\varepsilon)$ -tilting generator for \mathcal{R}' and there is an algebra isomorphism*

$$\mu : B \xrightarrow{\sim} \text{End}_{\mathcal{R}'}(T')^{\text{op}} \quad (4.16)$$

sending $y \in B$ to $\mu(y) : T' \rightarrow T', v \mapsto vy$. Equivalently, there is a coalgebra isomorphism

$$\mu^* : \text{Coend}_{\mathcal{R}'}(T') \xrightarrow{\sim} C, \quad c_{r,s}^{(i)} \mapsto \tilde{c}_{r,s}^{(i)} \quad (4.17)$$

where $c_{r,s}^{(i)}$ is the element of $\text{Coend}_{\mathcal{R}'}(T')$ corresponding to $v_s^{(i)} \otimes u_r^{(i)} \in T_i \otimes T_i^$ according to (2.13) for dual bases $v_1^{(i)}, \dots, v_{d(i)}^{(i)}$ for T_i and $u_1^{(i)}, \dots, u_{d(i)}^{(i)}$ for T_i^* , and $\tilde{c}_{r,s}^{(i)} \in C$ is defined so that the structure map of the right C -comodule T_i sends $v_s^{(i)} \mapsto \sum_{r=1}^{d(i)} v_r^{(i)} \otimes \tilde{c}_{r,s}^{(i)}$. So the Ringel dual of \mathcal{R}' relative to T' in the sense of Definition 4.26 is equivalent to the original category \mathcal{R} .*

(2) *Denote the Ringel duality functor for \mathcal{R}' relative to T' now by*

$$F^* := \text{Cohom}_{\mathcal{R}'}(T', ?) = \text{Hom}_{\mathcal{R}'}(?, T')^* : \text{Ind}(\mathcal{R}'_c) \rightarrow \text{Ind}(\mathcal{R}). \quad (4.18)$$

We have that $F^ \cong T \otimes_A ?$, hence, (F^*, F) is an adjoint pair; cf. Lemma 2.11.*

Proof. By Lemma 2.2, we have natural isomorphisms $\text{Hom}_C(T_i, C) \cong T_i^*$ as right B -modules, hence, $FC \cong T'$ as an (A, B) -bimodule. Since every $I(b)$ appears as a summand of the regular comodule, and $FI(b) \cong T'_{-\varepsilon}(b)$ by Theorem 4.25, we deduce that T' is a

$(-\varepsilon)$ -tilting generator for \mathcal{R}' . To see that $B \cong \text{End}_A(T')^{\text{op}}$, we use the fact that F is an equivalence on ∇ -filtered objects to deduce that

$$\text{End}_A(T')^{\text{op}} \cong \text{End}_A(FC)^{\text{op}} \cong \text{End}_C(C)^{\text{op}} \cong B,$$

using Lemma 2.2 again for the final algebra isomorphism. This produces the isomorphism μ . To deduce (4.17), we need to show that $\mu^*(c_{r,s}^{(i)})$ and $\tilde{c}_{r,s}^{(i)}$ take the same value on $y \in B$. The left hand side gives $c_{r,s}^{(i)}(\mu(y)) = v_s^{(i)}(u_s^{(i)}b)$. For the right hand side, we have that $yv_s^{(i)} = \sum_{r=1}^{d(i)} \tilde{c}_{r,s}^{(i)}(y)v_r^{(i)}$, so $c_{r,s}^{(i)}(y) = (yv_s^{(i)})u_r^{(i)}$. These are equal. This establishes (1). Then (2) follows from Lemma 2.11. \square

Corollary 4.30 (Upper to lower double centralizer property). *Let notation be as in Definition 4.26, and assume in addition that $\mathcal{R} = A\text{-mod}_{\text{fd}}$ for a locally finite-dimensional locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$. Let $T_i = e_i T$ and $T'_i := T_i^*$, so that $T' := \bigoplus_{i \in I} T'_i = T^{\otimes}$. This is a (B, A) -bimodule.*

(1) $T' = \bigoplus_{i \in I} T'_i$ is a $(-\varepsilon)$ -tilting generator for \mathcal{R}' and there is an algebra isomorphism

$$\mu : A \xrightarrow{\sim} \left(\bigoplus_{i,j \in I} \text{Hom}_{\mathcal{R}'}(T'_i, T'_j) \right)^{\text{op}} \quad (4.19)$$

sending $a \in e_i A e_j$ to $\mu(a) : T'_i \rightarrow T'_j, v \mapsto va$. So the Ringel dual of \mathcal{R}' relative to T' in the sense of Definition 4.24 is equivalent to the original category \mathcal{R} .

(2) Denote the Ringel duality functor for \mathcal{R}' relative to T' now by

$$G_* := \bigoplus_{i \in I} \text{Hom}_{\mathcal{R}'}(T'_i, ?) : \text{Ind}(\mathcal{R}') \rightarrow \text{Ind}(\mathcal{R}_c). \quad (4.20)$$

We have that $G \cong T' \otimes_A ?$, hence, (G, G_*) is an adjoint pair.

Proof. Note that $G(Ae_i) = \text{Hom}_A(Ae_i, T)^* = (e_i T)^* = T_i$. So Theorem 4.27 implies that $T = \bigoplus_{i \in I} T_i$ is a $(-\varepsilon)$ -tilting generator for \mathcal{R}' . Moreover,

$$\text{Hom}_{\mathcal{R}'}(T_i, T_j) = \text{Hom}_{\mathcal{R}'}(G(Ae_i), G(Ae_j)) \cong \text{Hom}_{\mathcal{R}}(Ae_i, Ae_j) = e_i Ae_j.$$

This proves (1) and then (2) follows from Lemma 2.11. \square

Remark 4.31. Combining Corollary 4.28 with the double centralizer properties just explained, one obtains a restricted version of semi-infinite Ringel duality giving a correspondence

$$\left\{ \begin{array}{l} \text{Tilting-bounded lower finite} \\ \text{highest weight categories} \end{array} \right\} \xleftrightarrow{\text{Ringel duality}} \left\{ \begin{array}{l} \text{Cartan-bounded upper finite} \\ \text{highest weight categories} \end{array} \right\}.$$

In the upper finite to lower finite direction, this appeared already in the work of Marko and Zubkov [MZ]. In more detail, if \mathcal{R} is the category of finite-dimensional modules over a descending quasi-hereditary pseudo-compact algebra in the sense of [MZ, Def. 3.19] and the indecomposable projectives in \mathcal{R} are of finite length as assumed in [MZ, §4], then \mathcal{R} is an essentially finite highest weight category with upper finite weight poset, hence, $\text{Env}(\mathcal{R})$ is a Cartan-bounded upper finite highest weight category. In this case, the indecomposable tilting modules $T(\lambda) \in \text{Env}(\mathcal{R})$ were constructed already in [MZ, §4], and the appropriate (lower finite) Ringel dual category appears in [MZ, §6]. Also [MZ, Lem. 6.5] establishes a double centralizer property which is equivalent to Corollary 4.30(1) for such categories.

In the setup of Definition 4.24, one can also define a functor

$$G := \text{Cohom}_{\mathcal{R}}(T, ?) = \text{Hom}_{\mathcal{R}}(?, T)^{\otimes} : \Delta_{\varepsilon}(\mathcal{R}) \rightarrow \nabla_{-\varepsilon}(\mathcal{R}'). \quad (4.21)$$

Theorem 4.25 plus an argument with duality like in Steps 11–12 of the proof of Theorem 4.10 shows that G is an equivalence of categories such that $G\Delta_{\varepsilon}(b) \cong \nabla'_{-\varepsilon}(b)$ and $GT_{\varepsilon}(b) \cong I'(b)$ for all $b \in \mathbf{B}$. Likewise, in the setup of Definition 4.26, one can also define

$$F := \text{Hom}_{\mathcal{R}}(T, ?) : \Delta_{\varepsilon}(\mathcal{R}) \rightarrow \nabla_{-\varepsilon}(\mathcal{R}'). \quad (4.22)$$

Theorem 4.27 plus an argument involving duality shows that F is an equivalence of categories such that $FI(b) \cong T'_{-\varepsilon}(b)$ and $F\nabla_{\varepsilon}(b) \cong \Delta'_{-\varepsilon}(b)$ for all $b \in \mathbf{B}$. These functors are needed to formulate the following, which is the semi-infinite counterpart of Corollary 4.12. The proof is similar to the finite case; see also Lemma 4.41 below.

Corollary 4.32. *If \mathcal{R} is a lower finite or an upper finite ε -stratified category and \mathcal{R}' is the Ringel dual category relative to some ε -tilting generator as above, the strata \mathcal{R}_{λ} and \mathcal{R}'_{λ} are equivalent for all $\lambda \in \Lambda$. More precisely:*

- (1) *If $\varepsilon(\lambda) = +$ the functor $F_{\lambda} := (j')^{\lambda} \circ (i')^{\lambda}_{\geq \lambda} \circ F \circ i_{\leq \lambda} \circ j_{*}^{\lambda} : \mathcal{R}_{\lambda} \rightarrow \mathcal{R}'_{\lambda}$ is an equivalence of categories taking $L_{\lambda}(b) = j^{\lambda}L(b)$ to $L'_{\lambda}(b) = (j')^{\lambda}L'(b)$.*
- (2) *If $\varepsilon(\lambda) = -$ the functor $G_{\lambda} := (j')^{\lambda} \circ (i')^{\lambda}_{\geq \lambda} \circ G \circ i_{\leq \lambda} \circ j_i^{\lambda} : \mathcal{R}_{\lambda} \rightarrow \mathcal{R}'_{\lambda}$ is an equivalence of categories taking $L_{\lambda}(b) = j^{\lambda}L(b)$ to $L'_{\lambda}(b) = (j')^{\lambda}L'(b)$.*

In view of Corollary 4.4, Corollary 4.13 can be applied also in any lower finite ε -stratified category (without any need to appeal to semi-infinite Ringel duality). In particular, if \mathcal{R} is a lower finite $+$ -stratified (resp., $-$ -stratified) category then all $V \in \nabla(\mathcal{R})$ (resp., $V \in \Delta(\mathcal{R})$) have finite $-$ -tilting resolutions (resp., finite $+$ -tilting coresolutions). Using Theorem 4.25, one sees that this assertion is equivalent to Lemma 3.43.

We have not investigated derived equivalences or any analog of Theorem 4.16 in the semi-infinite setting.

Proof of Theorem 4.25. We may assume that $\mathcal{R} = \text{comod}_{\text{fd}} C$ for a coalgebra C . Let $B := C^*$ be the dual algebra, so that \mathcal{R} is identified also with $B\text{-mod}_{\text{fd}}$. We can replace the ε -tilting generator $T = \bigoplus_{i \in I} T_i$ with any other. This just has the effect of transforming A into a Morita equivalent locally unital algebra. Consequently, without loss of generality, we may assume that $I = \mathbf{B}$ and $T = \bigoplus_{b \in \mathbf{B}} T_{\varepsilon}(b)$. Then

$$A = \left(\bigoplus_{a, b \in \mathbf{B}} \text{Hom}_{\mathcal{R}}(T_{\varepsilon}(a), T_{\varepsilon}(b)) \right)^{\text{op}}$$

is a pointed locally finite-dimensional locally unital algebra with (primitive) distinguished idempotents $\{e_b \mid b \in \mathbf{B}\}$. Let $P'(b) := Ae_b$ and $L'(b) := \text{hd } P'(b)$. Then $\mathcal{R}' = A\text{-mod}_{\text{fd}}$ is a Schurian category, the A -modules $\{L'(b) \mid b \in \mathbf{B}\}$ give a full set of pairwise inequivalent irreducible objects, and $P'(b)$ is a projective cover of $L'(b)$ in $\text{Ind}(\mathcal{R}'_{\varepsilon}) = A\text{-mod}$. It is immediate that $(\mathbf{B}, L', \rho, \Lambda, \geq)$ is a stratification of \mathcal{R}' . Let $\Delta'_{-\varepsilon}(b)$ and $\nabla'_{-\varepsilon}(b)$ be its $(-\varepsilon)$ -standard and $(-\varepsilon)$ -costandard objects. Also let $V(b) := F\nabla_{\varepsilon}(b)$. Now one checks that Steps 1–6 from the proof of Theorem 4.10 carry over to the present situation with very minor modifications. We will not rewrite these steps here, but cite them freely below. In particular, Step 6 establishes that \mathcal{R}' is an upper finite $(-\varepsilon)$ -stratified category. Also, $F\nabla_{\varepsilon}(b) \cong \Delta'_{-\varepsilon}(b)$ by Step 5. It just remains to show:

- F restricts to an equivalence of categories between $\nabla_{\varepsilon}^{\text{asc}}(\mathcal{R})$ and $\Delta_{-\varepsilon}^{\text{asc}}(\mathcal{R}')$.
- $FI(b) \cong T'_{-\varepsilon}(b)$, the indecomposable $(-\varepsilon)$ -tilting object of \mathcal{R}' labelled by $b \in \mathbf{B}$.

This requires some different arguments compared to the ones from Steps 7–10 in the proof of Theorem 4.10.

Let Ω be the directed poset consisting of all finite lower sets in Λ . Take $\omega = \Lambda^{\downarrow} \in \Omega$. Let $\nabla_{\varepsilon}(\mathcal{R}, \omega)$ be the full subcategory of $\nabla_{\varepsilon}(\mathcal{R})$ consisting of the ∇_{ε} -filtered objects with sections $\nabla_{\varepsilon}(b)$ for $b \in \mathbf{B}^{\downarrow} := \rho^{-1}(\Lambda^{\downarrow})$. Similarly, we define the subcategory $\Delta_{-\varepsilon}(\mathcal{R}', \omega)$ of $\Delta_{-\varepsilon}(\mathcal{R}')$. By Steps 2 and 5, F restricts to a well-defined functor

$$F : \nabla_{\varepsilon}(\mathcal{R}, \omega) \rightarrow \Delta_{-\varepsilon}(\mathcal{R}', \omega). \quad (4.23)$$

We claim that this is an equivalence of categories. To prove it, let $i : \mathcal{R}^{\downarrow} \rightarrow \mathcal{R}$ be the finite ε -stratified subcategory of \mathcal{R} associated to Λ^{\downarrow} . Let $e := \sum_{b \in \mathbf{B}^{\downarrow}} e_b \in A$. Then $T^{\downarrow} := \bigoplus_{b \in \mathbf{B}^{\downarrow}} T_{\varepsilon}(b)$ is an ε -tilting generator for \mathcal{R}^{\downarrow} . As $\text{End}_{\mathcal{R}^{\downarrow}}(T^{\downarrow})^{\text{op}} = eAe$, the Ringel dual $(\mathcal{R}^{\downarrow})'$ of \mathcal{R}^{\downarrow} relative to T^{\downarrow} is identified with the quotient category $(\mathcal{R}')^{\downarrow} := eAe\text{-mod}_{\text{fd}}$ of

\mathcal{R}' . Let $F^\downarrow := \text{Hom}_{\mathcal{R}}(T^\downarrow, ?) : \mathcal{R}^\downarrow \rightarrow (\mathcal{R}')^\downarrow$ be the corresponding Ringel duality functor. We also know from Theorem 3.42 that $(\mathcal{R}')^\downarrow$ is the finite $(-\varepsilon)$ -stratified quotient of \mathcal{R}' associated to Λ^\downarrow (which is a finite upper set in (Λ, \geq)). Let $j' : \mathcal{R}' \rightarrow (\mathcal{R}')^\downarrow$ be the quotient functor, i.e., the functor defined by multiplication by the idempotent e . For a right C -comodule V , we have that

$$F^\downarrow(i^!V) \cong \bigoplus_{b \in \mathbf{B}^\downarrow} \text{Hom}_{\mathcal{R}}(T_\varepsilon(b), i^!V) \cong e \bigoplus_{b \in \mathbf{B}} \text{Hom}_{\mathcal{R}}(T_\varepsilon(b), V) \cong j'(FV).$$

This shows that

$$F^\downarrow \circ i^! \cong j' \circ F, \quad (4.24)$$

so in particular following diagram commutes up to a natural isomorphism:

$$\begin{array}{ccc} \mathcal{R} & \xrightarrow{F} & \mathcal{R}' \\ i^! \downarrow & & \downarrow j' \\ \mathcal{R}^\downarrow & \xrightarrow{F^\downarrow} & (\mathcal{R}^\downarrow)' \equiv (\mathcal{R}')^\downarrow. \end{array}$$

By Theorem 4.10, F^\downarrow restricts to an equivalence $\nabla_\varepsilon(\mathcal{R}^\downarrow) \rightarrow \Delta_{-\varepsilon}((\mathcal{R}')^\downarrow)$. Also the restrictions $i^! : \nabla_\varepsilon(\mathcal{R}, \omega) \rightarrow \nabla_\varepsilon(\mathcal{R}^\downarrow)$ and $j' : \Delta_{-\varepsilon}(\mathcal{R}', \omega) \rightarrow \Delta_{-\varepsilon}((\mathcal{R}')^\downarrow)$ are equivalences. This is clear for $i^!$. To see it for j' , one shows using Theorem 3.42 that the left adjoint $(j')_!$ gives a quasi-inverse equivalence. Putting these things together, we deduce that (4.23) is an equivalence as claimed.

Now we can show that F defines an equivalence $F : \nabla_\varepsilon^{\text{asc}}(\mathcal{R}) \rightarrow \Delta_{-\varepsilon}^{\text{asc}}(\mathcal{R}')$. Take $V \in \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$. Then V has a distinguished ascending ∇_ε -flag $(V_\omega)_{\omega \in \Omega}$ indexed by the set Ω of finite lower sets in Λ . This is defined by setting $V_\omega := i^!V$ in the notation of the previous paragraph; see the proof of Theorem 3.56. As each comodule $T_\varepsilon(b)$ is finite-dimensional, hence, compact, the functor F commutes with direct limits. Hence, $FV \cong \varinjlim (FV_\omega)$. In fact, $(FV_\omega)_{\omega \in \Omega}$ is the data of an ascending $\Delta_{-\varepsilon}$ -flag in $FV \in \mathcal{R}'$. To see this, we have that $FV_\omega \in \Delta_{-\varepsilon}(\mathcal{R}')$ by the previous paragraph. For $\omega < v$ the quotient V_v/V_ω has a ∇_ε -flag thanks to Corollary 3.58, so $FV_v/FV_\omega \cong F(V_v/V_\omega)$ has a $\Delta_{-\varepsilon}$ -flag. We still need to show that FV is locally finite-dimensional. For this, we prove that $\dim \text{Hom}_A(FV, I'(b)) < \infty$ for each $b \in \mathbf{B}$. Since $I'(b)$ has a finite $\nabla_{-\varepsilon}$ -flag, this reduces to checking that $\dim \text{Hom}_A(FV, \nabla'_{-\varepsilon}(b)) < \infty$ for each b . To see this, pick a finite lower set ω containing $\rho(b)$. Then for $v > \omega$, FV_v/FV_ω has a $\nabla_{-\varepsilon}$ -flag with all sections different from $\nabla'_{-\varepsilon}(b)$, so $\text{Hom}_A(FV_v/FV_\omega, \nabla'_{-\varepsilon}(b)) = \text{Ext}_A^1(FV_v/FV_\omega, \nabla'_{-\varepsilon}(b)) = 0$. It follows that $\text{Hom}_A(FV_v, \nabla'_{-\varepsilon}(b)) \cong \text{Hom}_A(FV_\omega, \nabla'_{-\varepsilon}(b))$ and

$$\text{Hom}_A(FV, \nabla'_{-\varepsilon}(b)) = \text{Hom}_A(\varinjlim (FV_v), \nabla'_{-\varepsilon}(b)) \cong \text{Hom}_A(FV_\omega, \nabla'_{-\varepsilon}(b)),$$

which is finite-dimensional.

At this point, we have proved that F induces a well-defined functor

$$F : \nabla_\varepsilon^{\text{asc}}(\mathcal{R}) \rightarrow \Delta_{-\varepsilon}^{\text{asc}}(\mathcal{R}').$$

We prove that this is an equivalence by showing that the left adjoint $F^* := T \otimes_A ?$ to F gives a quasi-inverse. The left mate of (4.24) gives an isomorphism

$$i \circ (F^\downarrow)^* \cong F^* \circ (j')_!. \quad (4.25)$$

Combining this with Corollary 4.11, we deduce that F^* restricts to a quasi-inverse of the equivalence (4.23) for each $\omega \in \Omega$. Also, F^* commutes with direct limits, and again any $V' \in \Delta_{-\varepsilon}^{\text{asc}}(\mathcal{R}')$ has a distinguished ascending $\Delta_{-\varepsilon}$ -flag $(V'_\omega)_{\omega \in \Omega}$ as we saw in the proof of Theorem 3.37. These facts are enough to show that F^* restricts to a well-defined functor $F^* : \Delta_{-\varepsilon}^{\text{asc}}(\mathcal{R}') \rightarrow \nabla_\varepsilon^{\text{asc}}(\mathcal{R})$ which is quasi-inverse to F .

Finally, we check that $FI(b) \cong T'_{-\varepsilon}(b)$. Let $V := I(b)$ and $(V_\omega)_{\omega \in \Omega}$ be its distinguished ascending ∇_ε -flag indexed by the set Ω of finite lower sets in Λ as above. Using the same notation as above, for $\omega = \Lambda^\downarrow \in \Omega$ such that $\rho(b) \in \Lambda^\downarrow$, we know that V_ω is an injective hull of $L(b)$ in \mathcal{R}^\downarrow . Hence, by Theorem 4.10, $F^\downarrow V_\omega$ is the indecomposable $(-\varepsilon)$ -tilting object of \mathcal{R}^\downarrow labelled by b . From this, we see that the ascending $\Delta_{-\varepsilon}$ -flag

$(FV_\omega)_{\omega \in \Omega}$ in $FI(b)$ coincides with the distinguished ascending $\Delta_{-\varepsilon}$ -flag in $T'_{-\varepsilon}(b)$ from the construction from the proof of Theorem 4.18. \square

Proof of Theorem 4.27. We may assume that $\mathcal{R} = A\text{-mod}_{\text{fd}}$ for a pointed locally finite-dimensional locally unital algebra $A = \bigoplus_{a,b \in \mathbf{B}} e_a A e_b$, so that T is a locally finite-dimensional left A -module. Let $C := T^{\otimes} \otimes_A T$ viewed as a coalgebra according to (2.14). By Lemma 2.10 this coalgebra is the continuous dual of $B = \text{End}_A(T)^{\text{op}}$, and we may identify \mathcal{R} with the locally finite Abelian category $\text{comod}_{\text{fd}}\text{-}C$. Applying Lemma 2.11, the Ringel duality functor G becomes the functor $T^{\otimes} \otimes_A ? : A\text{-mod} \rightarrow \text{comod}\text{-}C$, with the comodule structure map of $GV := T^{\otimes} \otimes_A V$ being defined as in (2.17). Let

$$I'(b) := GT_\varepsilon(b), \quad L'(b) := \text{soc } I'(b), \quad \nabla'_{-\varepsilon}(b) := G\Delta_\varepsilon(b). \quad (4.26)$$

Each $I'(b)$ is an indecomposable injective right C -comodule, and $\{L'(b) \mid b \in \mathbf{B}\}$ is a full set of pairwise inequivalent irreducible C -comodules. To show that \mathcal{R}' is a lower finite $(-\varepsilon)$ -stratified category, we must show for each finite upper set Λ^\dagger in Λ that the Serre subcategory $(\mathcal{R}')^\dagger$ of \mathcal{R}' generated by $\{L'(b) \mid b \in \mathbf{B}^\dagger := \rho^{-1}(\Lambda^\dagger)\}$ is a finite $(-\varepsilon)$ -stratified category for the induced stratification $(\mathbf{B}^\dagger, L', \rho, \Lambda, \geq)$.

The functor G sends short exact sequences of objects in $\Delta_\varepsilon^{\text{asc}}(\mathcal{R})$ to short exact sequences in $\text{Ind}(\mathcal{R}')$. This follows because $\text{Hom}_{\mathcal{R}}(?, T)$ has this property thanks to the Ext^1 -vanishing from Lemma 3.36. Since $\Delta_\varepsilon(b) \hookrightarrow T_\varepsilon(b)$, we deduce that $\nabla'_{-\varepsilon}(b) \hookrightarrow I'(b)$. Thus, we have that $L'(b) = \text{soc } \nabla'_{-\varepsilon}(b)$.

Now let \mathcal{R}^\dagger be the Serre quotient of \mathcal{R} associated to some finite upper set $\Lambda^\dagger \subseteq \Lambda$ and let $j : \mathcal{R} \rightarrow \mathcal{R}^\dagger$ be the quotient functor. This is a finite ε -stratified category thanks to Theorem 3.42. In fact, $\mathcal{R}^\dagger = A^\dagger\text{-mod}_{\text{fd}}$ where $A^\dagger := eAe$ for $e := \sum_{b \in \mathbf{B}^\dagger} e_b$; the quotient functor j is the idempotent truncation functor defined by multiplying by e . By the upper finite analog of Corollary 4.5, $T^\dagger := eT$ is an ε -tilting generator for \mathcal{R}^\dagger . Let $B^\dagger := \text{End}_{A^\dagger}(T^\dagger)^{\text{op}}$ be its (finite-dimensional) endomorphism algebra. Then $(\mathcal{R}^\dagger)' := B^\dagger\text{-mod}_{\text{fd}}$ is the Ringel dual of \mathcal{R}^\dagger relative to T^\dagger . By the finite Ringel duality from Theorem 4.10, $(\mathcal{R}^\dagger)'$ is a finite $(-\varepsilon)$ -stratified category. Let $G^\dagger := \text{Cohom}_{\mathcal{R}}(T^\dagger, ?) = \text{Hom}_{\mathcal{R}}(?, T^\dagger)^* : \mathcal{R}^\dagger \rightarrow (\mathcal{R}^\dagger)'$ be its Ringel duality functor. The functor j defines an algebra homomorphism $\pi : B \rightarrow B^\dagger$, hence, we get a functor $\pi^* : (\mathcal{R}^\dagger)' \rightarrow \mathcal{R}'$. We claim that this gives an isomorphism identifying $(\mathcal{R}^\dagger)'$ with the subcategory $(\mathcal{R}')^\dagger$ of \mathcal{R}' . This will be proved in the next paragraph. Moreover, making this identification, we have that

$$i' \circ G^\dagger \cong G \circ j!, \quad (4.27)$$

i.e., the following diagram commutes up to natural isomorphism:

$$\begin{array}{ccc} \mathcal{R}^\dagger & \xrightarrow{G^\dagger} & (\mathcal{R}^\dagger)' \cong (\mathcal{R}')^\dagger \\ j! \downarrow & & \downarrow i' \\ \mathcal{R} & \xrightarrow{G} & \mathcal{R}'. \end{array}$$

This follows because the northeast composition is the functor $T^{\otimes} e \otimes_{eAe} ?$ while the southwest composition is $T^{\otimes} \otimes_A Ae \otimes_{eAe} ?$, and $T^{\otimes} e \cong T^{\otimes} \otimes_A Ae$ as bimodules. Since we already know that $(\mathcal{R}^\dagger)'$ is a finite $(-\varepsilon)$ -category, it follows that $(\mathcal{R}')^\dagger$ one too, with costandard objects

$$i'(G^\dagger \Delta_\varepsilon(b)) \cong G(j! \Delta_\varepsilon^\dagger(b)) \cong G\Delta_\varepsilon(b) = \nabla'_{-\varepsilon}(b)$$

thanks again to Theorem 4.10 plus Theorem 3.42(6).

To prove the claim, let $C^\dagger := (B^\dagger)^*$ be the (finite-dimensional) dual coalgebra so that $(\mathcal{R}^\dagger)' = \text{comod}_{\text{fd}}\text{-}C^\dagger$. Consider the short exact sequence

$$0 \longrightarrow Ae \otimes_{eAe} eT \longrightarrow T \longrightarrow Q \longrightarrow 0$$

which comes from the upper finite counterpart of Lemma 3.19(2); thus, $Q \in \Delta_\varepsilon^{\text{asc}}(\mathcal{R})$ and all of its sections are of the form $\Delta_\varepsilon(b)$ for $b \notin \mathbf{B}^\dagger$, while $Ae \otimes_{eAe} eT \in \Delta_\varepsilon(\mathcal{R})$ has

sections of the form $\Delta_\varepsilon(b)$ for $b \in \mathbf{B}^\dagger$. Applying G and using the exactness noted in the second paragraph of the proof, we get a short exact sequence

$$0 \longrightarrow C^\dagger \longrightarrow C \longrightarrow GQ \rightarrow 0.$$

The first map $C^\dagger \rightarrow C$ here is dual to the algebra homomorphism $\pi : B \rightarrow B^\dagger$, so it is a coalgebra homomorphism. It identifies $(\mathcal{R}^\dagger)'$ with the the Abelian subcategory $\text{comod}_{\text{fd}}\text{-}C^\dagger$ of $\mathcal{R}' = \text{comod}_{\text{fd}}\text{-}C$. Note also that the irreducible objects of \mathcal{R}' are $\{L'(b) \mid b \in \mathbf{B}'\}$. To complete the proof of the claim, it suffices using Lemma 2.25 to show that the socle of GQ only has constituents of the form $L'(b)$ for $b \notin \mathbf{B}^\dagger$. Fix an ascending Δ_ε -flag $(V_\omega)_{\omega \in \Omega}$ in Q . As G commutes with direct limits, we deduce that $GQ = \varinjlim(GV_\omega)$. The sections in a Δ_ε -flag in V_ω are $\Delta_\varepsilon(b)$ for $b \notin \mathbf{B}^\dagger$, hence, GV_ω has a $\nabla_{-\varepsilon}$ -flag with sections $\nabla'_{-\varepsilon}(b)$ for $b \notin \mathbf{B}^\dagger$. It follows that $\text{soc}(GV_\omega)$ is of the desired form for each ω , hence, the socle of GQ is too.

We can now complete the proof of the theorem. We have shown already that \mathcal{R}' is a lower finite $(-\varepsilon)$ -stratified category. Theorem 4.10 plus Corollary 4.4 shows for Λ^\dagger chosen to contain $\rho(b)$ that

$$T'_{-\varepsilon}(b) \cong G^\dagger(jP(b)) \cong G(j!(jP(b))) \cong GP(b).$$

Also, for $a, b \in \mathbf{B}^\dagger$, we have that

$$\text{Hom}_{\mathcal{R}'}(T'_{-\varepsilon}(a), T'_{-\varepsilon}(b)) \cong \text{Hom}_{(\mathcal{R}')^\dagger}(T'_{-\varepsilon}(a), T'_{-\varepsilon}(b)) \cong \text{Hom}_{A^\dagger}(A^\dagger e_a, A^\dagger e_b) \cong e_a A e_b.$$

These things are true for all choices of Λ^\dagger , so we see that the Ringel dual of \mathcal{R}' relative to $\bigoplus_{b \in \mathbf{B}} T'_{-\varepsilon}(b)$ is the original category $\mathcal{R} = A\text{-mod}_{\text{fd}}$. This puts us in the situation of Corollary 4.29, and finally we invoke that corollary (whose proof did not depend on Theorem 4.27) to establish that $G : \Delta_\varepsilon^{\text{asc}}(\mathcal{R}) \rightarrow \nabla_{-\varepsilon}^{\text{asc}}(\mathcal{R}')$ is an equivalence. \square

4.5. Essentially finite Ringel duality. To complete our account of infinite versions of Ringel duality, it remains to discuss the essentially finite case. For this, we impose the tilting-bounded assumption from Definition 4.20.

Definition 4.33. Assume \mathcal{R} is an essentially finite ε -stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Assume in addition that \mathcal{R} is tilting-bounded. An ε -tilting generator for \mathcal{R} means an object $T = \bigoplus_{j \in J} T_j \in \text{Env}(\mathcal{R})$ with a given decomposition as a direct sum of objects $T_j \in \text{Tilt}_\varepsilon(\mathcal{R})$ such that each $T_\varepsilon(b)$ appears as an indecomposable summand of T with multiplicity that is non-zero and finite. Then we define the *Ringel dual* of \mathcal{R} relative to T to be the category $\mathcal{R}' := B\text{-mod}_{\text{fd}}$ where

$$B := \left(\bigoplus_{i, j \in J} \text{Hom}_{\mathcal{R}}(T_i, T_j) \right)^{\text{op}}.$$

We denote the system of distinguished idempotents of B arising from the identity endomorphisms of each T_j by $\{f_j \mid j \in J\}$. Also define the two Ringel duality functors

$$F := \bigoplus_{j \in J} \text{Hom}_{\mathcal{R}}(T_j, ?) : \mathcal{R} \rightarrow \mathcal{R}', \quad (4.28)$$

$$G := \text{Cohom}_{\mathcal{R}}(T, ?) = \text{Hom}_{\mathcal{R}}(?, T)^* : \mathcal{R} \rightarrow \mathcal{R}'. \quad (4.29)$$

Theorem 4.34 (Essentially finite Ringel duality). *In the setup of Definition 4.33, the Ringel dual category \mathcal{R}' is a tilting-bounded essentially finite $(-\varepsilon)$ -stratified category with stratification $(\mathbf{B}, L', \rho, \Lambda, \geq)$ and distinguished objects*

$$\begin{aligned} P'(b) &= FT_\varepsilon(b), & I'(b) &= GT_\varepsilon(b), & L'(b) &= \text{hd } P'(b) \cong \text{soc } I'(b), \\ \Delta'_{-\varepsilon}(b) &= F\nabla_\varepsilon(b), & \nabla'_{-\varepsilon}(b) &= G\Delta_\varepsilon(b), & T'_{-\varepsilon}(b) &= FI(b) \cong GP(b). \end{aligned}$$

The restrictions $F : \nabla_\varepsilon(\mathcal{R}) \rightarrow \Delta_{-\varepsilon}(\mathcal{R}')$ and $G : \Delta_\varepsilon(\mathcal{R}) \rightarrow \nabla_{-\varepsilon}(\mathcal{R}')$ are equivalences.

Proof. We may assume that $\mathcal{R} = A\text{-mod}_{\text{fd}}$ for an essentially finite-dimensional pointed locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$. Replacing the ε -tilting generator $T = \bigoplus_{j \in J} T_j$ by any other changes B to a Morita equivalent algebra, so we may as well assume simply that $J = \mathbf{B}$ and $T = \bigoplus_{b \in \mathbf{B}} T_\varepsilon(b)$. Then the algebra $B = \bigoplus_{a,b \in \mathbf{B}} f_a B f_b$ is a pointed locally unital algebra. The assumption that \mathcal{R} is tilting-bounded implies that

$$\sum_{a \in \mathbf{B}} \dim \text{Hom}_{\mathcal{R}}(T_\varepsilon(a), T_\varepsilon(b)) < \infty, \quad \sum_{b \in \mathbf{B}} \dim \text{Hom}_{\mathcal{R}}(T_\varepsilon(a), T_\varepsilon(b)) < \infty$$

for each $a, b \in \mathbf{B}$. Thus, B is essentially finite-dimensional, i.e., \mathcal{R}' is essentially finite Abelian. The module $P'(b) := Af_b$ is an indecomposable projective A -module, and

$$\{L'(b) := \text{hd } P'(b) \mid b \in \mathbf{B}\}$$

is a full set of pairwise inequivalent irreducibles. Now $(\mathbf{B}, L', \rho, \Lambda, \geq)$ defines a stratification of \mathcal{R}' . One checks that Steps 1–12 from the proof of Theorem 4.10 all go through essentially unchanged in the present setting. This completes the proof except for one point: we must observe finally that \mathcal{R}' is tilting-bounded. This follows because the relevant matrix from Definition 4.20 (with each $T_\varepsilon(b)$ now being replaced by $T'_{-\varepsilon}(b)$) is the Cartan matrix

$$\left(\dim \text{Hom}_A(P(a), P(b)) \right)_{a,b \in \mathbf{B}}$$

of A . Its rows and columns have only finitely many non-zero entries as A is essentially finite-dimensional. \square

Corollary 4.35 (Essentially finite double centralizer property). *Continuing in the general setup of Definition 4.33, suppose that the ε -stratified category \mathcal{R} is $A\text{-mod}_{\text{fd}}$ for an essentially finite-dimensional locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$, so that $T = \bigoplus_{j \in J} T_j$ is an (A, B) -bimodule. For $i \in I$, let $T'_i := (e_i T)^* \in B\text{-mod}_{\text{fd}}$, so that $T' := \bigoplus_{i \in I} T'_i$ is a (B, A) -bimodule.*

- (1) *The module $T' = \bigoplus_{i \in I} T'_i$ is a $(-\varepsilon)$ -tilting generator for $\mathcal{R}' = B\text{-mod}_{\text{fd}}$ and there is an algebra isomorphism*

$$\mu : A \xrightarrow{\sim} \left(\bigoplus_{i,j \in I} \text{Hom}_{\mathcal{R}'}(T'_i, T'_j) \right)^{\text{op}} \quad (4.30)$$

sending $a \in e_i A e_j$ to $\mu(a) : T'_i \rightarrow T'_j, t \mapsto ta$. So the Ringel dual of \mathcal{R}' relative to $T' = \bigoplus_{i \in I} T'_i$ is equivalent to the original category \mathcal{R} .

- (2) *Denote the Ringel duality functors from \mathcal{R}' to \mathcal{R} by*

$$G_* := \bigoplus_{i \in I} \text{Hom}_{\mathcal{R}'}(T'_i, ?) : \mathcal{R}' \rightarrow \mathcal{R}, \quad (4.31)$$

$$F^* := \text{Cohom}_{\mathcal{R}}(T', ?) = \text{Hom}_{\mathcal{R}}(?, T')^* : \mathcal{R}' \rightarrow \mathcal{R}. \quad (4.32)$$

respectively. We have that $F^ \cong T \otimes_B ?$ and $G \cong T' \otimes_A ?$, hence, (F^*, F) and (G, G_*) are adjoint pairs.*

Proof. For (1), note that $\bigoplus_{i \in I} G(Ae_i)$ is a $(-\varepsilon)$ -tilting generator for \mathcal{R}' since $GP(b) \cong T'_{-\varepsilon}(b)$ for $b \in \mathbf{B}$. Actually, $G(Ae_i) = \text{Hom}_A(Ae_i, T)^* \cong (e_i T)^* = T'_i$. Thus, $T' = \bigoplus_{i \in I} T'_i$ is a $(-\varepsilon)$ -tilting generator for \mathcal{R}' . To obtain the isomorphism between A and the locally finite endomorphism algebra of T' , apply the functor G to the canonical isomorphism $A \cong \left(\bigoplus_{i,j \in I} \text{Hom}_A(Ae_i, Ae_j) \right)^{\text{op}}$. To prove (2), we note first that $F^*(Bf_j) \cong T \otimes_B Bf_j$. It then follows that $F^*(V) \cong T \otimes_B V$ on any finite-dimensional B -module V by taking a resolution $P_2 \rightarrow P_1 \rightarrow V \rightarrow 0$ in which P_1, P_2 are direct sums of modules of the form Bf_j , then using the Five Lemma. The argument for G is similar. \square

We leave it to the reader to adapt Corollary 4.12 to the essentially finite setting.

4.6. Tilting-rigidity. We begin by recalling some well-known definitions:

(QF) A finite Abelian category \mathcal{R} is *quasi-Frobenius* if all projective objects are injective. In that case, there is a unique bijection $\nu : \mathbf{B} \rightarrow \mathbf{B}$, the *Nakayama permutation*, such that

$$P(b) \cong I(\nu(b))$$

for each $b \in \mathbf{B}$, where $P(b)$ and $I(b)$ are projective covers and injective hulls of the irreducible objects $\{L(b) \mid b \in \mathbf{B}\}$.

(WS) A finite Abelian category \mathcal{R} is *weakly symmetric* if it is quasi-Frobenius with Nakayama permutation being the identity function. Equivalently, $P(b) \cong I(b)$ for all $b \in \mathbf{B}$.

(S) A finite Abelian category \mathcal{R} is *symmetric* if there is a natural isomorphism of vector spaces

$$\mathrm{Hom}_{\mathcal{R}}(P, V) \cong \mathrm{Hom}_{\mathcal{R}}(V, P)^* \quad (4.33)$$

for all $P, V \in \mathcal{R}$ with P projective.

These are equivalent to saying that every algebra realization A of \mathcal{R} is quasi-Frobenius, Frobenius, or symmetric, respectively; see [GHK, §4.4] and [Ric, Th. 3.1]. Of course, (QF) \Rightarrow (WS) \Rightarrow (S). We are going to investigate some properties of fully stratified categories which involve the properties (QF), (WS) and (S) at the level of strata.

We assume from now on that \mathcal{R} is a fully stratified category, by which we mean a fully stratified category of any one of the four types, finite, essentially finite, upper finite or lower finite. We use the usual notation $(\mathbf{B}, L, \rho, \Lambda, \leq)$ for its stratification.

Definition 4.36. Let \mathcal{R} be a fully stratified category. We say that \mathcal{R} is *tilting-rigid* if

$$\mathcal{Tilt}_+(\mathcal{R}) = \mathcal{Tilt}_-(\mathcal{R}).$$

For this to make sense in the essentially finite case, it is necessary to assume implicitly that \mathcal{R} is tilting-bounded in the sense of Definition 4.20 for some choice (equivalently, all choices) of sign function ε .

Highest weight categories are automatically tilting-rigid for trivial reasons, so that Definition 4.36 is not needed when working just with highest weight categories. The importance of tilting-rigidity first became apparent in the context of fibered highest weight categories in [MO], [FM], where it is formulated as the property “tilting = cotilting”. The following lemma shows in a tilting-rigid category that the subcategories $\mathcal{Tilt}_\varepsilon(\mathcal{R})$ coincide for all choices of ε , so that we can denote them all simply by $\mathcal{Tilt}(\mathcal{R})$.

Theorem 4.37 (Tilting-rigid categories have quasi-Frobenius strata). *Let \mathcal{R} be a tilting-rigid fully stratified category. There is a unique bijection $\nu : \mathbf{B} \rightarrow \mathbf{B}$ such that*

$$T_+(b) \cong T_-(\nu(b)).$$

For $\lambda \in \Lambda$, this function leaves $\mathbf{B}_\lambda \subseteq \mathbf{B}$ invariant, and the stratum \mathcal{R}_λ is quasi-Frobenius with Nakayama permutation $\nu|_{\mathbf{B}_\lambda}$. Moreover, for any sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$, we have that

$$T_\varepsilon(b) \cong \begin{cases} T_+(b) & \text{if } \varepsilon(\lambda) = +, \\ T_+(\nu^{-1}(b)) & \text{if } \varepsilon(\lambda) = -. \end{cases} \quad (4.34)$$

Proof. There is obviously a unique function $\nu : \mathbf{B} \rightarrow \mathbf{B}$ such that $T_+(b) \cong T_-(\nu(b))$. This function is injective and leaves each of the finite subsets \mathbf{B}_λ invariant, hence, it is actually a bijection. To see that \mathcal{R}_λ is quasi-Frobenius with $\nu|_{\mathbf{B}_\lambda}$ as its Nakayama permutation, we must show that $P_\lambda(b) \cong I_\lambda(\nu(b))$ for each $b \in \mathbf{B}_\lambda$. This follows using $T_+(b) \cong T_-(\nu(b))$ together with Theorem 4.2(3) or Theorem 4.18(3) (which one depends on the particular setting we are in). Finally, take $b \in \mathbf{B}_\lambda$ and a sign function ε . Then $T_+(b) \cong T_-(\nu(b))$ has both a Δ -flag and a ∇ -flag, hence, it has a Δ_ε -flag and a ∇_ε -flag. It follows that it is isomorphic to $T_\varepsilon(b')$ for a unique $b' \in \mathbf{B}_\lambda$. Applying j^λ and using Theorems 4.2 or 4.18 again gives that $b' = b$ if $\varepsilon(\lambda) = +$ or $b' = \nu(b)$ if $\varepsilon(\lambda) = -$, and the formula (4.34) follows. \square

The argument used to prove the next lemma is based on the proof of [CM, Th. 2.2]. Note this proves Conjecture 4.15 assuming an additional hypothesis on the strata.

Lemma 4.38. *Suppose that \mathcal{R} is a finite fully stratified category and $\varepsilon : \Lambda \rightarrow \{\pm\}$ is some given sign function.*

- (1) *Assume that $L_\lambda(a)$ is isomorphic to a subobject of a projective object in \mathcal{R}_λ for all $a \in \mathbf{B}_\lambda$ and $\lambda \in \Lambda$ with $\varepsilon(\lambda) = +$. Then for $b \in \mathbf{B}$, $T_\varepsilon(b)$ has finite injective dimension if and only if $T_\varepsilon(b) \in \mathcal{Tilt}_-(\mathcal{R})$.*
- (2) *Assume that $L_\lambda(a)$ is isomorphic to a quotient of an injective object in \mathcal{R}_λ for all $a \in \mathbf{B}_\lambda$ and $\lambda \in \Lambda$ with $\varepsilon(\lambda) = -$. Then for $b \in \mathbf{B}$, $T_\varepsilon(b)$ has finite projective dimension if and only if $T_\varepsilon(b) \in \mathcal{Tilt}_+(\mathcal{R})$.*

Proof. We just prove (1), (2) being the equivalent dual statement. If $T_\varepsilon(b) \in \mathcal{Tilt}_-(\mathcal{R})$ then $T_\varepsilon(b)$ has a ∇ -flag, so it has finite injective dimension thanks to Corollary 3.24. Conversely, suppose that $T_\varepsilon(b)$ has finite injective dimension. Since $T_\varepsilon(b) \in \mathcal{Tilt}_\varepsilon(b)$, it has both a Δ_ε -flag and a ∇_ε -flag. Hence, as \mathcal{R} is fully stratified, it has both a $\bar{\Delta}$ -flag and a $\bar{\nabla}$ -flag. To show that $T_\varepsilon(b) \in \mathcal{Tilt}_-(\mathcal{R})$, it remains to show that $T_\varepsilon(b)$ has a ∇ -flag. This follows from the homological criterion (Theorem 3.11) if we can show that $\text{Ext}_{\mathcal{R}}^1(\bar{\Delta}(c), T_\varepsilon(b)) = 0$ for all $c \in \mathbf{B}$. By assumption, $T_\varepsilon(b)$ has finite injective dimension, so there is a greatest d such that $\text{Ext}_{\mathcal{R}}^d(\bar{\Delta}(a), T_\varepsilon(b)) \neq 0$ for some $a \in \mathbf{B}$. Now the goal is to show that $d = 0$.

Suppose for a contradiction that $d \neq 0$. Since $\text{Ext}_{\mathcal{R}}^d(\Delta_\varepsilon(a), T_\varepsilon(b)) = 0$, we must have that $a \in \mathbf{B}_\lambda$ for λ with $\varepsilon(\lambda) = +$. By the assumption on strata, there exists $a' \in \mathbf{B}_\lambda$ such that $L_\lambda(a) \hookrightarrow P_\lambda(a')$. Let $0 = V_0 < \dots < V_n = \Delta(a')$ be the $\bar{\Delta}$ -flag for $\Delta(a')$ obtained by applying the exact functor j_1^λ to a composition series for $P_\lambda(a')$ chosen so that its bottom section is isomorphic to $L_\lambda(a)$. For each $r = 1, \dots, n$ we have that $V_r/V_{r-1} \cong \bar{\Delta}(a_r)$ for some $a_r \in \mathbf{B}_\lambda$ with $a_1 = a$. Applying $\text{Hom}_R(?, T_\varepsilon(b))$ to the short exact sequence $0 \rightarrow V_{r-1} \rightarrow V_r \rightarrow \bar{\Delta}(a_r) \rightarrow 0$ and using $\text{Ext}_{\mathcal{R}}^{d+1}(\bar{\Delta}(a_r), T_\varepsilon(b)) = 0$ gives a surjection $\text{Ext}_{\mathcal{R}}^d(V_r, T_\varepsilon(b)) \rightarrow \text{Ext}_{\mathcal{R}}^d(V_{r-1}, T_\varepsilon(b))$. Since $\text{Ext}_{\mathcal{R}}^d(V_1, T_\varepsilon(b)) \neq 0$ by the choice of a , we deduce that $\text{Ext}_{\mathcal{R}}^d(V_r, T_\varepsilon(b)) \neq 0$ for all $r = 1, \dots, n$. Taking $r = n$ gives $\text{Ext}_{\mathcal{R}}^d(\Delta(a'), T_\varepsilon(b)) \neq 0$. This is a contradiction since $T_\varepsilon(b)$ has a $\bar{\nabla}$ -flag. \square

The following extends [CM, Th. 2.2] from fibered highest weight categories to fully stratified categories; cf. Remark 3.30.

Theorem 4.39 (Homological criterion for tilting-rigidity). *For a finite fully stratified category \mathcal{R} , the following properties are equivalent:*

- (i) \mathcal{R} is tilting-rigid;
- (ii) \mathcal{R} is Gorenstein⁸ and all of its strata are quasi-Frobenius;
- (iii) \mathcal{R} is Gorenstein and for each $\lambda \in \Lambda$ and $b \in \mathbf{B}_\lambda$ the irreducible object $L_\lambda(b)$ appears in the socle of some projective in \mathcal{R}_λ ;
- (iii') \mathcal{R} is Gorenstein and for each $\lambda \in \Lambda$ and $b \in \mathbf{B}_\lambda$ the irreducible object $L_\lambda(b)$ appears in the head of some injective in \mathcal{R}_λ .

Proof. We may assume that $\mathcal{R} = A\text{-mod}_{\text{fd}}$ for a finite-dimensional algebra A .

(i) \Rightarrow (ii). All strata are quasi-Frobenius by Theorem 4.37. The injective left A -module A^* has a finite $-$ -tilting resolution $0 \rightarrow T_n \rightarrow \dots \rightarrow T_1 \rightarrow T_0 \rightarrow A^* \rightarrow 0$ by Corollary 4.14. As \mathcal{R} is tilting-rigid, this is also a finite $+$ -tilting-resolution, so each T_i has a Δ -flag. Using Corollary 3.24, it follows that each T_i has finite projective dimension. We deduce that A^* has finite projective dimension by arguing as in the proof of [Wei, Th. 4.3.1]; cf. the proof of (2) \Rightarrow (1) from [CM, Th. 2.2]. The dual argument gives that A has finite injective dimension. Hence, A is Gorenstein.

⁸All projectives have finite injective dimension and all injectives have finite projective dimension.

(ii) \Rightarrow (iii), (iii'). This follows immediately since $P_\lambda(b) \cong I_\lambda(\nu(b))$ for all $b \in \mathbf{B}_\lambda$, where ν is the Nakayama permutation.

(iii) \Rightarrow (i). It suffices to show that each $T_+(b)$ belongs to $\mathcal{Tilt}_-(\mathcal{R})$. As $\bigoplus_{b \in \mathbf{B}} T_+(b)$ is tilting in the general sense of tilting theory (cf. the discussion before Conjecture 4.15), the assumption that A is Gorenstein together with [HU, Lem. 1.3] implies that $\bigoplus_{b \in \mathbf{B}} T_+(b)$ is cotilting. Hence, it has finite injective dimension, so each $T_+(b)$ has finite injective dimension. Then we apply Lemma 4.38(1) with $\varepsilon = +$.

(iii') \Rightarrow (i). This follows by the dual argument to the proof of (iii) \Rightarrow (i). \square

Corollary 4.40. *If \mathcal{R} is a finite fibered highest weight category, it is tilting-rigid if and only if it is Gorenstein.*

Proof. In a fibered highest weight category each stratum has a unique irreducible object (up to isomorphism). Therefore the second parts of (iii) and (iii') in Theorem 4.39 hold automatically. \square

Now we are going to consider the Ringel dual \mathcal{R}' of a tilting-rigid fully stratified category \mathcal{R} as in Definitions 4.9, 4.24, 4.26 or 4.33 (depending on the setting). These definitions all involve the choice of a sign function ε and the choice of an ε -tilting generator T . By (4.34), an ε -tilting generator for some choice of ε is an ε -tilting generator for all ε , so it makes sense to drop the prefix ε , referring to T simply as a tilting generator. Fixing such a choice, let \mathcal{R}' be the corresponding Ringel dual category, and let F and G be the Ringel duality functors from those definitions together with (4.21) and (4.22) in the lower finite and upper finite cases, respectively. Note these functors only depend on the choice of tilting generator, not on the choice of sign function ε , i.e., they are the same functors for all ε . For each $\lambda \in \Lambda$, there are now *two* equivalences of categories

$$F_\lambda = (j')^\lambda \circ (i')'_{\geq \lambda} \circ F \circ i_{\leq \lambda} \circ j_*^\lambda : \mathcal{R}_\lambda \rightarrow \mathcal{R}'_\lambda, \quad (4.35)$$

$$G_\lambda = (j')^\lambda \circ (i')^*_{\geq \lambda} \circ G \circ i_{\leq \lambda} \circ j_!^\lambda : \mathcal{R}_\lambda \rightarrow \mathcal{R}'_\lambda \quad (4.36)$$

between strata; see Corollary 4.12 (which also holds in the essentially finite case) and Corollary 4.32. The following lemma gives a more explicit description of these functors.

Lemma 4.41. *Let \mathcal{R} be a finite, tilting-bounded essentially finite, upper finite or lower finite ε -stratified category with the usual stratification $(L, \mathbf{B}, \rho, \Lambda, \leq)$. Suppose that \mathcal{R}' is the Ringel dual of \mathcal{R} with respect to some given tilting generator $T = \bigoplus_{i \in I} T_i$ such that the index set I contains \mathbf{B} and T_b ($b \in \mathbf{B}$) is a direct sum of $T_\varepsilon(b)$ and copies of $T_\varepsilon(c)$ for $c \in \mathbf{B}$ with $\rho(c) < \rho(b)$. For $\lambda \in \Lambda$, let $T_\lambda := \bigoplus_{b \in \mathbf{B}_\lambda} T_b \in \mathcal{R}_{\leq \lambda}$. There is an algebra isomorphism*

$$\phi_\lambda : A_\lambda \xrightarrow{\sim} \text{End}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda)^{\text{op}}$$

between the natural algebra realization A_λ for the stratum \mathcal{R}'_λ and the endomorphism algebra of $j^\lambda T_\lambda \in \mathcal{R}_\lambda$. Moreover:

- (1) *If $\varepsilon(\lambda) = +$ then $F_\lambda \cong \text{Hom}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda, ?) : \mathcal{R}_\lambda \rightarrow A_\lambda\text{-mod}_{\text{fd}}$ with the action of A_λ defined via ϕ_λ .*
- (2) *If $\varepsilon(\lambda) = -$ then $G_\lambda \cong \text{Hom}_{\mathcal{R}_\lambda}(?, j^\lambda T_\lambda)^* : \mathcal{R}_\lambda \rightarrow A_\lambda\text{-mod}_{\text{fd}}$ with the action of A_λ defined via ϕ_λ .*

Proof. We just explain the argument in detail if \mathcal{R} is a finite ε -stratified category; the other cases are similar but there are minor notational differences. We have that $\mathcal{R}' = A\text{-mod}_{\text{fd}}$ for $A := \text{End}_{\mathcal{R}}(T)^{\text{op}}$. The functors F and G are $\text{Hom}_{\mathcal{R}}(T, ?)$ and $\text{Hom}_{\mathcal{R}}(?, T)^*$, respectively. Let $e_b \in A$ be the projection of T onto $T_\varepsilon(b)$ and set $e_\lambda := \sum_{b \in \mathbf{B}_\lambda} e_b$. Let $A_{\geq \lambda}$ be the quotient of A by the two-sided ideal generated by the idempotents $\{e_\mu \mid \mu \in \Lambda \text{ with } \mu \not\geq \lambda\}$. This is the natural realization of the Serre subcategory $\mathcal{R}'_{\geq \lambda}$ of \mathcal{R}' . Then the stratum \mathcal{R}'_λ is realized by the basic finite-dimensional algebra $A_\lambda := \bar{e}_\lambda A_{\geq \lambda} \bar{e}_\lambda$, where we write \bar{x} for the canonical image of $x \in A$ under the quotient map $A \twoheadrightarrow A_{\geq \lambda}$.

The idempotents $\{\bar{e}_b \mid b \in \mathbf{B}_\lambda\}$ are representatives for the conjugacy classes of primitive idempotents in A_λ .

By Theorem 4.2(3), $j^\lambda T_\lambda$ is a minimal projective generator for \mathcal{R}_λ if $\varepsilon(\lambda) = +$ or a minimal injective cogenerator for \mathcal{R}_λ if $\varepsilon(\lambda) = -$. In either case, $\text{End}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda)^{\text{op}}$ is the basic algebra realizing the stratum \mathcal{R}_λ . Since \mathcal{R}_λ and \mathcal{R}'_λ are equivalent, it follows that $A_\lambda \cong \text{End}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda)^{\text{op}}$. However, the argument so far does not produce the desired explicit isomorphism ϕ_λ between these algebras. To obtain this, since we have already seen that the dimensions agree, it suffices to construct a surjective algebra homomorphism $\phi_\lambda : A_\lambda \twoheadrightarrow \text{End}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda)^{\text{op}}$.

Let $\mathcal{R}_{\geq \lambda}$ be the Serre quotient of \mathcal{R} associated to the upper set $(\lambda, \infty]$, so that $\mathcal{R}_{\geq \lambda}$ has irreducible objects labelled by $\mathbf{B}_{\geq \lambda}$. Denote the quotient functor by $j^{\geq \lambda} : \mathcal{R} \rightarrow \mathcal{R}_{\geq \lambda}$. The functor $j^{\geq \lambda}$ defines an algebra homomorphism

$$A = \text{End}_{\mathcal{R}}(T)^{\text{op}} \rightarrow \text{End}_{\mathcal{R}_{\geq \lambda}}(j^{\geq \lambda} T)^{\text{op}}. \quad (4.37)$$

This homomorphism is surjective. To see this, Corollary 3.19(2) gives a short exact sequence $0 \rightarrow j_!^{\geq \lambda} j^{\geq \lambda} T \rightarrow T \rightarrow Q \rightarrow 0$ in which Q has a Δ_ε -flag. Applying $\text{Hom}_{\mathcal{R}}(?, T)$ to this gives surjectivity of the first map below:

$$\text{Hom}_{\mathcal{R}}(T, T) \twoheadrightarrow \text{Hom}_{\mathcal{R}}(j_!^{\geq \lambda} j^{\geq \lambda} T, T) \xrightarrow{\sim} \text{Hom}_{\mathcal{R}_{\geq \lambda}}(j^{\geq \lambda} T, j^{\geq \lambda} T).$$

The second map comes from the adjunction. The composite is the map (4.37), so indeed it is surjective. Now we note that this map sends each e_μ for $\mu \not\geq \lambda$ to zero, so it factors through the quotient $A \twoheadrightarrow A_{\geq \lambda}$ to give a surjective homomorphism $A_{\geq \lambda} \twoheadrightarrow \text{End}_{\mathcal{R}_{\geq \lambda}}(j^{\geq \lambda} T)^{\text{op}}$. Then we restrict to $\bar{e}_\lambda A_{\geq \lambda} \bar{e}_\lambda$ to obtain the homomorphism ϕ_λ .

It just remains to prove (1) and (2). The universal property of Serre quotients produces a unique fully faithful functor i_λ making the following diagram of functors commute:

$$\begin{array}{ccc} \mathcal{R} & \xrightarrow{j^{\geq \lambda}} & \mathcal{R}_{\geq \lambda} \\ i_{\leq \lambda} \uparrow & & \uparrow i_\lambda \\ \mathcal{R}_{\leq \lambda} & \xrightarrow{j^\lambda} & \mathcal{R}_\lambda. \end{array}$$

Thus, $j^{\geq \lambda} \circ i_{\leq \lambda} \cong i_\lambda \circ j^\lambda$. Composing on the left with $j_*^{\geq \lambda}$ and on the right with j_*^λ , using that $j^\lambda \circ j_*^\lambda \cong \text{Id}$ and $j_*^{\geq \lambda} \circ j^{\geq \lambda} \cong \text{Id}$ on objects in the image of $i_{\leq \lambda} \circ j_*^\lambda$, we deduce that

$$j_*^{\geq \lambda} \circ i_\lambda \cong i_{\leq \lambda} \circ j_*^\lambda. \quad (4.38)$$

Using this, we have that

$$F_\lambda \cong (j')^\lambda((i')_{\geq \lambda}^! \text{Hom}_{\mathcal{R}}(T, j_*^{\geq \lambda}(i_\lambda?))) \cong \bar{e}_\lambda \text{Hom}_{\mathcal{R}_{\geq \lambda}}(j^{\geq \lambda} T, i_\lambda?) = \text{Hom}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda, ?),$$

proving (1). The proof of (2) is similar, using the isomorphism $j_!^{\geq \lambda} \circ i_\lambda \cong i_{\leq \lambda} \circ j_!^\lambda$ in place of (4.38). \square

Returning to the setup before the lemma, so \mathcal{R} is a tilting-rigid fully stratified category and \mathcal{R}' is its Ringel dual relative to some tilting generator T , we next discuss the labelling of irreducible objects in \mathcal{R}' . In the general tilting-rigid setting, this depends on a choice of sign function ε , since one needs to fix a specific labelling $\{T_\varepsilon(b) \mid b \in \mathbf{B}\}$ of the isomorphism classes of indecomposable summands of T . Put another way, the labelling of irreducible objects in \mathcal{R}' depends on a labelling $\{L'_b(\lambda) \mid b \in \mathbf{B}_\lambda\}$ of irreducible objects in each of the strata \mathcal{R}'_λ , which we do given a choice of ε by declaring that

$$L'_\lambda(b) := \begin{cases} F_\lambda L_\lambda(b) & \text{if } \varepsilon(\lambda) = +, \\ G_\lambda L_\lambda(b) & \text{if } \varepsilon(\lambda) = -. \end{cases} \quad (4.39)$$

In the next theorem, we see for the first time the advantage of assuming that all of the strata of \mathcal{R} are symmetric, or at least weakly symmetric, since then the labelling of irreducibles in \mathcal{R}' does not depend on the choice of ε here.

Theorem 4.42 (Ringel duality for tilting-rigid fully stratified categories). *Let \mathcal{R} be a tilting-rigid fully stratified category. The Ringel dual \mathcal{R}' of \mathcal{R} with respect to some tilting generator is again tilting-rigid. Moreover, the following hold for $\lambda \in \Lambda$:*

- (1) \mathcal{R}_λ is weakly symmetric if and only if $F_\lambda L_\lambda(b) \cong G_\lambda L_\lambda(b)$ for all $b \in \mathbf{B}_\lambda$.
- (2) \mathcal{R}_λ is symmetric if and only if $F_\lambda \cong G_\lambda$.

Proof. Taking $\varepsilon = +$ in the appropriate Ringel duality theorem (one of Theorems 4.10, 4.25, 4.27 or 4.34) gives that \mathcal{R}' is $--$ -stratified with indecomposable $--$ -tilting objects $\{FI(b) \mid b \in \mathbf{B}\}$ in the finite, lower finite or essentially finite cases and $\{GP(b) \mid b \in \mathbf{B}\}$ in the finite, upper finite or essentially finite cases. Taking $\varepsilon = -$ gives that \mathcal{R}' is $+-$ -stratified with indecomposable $+-$ -tilting objects $\{FI(b) \mid b \in \mathbf{B}\}$ in the finite, lower finite or essentially finite cases and $\{GP(b) \mid b \in \mathbf{B}\}$ in the finite, upper finite or essentially finite cases. It follows \mathcal{R}' is fully stratified and its indecomposable $--$ -tilting objects and $+-$ -tilting objects are the same, i.e., $\mathcal{T}ilt_+(\mathcal{R}') = \mathcal{T}ilt_-(\mathcal{R}')$ and \mathcal{R}' is tilting-rigid.

To prove (1) and (2), let ε be any sign function. We may assume that the tilting generator is $T = \bigoplus_{b \in \mathbf{B}} T_\varepsilon(b)$. Let $T_\lambda := \bigoplus_{b \in \mathbf{B}_\lambda} T_\varepsilon(b)$ and $A_\lambda \cong \text{End}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda)^{\text{op}}$ be as in Lemma 4.41. Using the explicit descriptions of F_λ and G_λ from Lemma 4.41(1)–(2), we deduce that $F_\lambda L_\lambda(b) \cong G_\lambda L_\lambda(b)$ if and only if

$$\text{Hom}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda, L_\lambda(b)) \cong \text{Hom}_{\mathcal{R}_\lambda}(L_\lambda(b), j^\lambda T_\lambda)^*$$

as left A_λ -modules (notation as in Lemma 4.41). The left hand side is the irreducible A_λ -module associated to the primitive idempotent that is the projection of $j^\lambda T_\lambda$ onto the summand isomorphic to $P_\lambda(b)$, and the right hand side is the irreducible A_λ -module associated to the primitive idempotent that is the projection of $j^\lambda T_\lambda$ onto the summand isomorphic to $I_\lambda(b)$. Thus, these modules are isomorphic for all $b \in \mathbf{B}_\lambda$ if and only if $P_\lambda(b) \cong I_\lambda(b)$ for all $b \in \mathbf{B}_\lambda$, i.e., the Nakayama permutation of \mathcal{R}_λ is the identity, and \mathcal{R}_λ is weakly symmetric. This proves (1).

To prove (2), using Lemma 4.41 again, we have that $F_\lambda \cong G_\lambda$ if and only if there is a natural isomorphism of left A_λ -modules

$$\text{Hom}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda, V) \cong \text{Hom}_{\mathcal{R}_\lambda}(V, j^\lambda T_\lambda)^*$$

for $V \in \mathcal{R}_\lambda$. Since $j^\lambda T_\lambda$ is a projective generator for \mathcal{R}_λ and $A_\lambda = \text{End}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda)^{\text{op}}$, there is such an A_λ -module isomorphism if and only if there is a natural vector space isomorphism as in (4.33) for all $P, V \in \mathcal{R}_\lambda$ with P projective, i.e., \mathcal{R}_λ is symmetric according to the definition we gave earlier. \square

In the sequel, we will only consider tilting-rigid fully stratified categories with the additional property that all strata are weakly symmetric. By Theorem 4.37, a tilting-rigid fully stratified category has this property if and only if $\nu = \text{id}$. Thus, a fully stratified category is tilting-rigid with weakly symmetric strata if and only if

$$T_+(b) \cong T_-(b) \tag{4.40}$$

for all $b \in \mathbf{B}$. In that case, $T_+(b) \cong T_\varepsilon(b)$ for all sign functions ε , so that one can simply write $T(b)$ in place of $T_\varepsilon(b)$. Moreover, if \mathcal{R}' is the Ringel dual category to \mathcal{R} with respect to some tilting generator, the irreducible objects of \mathcal{R}' are labelled unambiguously by the set \mathbf{B} ; the induced labelling of irreducible objects of the stratum \mathcal{R}'_λ satisfies

$$L'_\lambda(b) \cong F_\lambda L_\lambda(b) \cong G_\lambda L_\lambda(b) \tag{4.41}$$

for all $\lambda \in \Lambda$ and $b \in \mathbf{B}_\lambda$.

4.7. Bases for morphism spaces between Δ - and ∇ -filtered objects. In this subsection, we explain how to extend the construction of [AST, Th. 3.1] first to ε -stratified and then to fully stratified categories. These results will be used in the next section to construct triangular bases for endomorphism algebras of tilting generators.

Theorem 4.43. *Let \mathcal{R} be a finite, lower finite or tilting-bounded essentially finite ε -stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Suppose for each $b \in \mathbf{B}$ that we are given $T_b \in \text{Tilt}_\varepsilon(\mathcal{R})$ such that T_b is a direct sum of $T_\varepsilon(b)$ and copies of $T_\varepsilon(c)$ for c with $\rho(c) < \rho(b)$. Take $M \in \Delta_\varepsilon(\mathcal{R})$ and $N \in \nabla_\varepsilon(\mathcal{R})$. For each $b \in \mathbf{B}$, choose an embedding $\iota_b : \Delta_\varepsilon(b) \hookrightarrow T_b$, a projection $\pi_b : T_b \twoheadrightarrow \nabla_\varepsilon(b)$, and subsets*

$$Y_b \subset \text{Hom}_{\mathcal{R}}(M, T_b), \quad X_b \subset \text{Hom}_{\mathcal{R}}(T_b, N)$$

so that $\{\bar{y} := \pi_b \circ y \mid y \in Y_b\}$ is a basis for $\text{Hom}_{\mathcal{R}}(M, \nabla_\varepsilon(b))$ and $\{\bar{x} := x \circ \iota_b \mid x \in X_b\}$ is a basis for $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), N)$, as illustrated by the diagram:

$$\begin{array}{ccccc}
 & & \Delta_\varepsilon(b) & & \\
 & & \downarrow \iota_b & \searrow \bar{x} & \\
 M & \xrightarrow{y} & T_b & \xrightarrow{x} & N \\
 & \searrow \bar{y} & \downarrow \pi_b & & \\
 & & \nabla_\varepsilon(b) & &
 \end{array} \tag{4.42}$$

Then the morphisms $x \circ y$ for all $(y, x) \in \bigcup_{b \in \mathbf{B}} Y_b \times X_b$ give a basis for $\text{Hom}_{\mathcal{R}}(M, N)$.

Proof. We proceed by induction on $\ell_{\Delta_\varepsilon}(M) + \ell_{\nabla_\varepsilon}(N)$ where $\ell_{\Delta_\varepsilon}(M) := \sum_{b \in \mathbf{B}} (M : \Delta_\varepsilon(b))$ and $\ell_{\nabla_\varepsilon}(N) := \sum_{b \in \mathbf{B}} (N : \nabla_\varepsilon(b))$. The base case is this number is zero, hence, $M = N = 0$ too, which is trivial. For the induction step, we can replace \mathcal{R} by the Serre subcategory of \mathcal{R} associated to the lower set of Λ generated by all $\{\lambda \mid (M : \Delta_\varepsilon(b)) + (N : \nabla_\varepsilon(b)) \neq 0 \text{ for some } b \in \mathbf{B}_\lambda\}$ to assume that there is some maximal element $\lambda \in \Lambda$ such that $(M : \Delta_\varepsilon(b)) + (N : \nabla_\varepsilon(b)) \neq 0$ for some $b \in \mathbf{B}_\lambda$. Then we let $\Lambda^\perp := \Lambda \setminus \{\lambda\}$, $\mathbf{B}^\perp := \rho^{-1}(\Lambda^\perp)$, and $i : \mathcal{R}^\perp \rightarrow \mathcal{R}$ be the natural inclusion of the corresponding Serre subcategory of \mathcal{R} . Let $j : \mathcal{R} \rightarrow \mathcal{R}_\lambda$ be the quotient functor.

In this paragraph, we treat the special case $N \in \mathcal{R}^\perp$. Let $M^\perp := i^*M$. Note by the choice of λ that $|\ell_{\Delta_\varepsilon}(M^\perp) + \ell_{\nabla_\varepsilon}(N)| < \ell_{\Delta_\varepsilon}(M) + \ell_{\nabla_\varepsilon}(N)$. By (3.10) and Theorem 3.17(2), we have that $M^\perp \in \Delta_\varepsilon(\mathcal{R}^\perp)$, and there is a short exact sequence $0 \rightarrow K \rightarrow M \rightarrow M^\perp \rightarrow 0$ where K has a Δ_ε -flag with sections of the form $\Delta_\varepsilon(b)$ for $b \in \mathbf{B}_\lambda$. It follows that the natural inclusion $\text{Hom}_{\mathcal{R}}(M^\perp, N) \hookrightarrow \text{Hom}_{\mathcal{R}}(M, N)$ is an isomorphism. For $b \in \mathbf{B}^\perp$, all of the morphisms $\{y : M \rightarrow T_b \mid y \in Y_b\}$ factor through M^\perp too. Hence, we can apply the induction hypothesis to deduce that the morphisms $x \circ y$ for all $(y, x) \in \bigcup_{b \in \mathbf{B}^\perp} Y_b \times X_b$ give a basis for $\text{Hom}_{\mathcal{R}}(M^\perp, N) = \text{Hom}_{\mathcal{R}}(M, N)$. Since $X_b = \emptyset$ for $b \in \mathbf{B}_\lambda$, we have that $\bigcup_{b \in \mathbf{B}} Y_b \times X_b = \bigcup_{b \in \mathbf{B}^\perp} Y_b \times X_b$, so this is just what is needed.

Now suppose that $N \notin \mathcal{R}^\perp$ and let $N^\perp := i^!N \in \mathcal{R}^\perp$. We again have that $|\ell_{\Delta_\varepsilon}(M^\perp) + \ell_{\nabla_\varepsilon}(N^\perp)| < \ell_{\Delta_\varepsilon}(M) + \ell_{\nabla_\varepsilon}(N)$. By (3.10) and Theorem 3.17(4), we have that $N^\perp \in \nabla_\varepsilon(\mathcal{R}^\perp)$, and there is a short exact sequence $0 \rightarrow N^\perp \rightarrow N \xrightarrow{\pi} Q \rightarrow 0$ where Q has a ∇_ε -flag with sections of the form $\nabla_\varepsilon(b)$ for $b \in \mathbf{B}_\lambda$. Applying $\text{Hom}_{\mathcal{R}}(M, ?)$ to this and using Theorem 3.14 gives a short exact sequence

$$0 \rightarrow \text{Hom}_{\mathcal{R}}(M, N^\perp) \rightarrow \text{Hom}_{\mathcal{R}}(M, N) \rightarrow \text{Hom}_{\mathcal{R}}(M, Q) \rightarrow 0.$$

For $b \in \mathbf{B}^\perp$, the morphisms $\{x : T_b \rightarrow N \mid x \in X_b\}$ have image contained in N^\perp and are lifts of a basis for $\text{Hom}_{\mathcal{R}^\perp}(\Delta_\varepsilon(b), N^\perp)$. By induction, we get that $\text{Hom}_{\mathcal{R}}(M, N^\perp)$ has basis given by the compositions $x \circ y$ for all $(y, x) \in \bigcup_{b \in \mathbf{B}^\perp} Y_b \times X_b$. In view of this and the above short exact sequence, we are therefore reduced to showing that the morphisms $\pi \circ x \circ y$ for $(y, x) \in \bigcup_{b \in \mathbf{B}_\lambda} Y_b \times X_b$ give a basis for $\text{Hom}_{\mathcal{R}}(M, Q)$. We have that $Q \cong j_*j^!Q$ by Corollary 3.19(1), hence, the exact quotient functor j defines isomorphisms $\text{Hom}_{\mathcal{R}}(M, Q) \xrightarrow{\sim} \text{Hom}_{\mathcal{R}_\lambda}(jM, jQ)$. Similarly, $\text{Hom}_{\mathcal{R}}(M, \nabla_\varepsilon(b)) \xrightarrow{\sim} \text{Hom}_{\mathcal{R}_\lambda}(jM, j\nabla_\varepsilon(b))$ and $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), N) \xrightarrow{\sim} \text{Hom}_{\mathcal{R}_\lambda}(j\Delta_\varepsilon(b), jN)$ for $b \in \mathbf{B}_\lambda$. Moreover, $j\pi : jN \rightarrow jQ$ is an isomorphism. Thus, we are reduced to showing that the morphisms $jx \circ jy$ give a basis for $\text{Hom}_{\mathcal{R}_\lambda}(jM, jN)$ for all $(y, x) \in \bigcup_{b \in \mathbf{B}_\lambda} Y_b \times X_b$. The sets of morphisms

$\bar{Y}_b := \{jy : jM \rightarrow jT_b \mid y \in Y_b\}$ and $\bar{X}_b := \{jx : jT_b \rightarrow jN \mid x \in X_b\}$ appearing here are characterized equivalently as lifts of bases for $\text{Hom}_{\mathcal{R}_\lambda}(jM, j\nabla_\varepsilon(b))$ and $\text{Hom}_{\mathcal{R}_\lambda}(j\Delta_\varepsilon(b), jN)$, respectively. Let $\bar{M} := jM$ and $\bar{N} := jN$.

To complete the proof, we consider the two cases $\varepsilon(\lambda) = +$ and $\varepsilon(\lambda) = -$ separately. The arguments are similar, so we just explain the former. In this case, for $b \in \mathbf{B}_\lambda$, we have that $j\nabla_\varepsilon(b) \cong L_\lambda(b)$ and $j\Delta_\varepsilon(b) \cong P_\lambda(b) \cong jT_b$ by Theorem 4.2(3). The module \bar{M} is projective in \mathcal{R}_λ . We are trying to show that the morphisms $\bar{x} \circ \bar{y}$ for all $(\bar{y}, \bar{x}) \in \bigcup_{b \in \mathbf{B}_\lambda} \bar{Y}_b \times \bar{X}_b$ give a basis for $\text{Hom}_{\mathcal{R}_\lambda}(\bar{M}, \bar{N})$ where:

- $\bar{Y}_b \subset \text{Hom}_{\mathcal{R}_\lambda}(\bar{M}, P_\lambda(b))$ is a set lifting a basis of $\text{Hom}_{\mathcal{R}_\lambda}(\bar{M}, L_\lambda(b))$;
- \bar{X}_b is a basis of $\text{Hom}_{\mathcal{R}_\lambda}(P_\lambda(b), \bar{N})$.

Since \bar{M} is projective, the proof reduces to the case that $\bar{M} = P_\lambda(b)$, when the assertion is clear. \square

The following restatement in the special case of a highest weight categories recovers [AST, Th. 3.1].

Corollary 4.44. *Let \mathcal{R} be a finite, lower finite or tilting-bounded essentially finite highest weight category with poset (Λ, \leq) and labelling function L . Suppose for each $\lambda \in \Lambda$ that we are given $T_\lambda \in \text{Tilt}(\mathcal{R})$ such that T_λ is a direct sum of $T(\lambda)$ and copies of $T(\mu)$ for $\mu < \lambda$. Take $M \in \Delta(\mathcal{R})$ and $N \in \nabla(\mathcal{R})$. For each $\lambda \in \Lambda$, choose an embedding $\iota_\lambda : \Delta(\lambda) \hookrightarrow T_\lambda$, a projection $\pi_\lambda : T_\lambda \twoheadrightarrow \nabla(\lambda)$, and subsets*

$$Y_\lambda \subset \text{Hom}_{\mathcal{R}}(M, T_\lambda), \quad X_\lambda \subset \text{Hom}_{\mathcal{R}}(T_\lambda, N)$$

so that $\{\bar{y} := \pi_\lambda \circ y \mid y \in Y_\lambda\}$ is a basis for $\text{Hom}_{\mathcal{R}}(M, \nabla(\lambda))$ and $\{\bar{x} := x \circ \iota_\lambda \mid x \in X_\lambda\}$ is a basis for $\text{Hom}_{\mathcal{R}}(\Delta(\lambda), N)$. Then the morphisms $\bar{x} \circ \bar{y}$ for all $(y, x) \in \bigcup_{\lambda \in \Lambda} Y_\lambda \times X_\lambda$ give a basis for $\text{Hom}_{\mathcal{R}}(M, N)$.

For tilting-rigid fully stratified categories, there is a more refined version of Theorem 4.43.

Theorem 4.45. *Let \mathcal{R} be a finite, lower finite or essentially finite fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ such that \mathcal{R} is tilting-rigid with weakly symmetric strata. Suppose for each $b \in \mathbf{B}$ that we are given $T_b \in \text{Tilt}(\mathcal{R})$ such that T_b is a direct sum of $T(b)$ and copies of $T(c)$ for c with $\rho(c) < \rho(b)$. Take $M \in \Delta(\mathcal{R})$ and $N \in \nabla(\mathcal{R})$. For $a, b \in \mathbf{B}$, choose embeddings $\iota_a : \Delta(a) \hookrightarrow T_a$, $\bar{\iota}_b : \bar{\Delta}(b) \hookrightarrow T_b$, projections $\bar{\pi}_a : T_a \twoheadrightarrow \bar{\nabla}(a)$, $\pi_b : T_b \twoheadrightarrow \nabla(b)$, and subsets*

$$Y_a \subset \text{Hom}_{\mathcal{R}}(M, T_a), \quad H(a, b) \subset \text{Hom}_{\mathcal{R}}(T_a, T_b), \quad X_b \subset \text{Hom}_{\mathcal{R}}(T_b, N)$$

so that $\{\bar{y} := \bar{\pi}_a \circ y \mid y \in Y_a\}$ is a basis for $\text{Hom}_{\mathcal{R}}(M, \bar{\nabla}(a))$, $\{\bar{h} := \pi_b \circ h \circ \iota_a \mid h \in H(a, b)\}$ is a basis for $\text{Hom}_{\mathcal{R}}(\Delta(a), \nabla(b))$, and $\{\bar{x} := x \circ \bar{\iota}_b \mid x \in X_b\}$ is a basis for $\text{Hom}_{\mathcal{R}}(\bar{\Delta}(b), N)$, as illustrated by the diagram:

$$\begin{array}{ccccc}
 \Delta(a) & \xrightarrow{\bar{h}} & \nabla(b) & & \\
 \downarrow \iota_a & & \uparrow \pi_b & & \\
 M & \xrightarrow{y} & T_a & \xrightarrow{h} & T_b & \xrightarrow{x} & N \\
 \searrow \bar{y} & & \downarrow \bar{\pi}_a & & \uparrow \bar{\iota}_b & & \nearrow \bar{x} \\
 & & \bar{\nabla}(a) & & \bar{\Delta}(b) & &
 \end{array} \tag{4.43}$$

Then the morphisms $\bar{x} \circ \bar{h} \circ \bar{y}$ for all $(y, h, x) \in \bigcup_{a, b \in \mathbf{B}} Y_a \times H(a, b) \times X_b$ give a basis for $\text{Hom}_{\mathcal{R}}(M, N)$.

Proof. This follows by the same strategy as was used in the proof of Theorem 4.43. The only substantial difference is in the final paragraph of the proof. By that point, we have reduced to showing for projective and injective objects $\bar{M}, \bar{N} \in \mathcal{R}_\lambda$, respectively, that

the morphisms $\bar{x} \circ \bar{h} \circ \bar{y}$ for all $(\bar{y}, \bar{h}, \bar{x}) \in \bigcup_{a,b \in \mathbf{B}_\lambda} \bar{Y}_a \times \bar{H}(a,b) \times \bar{X}_b$ give a basis for $\text{Hom}_{\mathcal{R}_\lambda}(\bar{M}, \bar{N})$ where:

- $\bar{Y}_a \subset \text{Hom}_{\mathcal{R}_\lambda}(\bar{M}, P_\lambda(a))$ is a set lifting a basis of $\text{Hom}_{\mathcal{R}_\lambda}(\bar{M}, L_\lambda(a))$;
- $\bar{H}(a,b)$ is a basis for $\text{Hom}_{\mathcal{R}_\lambda}(P_\lambda(a), I_\lambda(b))$;
- $\bar{X}_b \subset \text{Hom}_{\mathcal{R}_\lambda}(I_\lambda(b), \bar{N})$ is a set lifting a basis of $\text{Hom}_{\mathcal{R}_\lambda}(L_\lambda(b), \bar{N})$.

Using that \bar{M} is projective and \bar{N} is injective, the proof of this reduces to the case that $\bar{M} = P_\lambda(a)$ and $\bar{N} = I_\lambda(b)$, when the assertion is clear. \square

4.8. Chevalley dualities. Finally, in this section we discuss some further aspects of Ringel duality. These results will be used in the next section to construct symmetric triangular bases for endomorphism algebras of tilting generators. Like in §4.6, the phrase “fully stratified category” means a fully stratified category \mathcal{R} that is either finite, essentially finite, upper finite or lower finite.

Given a finite-dimensional algebra A and an algebra anti-automorphism $\sigma : A \rightarrow A$, there is a contravariant autoequivalence

$$?^\circledast : A\text{-mod}_{\text{fd}} \rightarrow A\text{-mod}_{\text{fd}} \quad (4.44)$$

taking V to its linear dual V^* viewed as a left module by restricting the natural right action along σ . If \mathcal{R} is a finite Abelian category and $?^\vee : \mathcal{R} \rightarrow \mathcal{R}$ is a contravariant autoequivalence, we call a pair (A, σ) consisting of a finite-dimensional algebra A and an anti-automorphism σ a *realization* of $(\mathcal{R}, ?^\vee)$ if there is an equivalence of categories $F : \mathcal{R} \rightarrow A\text{-mod}_{\text{fd}}$ such that $F \circ ?^\vee \cong ?^\circledast \circ F$. The following lemma shows that any contravariant autoequivalence of \mathcal{R} admits a realization in this sense. In fact, we will only ever consider contravariant autoequivalences that preserve isomorphism classes of irreducible objects, in which case we can say a little more about σ as explained at the end of the lemma.

Lemma 4.46. *Let A be a finite-dimensional algebra. Suppose that $?^\vee$ is a contravariant autoequivalence of $A\text{-mod}_{\text{fd}}$. Then there exists an algebra anti-automorphism $\sigma : A \rightarrow A$ such that $?^\vee \cong ?^\circledast$. Moreover, if $?^\vee$ preserves isomorphism classes of irreducible A -modules, then σ can be chosen so that it fixes each of a given set $\{e_i \mid i \in I\}$ of mutually orthogonal idempotents in A .*

Proof. Consider the functor $F := ?^* \circ ?^\vee : A\text{-mod}_{\text{fd}} \rightarrow A^{\text{op}}\text{-mod}_{\text{fd}}$. Since this is right exact and preserves direct sums, we have that $F \cong FA \otimes_A ?$ where FA is the (A^{op}, A) -bimodule obtained by applying F to the regular (A, A) -bimodule A . Note that the right action of $x \in A$ on FA here is defined by applying F to the left A -module homomorphism $r_x : A \rightarrow A, a \mapsto ax$.

Viewing A as a left A^{op} -module with action $x \cdot y := yx$, we claim that $FA \cong A$ as left A^{op} -modules. To see this, let $\{L(b) \mid b \in \mathbf{B}\}$ be a full set of pairwise inequivalent irreducible left A -modules. Then $A \cong \bigoplus_{b \in \mathbf{B}} P(b)^{\oplus \dim L(b)}$ as left A -modules, where $P(b)$ is the projective cover of $L(b)$. Let $\mathbf{B} \rightarrow \mathbf{B}, b \mapsto b'$ be the bijection defined from $L(b)^\vee \cong L(b')$. Then $P(b)^\vee \cong I(b')$, the injective hull of $L(b')$. Hence $FP(b) \cong I(b')^*$ as left A^{op} -modules. Here, $I(b)^*$ is the projective cover of the left A^{op} -module $L(b)^*$. Using that $\dim L(b) = \dim L(b)^*$, we deduce that

$$FA \cong \bigoplus_{b \in \mathbf{B}} (I(b')^*)^{\oplus \dim L(b)} \cong \bigoplus_{b \in \mathbf{B}} (I(b)^*)^{\oplus \dim L(b)^*} \cong A$$

as left A^{op} -modules. This proves the claim. Similarly, under the additional hypothesis that $?^\vee$ preserves isomorphism classes of irreducible objects and we are given mutually orthogonal idempotents $\{e_i \mid i \in I\}$, we get that $F(Ae_i) \cong e_i A$ as left A^{op} -modules for each $i \in I$.

Now we let $\phi : FA \xrightarrow{\sim} A$ be some choice of a left A^{op} -module isomorphism. When the additional hypothesis holds, we may pick this so that it restricts to isomorphisms $F(Ae_i) \xrightarrow{\sim} e_i A$ for each $i \in I$. Transporting the right A -module structure on FA through

ϕ , we make the left A^{op} -module A into an (A^{op}, A) -bimodule, which we will denote by $A_{\sigma^{-1}}$. Explicitly, left action of $x \in A^{\text{op}}$ on $y \in A_{\sigma^{-1}}$ is given by $x \cdot y := yx$ as in the previous paragraph, while the new right action of $x \in A$ is by $y \cdot x := (\phi \circ ((r_x)^\vee)^* \circ \phi^{-1})(y)$. Since $\text{End}_{A^{\text{op}}}(A) \cong A$, this right action of x can be written as left multiplication by a unique element $x' \in A$. The resulting map $A \rightarrow A, x \mapsto x'$ is an algebra anti-automorphism. Let $\sigma : A \rightarrow A$ be the inverse anti-automorphism. Note then that the right action of $x \in A$ on $y \in A_{\sigma^{-1}}$ is by $y \cdot x = \sigma^{-1}(x)y$, explaining our earlier choice of notation. When the additional hypothesis holds, the choice of ϕ ensures that $(e_i)' = e_i$ for $i \in I$, hence, $\sigma(e_i) = e_i$ for each $i \in I$.

For a left A -module V , let ${}_\sigma V$ be V viewed instead as a left A^{op} -module by restricting along σ . Then ${}_\sigma A$ is an (A^{op}, A) -bimodule which is isomorphic via $\sigma : A_{\sigma^{-1}} \xrightarrow{\sim} {}_\sigma A$ to the (A^{op}, A) -bimodule $A_{\sigma^{-1}} \cong FA$ from the previous paragraph. Thus, we have shown that $F \cong A_{\sigma^{-1}} \otimes_A ? \cong {}_\sigma A \otimes_A ? \cong {}_\sigma ? : A\text{-mod}_{\text{fd}} \rightarrow A^{\text{op}}\text{-mod}_{\text{fd}}$. Applying $?^*$ gives finally that $?^\vee \cong ?^\circledast$. \square

Remark 4.47. In the setup of Lemma 4.46, assume that $?^\vee$ preserves isomorphism classes of irreducible A -modules. Then we can take the set of mutually orthogonal idempotents at the end of the lemma to be a mutually orthogonal set $\{e_b \mid b \in \mathbf{B}\}$ of representatives for the conjugacy classes of primitive idempotents in A . Then the lemma shows that we can choose the anti-automorphism σ so that $\sigma(e_b) = e_b$ for all $b \in \mathbf{B}$. Conversely, if $\sigma : A \rightarrow A$ is an anti-automorphism fixing such a set of representatives for the conjugacy classes of primitive idempotents on A , it is obvious that the contravariant autoequivalence $?^\circledast$ preserves isomorphism classes of irreducible A -modules.

To adapt the above from finite Abelian categories to essentially finite Abelian categories, Schurian categories or locally finite Abelian categories, we need the following definitions:

- If $A = \bigoplus_{i,j \in I} e_i A e_j$ is an essentially or locally finite-dimensional locally unital algebra, a locally unital algebra anti-automorphism $\sigma : A \rightarrow A$ gives rise to a contravariant autoequivalence $?^\circledast$ of the categories $A\text{-mod}_{\text{fd}}$ or $A\text{-mod}_{\text{fd}}$, respectively. This is defined by first applying the usual duality from left modules to right modules, either $?^* : A\text{-mod}_{\text{fd}} \rightarrow \text{mod}_{\text{fd}}\text{-}A$ or $?^\circledast : A\text{-mod}_{\text{fd}} \rightarrow \text{mod}_{\text{fd}}\text{-}A$ depending on the case, and then converting right modules back to left modules by restricting along σ .
- If A is a pseudo-compact topological algebra, that is, $A \cong C^*$ for a coalgebra C , an algebra anti-automorphism $\sigma : A \rightarrow A$ gives rise to a contravariant autoequivalence $?^\circledast$ of $A\text{-mod}_{\text{fd}} \cong \text{comod}_{\text{fd}}\text{-}C$. Note in this case that σ is necessarily continuous so that it is the dual of a coalgebra anti-automorphism $\sigma^* : C \rightarrow C$; the definition of the duality $?^\circledast$ could also be formulated in terms of comodules using σ^* .

Then given an essentially finite Abelian category, a Schurian category, or a locally finite Abelian category \mathcal{R} with a contravariant autoequivalence $?^\vee$, a *realization* of $(\mathcal{R}, ?^\vee)$ means a pair (A, σ) consisting of an algebra A and an anti-automorphism $\sigma : A \rightarrow A$ of the appropriate type such that $?^\circledast \circ F \cong F \circ ?^\vee$ for some equivalence F from \mathcal{R} to $A\text{-mod}_{\text{fd}}$, $A\text{-mod}_{\text{fd}}$ or $A\text{-mod}_{\text{fd}}$, respectively. The following lemmas are analogs of Lemma 4.46 in each of these new settings.

Lemma 4.48. *Suppose that $A = \bigoplus_{i,j \in I} e_i A e_j$ is either an essentially or a locally finite-dimensional locally unital algebra. Let $?^\vee$ be a contravariant autoequivalence of $A\text{-mod}_{\text{fd}}$ or $A\text{-mod}_{\text{fd}}$, respectively, which preserves isomorphism classes of irreducible objects. There exists a locally unital algebra anti-automorphism $\sigma : A \rightarrow A$ such that $?^\vee \cong ?^\circledast$.*

Proof. In the locally finite-dimensional case, let $F := ?^\circledast \circ ?^\vee : A\text{-mod}_{\text{fd}} \rightarrow A^{\text{op}}\text{-mod}_{\text{fd}}$. Viewing $\bigoplus_{i \in I} F(Ae_i)$ as an (A^{op}, A) -bimodule in the natural way, we have $\text{tat } F \cong$

$(\bigoplus_{i \in I} F(Ae_i)) \otimes_A ?$. Then we observe for each $i \in I$ that $F(Ae_i) \cong e_i A$ as left A^{op} -modules as $?^\vee$ preserves isomorphism classes of irreducibles. Now argue as in proof of Lemma 4.46. The essentially finite-dimensional case is similar. \square

Lemma 4.49. *Suppose that A is a pseudo-compact topological algebra. Let $?^\vee$ be a contravariant autoequivalence of $A\text{-mod}_{\text{fd}}$ which preserves isomorphism classes of irreducible objects. Then there exists an algebra anti-automorphism $\sigma : A \rightarrow A$ such that $?^\vee \cong ?^\circledast$. Moreover, given a family $\{e_i \mid i \in I\}$ of mutually orthogonal idempotents in A , σ can be chosen so that $\sigma(e_i) = e_i$ for all $i \in I$.*

Proof. The functor $?^\vee : A\text{-mod}_{\text{fd}} \rightarrow A\text{-mod}_{\text{fd}}$ extends to $?^\vee : A\text{-mod}_{\text{pc}} \rightarrow A\text{-mod}_{\text{ds}}$ with $(\varinjlim V_\omega)^\vee := \varinjlim (V_\omega^\vee)$, taking limits over finite-dimensional submodules $V_\omega \leq V$. Composing with $?^*$ gives an equivalence $F := ?^* \circ ?^\vee : A\text{-mod}_{\text{pc}} \rightarrow A^{\text{op}}\text{-mod}_{\text{pc}}$. Moreover, for each $i \in I$ we have that $F(Ae_i) \cong e_i A$ as a (A^{op}, A) -bimodule as $?^\vee$ preserves isomorphism classes of irreducibles. Then we argue as in Lemma 4.46 to obtain an algebra anti-automorphism $\sigma : A \rightarrow A$ with $\sigma(e_i) = e_i$ for each $i \in I$ such that F is isomorphic to the functor $A\text{-mod}_{\text{pc}} \rightarrow A^{\text{op}}\text{-mod}_{\text{pc}}$ defined by restriction along σ . The lemma follows on composing with $?^*$ then restricting to $A\text{-mod}_{\text{fd}}$. \square

With these preliminaries in place, we can now prove a result which explains how to transfer a contravariant autoequivalence on a fully stratified category to its Ringel dual.

Theorem 4.50 (Dualities commute with Ringel duality). *Suppose that \mathcal{R} is a fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ such that \mathcal{R} is tilting-rigid with weakly symmetric strata, i.e., (4.40) holds. Assume also that \mathcal{R} possesses a contravariant autoequivalence $?^\vee$ which preserves isomorphism classes of irreducible objects. Then we have that $T(b)^\vee \cong T(b)$ for all $b \in \mathbf{B}$. Moreover, letting \mathcal{R}' be the Ringel dual category with respect to some choice of tilting generator and F, G be the usual Ringel duality functors, there is an induced contravariant autoequivalence $?^\wedge$ on \mathcal{R}' preserving isomorphism classes of irreducible objects such that*

$$F \circ ?^\vee \cong ?^\wedge \circ G, \quad G \circ ?^\vee \cong ?^\wedge \circ F \quad (4.45)$$

whenever these functors make sense (e.g., these isomorphisms always hold on $\Delta_\varepsilon(\mathcal{R})$ and on $\nabla_\varepsilon(\mathcal{R})$, respectively, for any choice of ε).

Proof. We just explain the proof in the case that \mathcal{R} is a finite fully stratified category, leaving the minor modifications needed in the other three cases to the reader. By Lemma 4.46, we may assume that $\mathcal{R} = A\text{-mod}_{\text{fd}}$ for a finite-dimensional algebra A and that $?^\vee : \mathcal{R} \rightarrow \mathcal{R}$ is the functor $?^\circledast$ taking a left A -module V to the dual right A -module viewed as a left module by restricting the natural right action along some given anti-automorphism $\sigma : A \xrightarrow{\sim} A$. (In the other three cases, one needs to use Lemmas 4.48–4.49 here in place of Lemma 4.46.)

Since $T_+(b)$ has a Δ -flag with $\Delta(b)$ at the bottom, and also a $\bar{\nabla}$ -flag, we see using Lemma 3.3 that $T_+(b)^\vee$ has a ∇ -flag with $\nabla(b)$ at the top, and also a $\bar{\Delta}$ -flag. So it is isomorphic to $T_-(b)$. As \mathcal{R} is tilting-rigid, $T(b) := T_+(b) \cong T_-(b)$, so we have shown that $T(b)^\vee \cong T(b)$ for all $b \in \mathbf{B}$.

We are given some full tilting module T defining the Ringel dual category \mathcal{R}' , i.e., $\mathcal{R}' = B\text{-mod}_{\text{fd}}$ for $B = \text{End}_A(T)^{\text{op}}$. From the previous paragraph, we get that $T \cong T^\vee$. Let $\phi : T \xrightarrow{\sim} T^\vee$ be an isomorphism of left A -modules. Equivalently, ϕ is the data of a non-degenerate pairing $\langle \cdot, \cdot \rangle : T \times T \rightarrow \mathbb{k}$ with $\langle v, w \rangle := \phi(v)(w)$, and we have that $\langle xv, w \rangle = \langle v, \sigma(x)w \rangle$ for $v, w \in T$, $x \in A$. Let $\tau : B \rightarrow B$ be the anti-automorphism of B defined so that $\langle vy, w \rangle = \langle v, w\tau(y) \rangle$ for $v, w \in T$, $y \in B$. It follows that ϕ is also an isomorphism of right B -modules for the right B -module structure on T^\vee obtained by restricting its natural left action on T^* along τ . Now we can define the contravariant autoequivalence $?^\wedge : B\text{-mod}_{\text{fd}} \rightarrow B\text{-mod}_{\text{fd}}$ to be $?^\circledast$.

In this paragraph, we check (4.45). We just prove the first of these isomorphisms; the latter follows from former (with the roles of A and B reversed) on taking adjoints. Take $V \in \mathcal{R}$. Then we have natural left B -module isomorphisms

$$(GV)^\wedge \cong \text{Hom}_A(V, T) \cong \text{Hom}_A(T^\vee, V^\vee) \cong \text{Hom}_A(T, V^\vee) = F(V^\vee),$$

as required. (On the space $\text{Hom}_A(V, T)$ here, the left B -module structure is defined by restricting the natural right action along τ .)

It remains to check that $?\wedge$ preserves isomorphism classes of irreducible objects in \mathcal{R}' . Since the strata are weakly symmetric, we have that

$$\nabla'(b)^\wedge \cong (G\Delta(b))^\wedge \cong F(\Delta(b)^\vee) \cong F\nabla(b) \cong \Delta'(b).$$

This implies that $L'(b)^\wedge \cong L'(b)$. \square

In examples coming from Lie theory, highest weight categories usually come equipped with dualities arising from anti-involutions which restrict to the identity on the Cartan part. The material in the rest of the subsection is an attempt to axiomatize the essential features of such dualities in the more general setting of fully stratified categories. We start with a definition which will be relevant at the level of strata.

Definition 4.51. Let A be a finite-dimensional algebra and $\sigma : A \rightarrow A$ be an anti-involution. We say that A is σ -symmetric if the following hold:

- (σ S1) There is a set $\{e_b \mid b \in \mathbf{B}\}$ of representatives for the conjugacy classes of primitive idempotents in A such that $\sigma(e_b) = e_b$ for all $b \in \mathbf{B}$.
- (σ S2) There is a non-degenerate associative symmetric bilinear form $(\cdot, \cdot) : A \times A \rightarrow \mathbb{k}$ such that $(x, y) = (\sigma(x), \sigma(y))$ for all $x, y \in A$.

If A is σ -symmetric in the sense of Definition 4.51 then it is a symmetric algebra in the usual sense. Moreover, every finitely generated projective left A -module P possesses a non-degenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$ such that $\langle xv, w \rangle = \langle v, \sigma(x)w \rangle$ for $v, w \in P, x \in A$; in particular, $P \cong P^\circledast$. To see this, we may assume without loss of generality that P is indecomposable and that $P = Ae$ for a σ -invariant primitive idempotent e . Then the form $\langle \cdot, \cdot \rangle : P \times P \rightarrow \mathbb{k}$ defined in terms of the given σ -symmetric form (\cdot, \cdot) on A by $\langle v, w \rangle := (\sigma(v), w)$ for $v, w \in P$ has these properties; it is non-degenerate because by associativity

$$A = eAe \oplus [eA(1-e) + (1-e)Ae] \oplus (1-e)A(1-e) \quad (4.46)$$

is an orthogonal decomposition of A with respect to (\cdot, \cdot) and the subspaces $eA(1-e)$ and $(1-e)Ae$ are isotropic.

The following lemma shows that σ -symmetry is preserved by Morita equivalence. The basic point underlying this is that if A is σ -symmetric and $e \in A$ is a σ -invariant idempotent, then σ restricts to an anti-involution of eAe . Moreover a σ -symmetric form (\cdot, \cdot) on A restricts to such a form on eAe so that eAe is also σ -symmetric; the non-degeneracy of this restriction follows from the orthogonal decomposition (4.46).

Lemma 4.52. *Let A be a finite-dimensional algebra which is σ -symmetric for some anti-involution σ . Let B be another finite-dimensional algebra that is Morita equivalent to A , so that there is an equivalence of categories $F : B\text{-mod}_{\text{fd}} \rightarrow A\text{-mod}_{\text{fd}}$. Then B possesses an anti-involution $\tau : B \rightarrow B$ such that $?\circledast \circ F \cong F \circ ?\circledast$, and B is τ -symmetric for any such anti-involution τ . Moreover, τ can be chosen in such a way that it fixes each of some given set $\{f_i \mid i \in I\}$ of mutually orthogonal idempotents in B .*

Proof. Let $\{e_b \mid b \in \mathbf{B}\}$ be a set of mutually orthogonal representatives for the conjugacy classes of primitive idempotents in A with $\sigma(e_b) = e_b$ for all b . Let $e := \sum_{b \in \mathbf{B}} e_b$. Then eAe is the basic algebra that is Morita equivalent to A , and it is σ -symmetric too. The functors $?\circledast$ on $A\text{-mod}_{\text{fd}}$ and $eAe\text{-mod}_{\text{fd}}$ obviously commute with the idempotent truncation functor giving an equivalence $A\text{-mod}_{\text{fd}} \rightarrow eAe\text{-mod}_{\text{fd}}$. All of this means that we can replace A with eAe if necessary to assume that A itself is *basic* with $1 = \sum_{b \in \mathbf{B}} e_b$ being

a decomposition of its identity element into mutually orthogonal σ -invariant primitive idempotents.

Now suppose that B is Morita equivalent to A via some given $F : B\text{-mod}_{\text{fd}} \rightarrow A\text{-mod}_{\text{fd}}$. Let $P := FB$ be the (A, B) -bimodule obtained by applying F to the regular (B, B) -bimodule. Note that $P = \bigoplus_{i \in I} Pf_i$ where $\{f_i \mid i \in I\}$ is the given set of mutually orthogonal idempotents in B ; we are assuming here that $\sum_{i \in I} f_i = 1_B$ which we can clearly do by adding one more idempotent to this set if necessary. As an A -module, we have for each $i \in I$ that $Pf_i \cong \bigoplus_{b \in \mathbf{B}} Ae_b^{\oplus d_i(b)}$ for integers $d_i(b) > 0$; the numbers $d(b) = \sum_{i \in I} d_i(b)$ are the dimensions of the irreducible B -modules. Moreover, $e_i B e_j \cong \text{End}_A(Pe_i, Pe_j)^{\text{op}}$. Fixing such isomorphisms, we may assume simply that $P = \bigoplus_{i \in I} Pf_i$ with $Pf_i = \bigoplus_{b \in \mathbf{B}} Ae_b^{\oplus d_i(b)}$, $B = \text{End}_A(P)^{\text{op}}$ with f_i being the projection of P onto the i -th summand Pf_i , and $F = P \otimes_B ?$.

Next we observe that $B = \text{End}_A(P)^{\text{op}}$ is isomorphic to an algebra of block matrices, with blocks indexed by the set $I \times \mathbf{B}$, and the block in the row indexed by (i, a) and column indexed by (j, b) being a $d_i(a) \times d_j(b)$ matrix with entries in $e_a A e_b$. The multiplication is just matrix multiplication combined with multiplication in A . From this description, it is clear that B possesses an anti-involution τ defined by taking the transpose of a matrix and applying σ to all of the entries of the result. For $i \in I, b \in \mathbf{B}$ and $1 \leq r \leq d_i(b)$, let $f_{i,b;r} \in B$ be the matrix with all entries equal to zero except for the r -th entry in its (i, b) -th diagonal block, which is equal to e_b . This is a primitive idempotent in B , and it is fixed by τ . This verifies the axiom (σ S1) for this particular anti-involution τ of B . Next we check that the axiom (σ S2) is satisfied. Let $\text{tr} : A \rightarrow \mathbb{k}, x \mapsto (1_A, x)$ be the trace function associated to a σ -symmetric form on A . Define $\text{tr}' : B \rightarrow \mathbb{k}$ by mapping a matrix in B to the sum of the scalars obtained by applying tr to each of its diagonal entries. Then let $(\cdot, \cdot)' : B \times B \rightarrow \mathbb{k}$ be the bilinear form defined from $(x, y)' := \text{tr}'(xy)$. This is a non-degenerate symmetric bilinear form on B with $(\tau(x), \tau(y))' = (x, y)'$.

It is clear that $F \circ ?^{\circledast} \cong ?^{\circledast} \circ F$ since F is isomorphic to the idempotent truncation functor defined by $f := \sum f_{i,b;1}$ summing over all $i \in I, b \in \mathbf{B}$ such that $d_i(b) \neq 0$. We also have that $f_i = \sum_{b \in \mathbf{B}} \sum_{r=1}^{d_i(b)} f_{i,b;r}$, so $\tau(f_i) = f_i$ for each $i \in I$. So we have now proved the existence of an anti-involution τ with all of the desired properties. It remains to note given another other anti-involution $\omega : B \rightarrow B$ with $F \circ ?^{\circledast} \cong ?^{\circledast} \circ F$ that $?^{\circledast} \circ ?^{\circledast} \cong \text{Id}$, hence, we have that $\omega \circ \tau = \gamma$ for some inner automorphism $\gamma : B \rightarrow B, x \mapsto uxu^{-1}$; equivalently, $\omega = \gamma \circ \tau$. If that is the case, then B is also ω -symmetric since the bilinear form $(\cdot, \cdot)'$ constructed in the previous paragraph also satisfies

$$(\omega(x), \omega(y))' = (u\tau(x)u^{-1}, u\tau(y)u^{-1}) = (\tau(x), \tau(y)) = (x, y)'$$

for $x, y \in A$. □

Definition 4.53. Let \mathcal{R} be a fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. We say that a contravariant autoequivalence $?^\vee$ of \mathcal{R} is a *Chevalley duality* if there is a realization (A, σ) of $(\mathcal{R}, ?^\vee)$ in which σ is a *Chevalley anti-involution*, meaning that $\sigma^2 = \text{id}$ and the following two properties hold:

- (Ch1) There exists a set $\{e_a \mid a \in \mathbf{B}\}$ of mutually orthogonal σ -invariant idempotents in A such that $\dim e_a L(b) = \delta_{a,b}$ for all $b \in \mathbf{B}$ with $\rho(b) \not\prec \rho(a)$; here, $L(b)$ is the irreducible A -module labelled by $b \in \mathbf{B}$.
- (Ch2) Let $A_{\leq \lambda}$ be the quotient of A by the two-sided ideal generated by the idempotents $\{e_a \mid a \in \mathbf{B} \text{ with } \rho(a) \not\leq \lambda\}$ and $A_\lambda := \bigoplus_{a,b \in \mathbf{B}_\lambda} \bar{e}_a A_{\leq \lambda} \bar{e}_b$. For each $\lambda \in \Lambda$, we require that A_λ possesses a non-degenerate associative symmetric bilinear form $(\cdot, \cdot)_\lambda$ such that $(\sigma_\lambda(x), \sigma_\lambda(y))_\lambda = (x, y)_\lambda$ for all $x, y \in A_\lambda$, where $\sigma_\lambda : A_\lambda \rightarrow A_\lambda$ is the anti-involution induced by σ .

In view of the following lemma, axiom (Ch2) is vacuous in the case that \mathcal{R} is a highest weight category, since then we have that $A_\lambda = \mathbb{k}$ and $\sigma_\lambda = \text{id}$.

Lemma 4.54. *Suppose that (A, σ) is a realization of $(\mathcal{R}, ?^\vee)$ in which σ is a Chevalley involution as in Definition 4.53. The algebra A_λ from (Ch2) is the basic algebra realizing the stratum \mathcal{R}_λ , and it is σ_λ -symmetric in the sense of Definition 4.51. We also have that $L(b)^\vee \cong L(b)$ for all $b \in \mathbf{B}$, i.e., Chevalley dualities preserve isomorphism classes of irreducible objects.*

Proof. Let I be the two-sided ideal of A generated by $\{e_a \mid a \in \mathbf{B} \text{ with } \rho(a) \not\leq \lambda\}$. We claim that $\mathcal{R}_{\leq \lambda}$ is the full subcategory of \mathcal{R} consisting of all objects V such that $IV = 0$. To see this, if $IV = 0$ then $e_a V = 0$ for all $a \in \mathbf{B}$ with $\rho(a) \not\leq \lambda$ then $[V : L(a)] = 0$ for all such a thanks to axiom (Ch1). So we have that $V \in \mathcal{R}_{\leq \lambda}$. Conversely, if $V \in \mathcal{R}_{\leq \lambda}$ and $\rho(a) \not\leq \lambda$ then the idempotent e_a is zero on every irreducible subquotient of V by (Ch1), hence, $e_a V = 0$. This implies that $IV = 0$.

By the claim, the algebra $A_{\leq \lambda} = A/I$ gives a realization of $\mathcal{R}_{\leq \lambda}$. Let \bar{e}_b denote the image of e_b in $A_{\leq \lambda}$. For $b \in \mathbf{B}_\lambda$, we have that $\bar{e}_b L(c) = \delta_{b,c}$ for all $c \in \mathbf{B}_\lambda$. This shows that the mutually orthogonal idempotents $\{\bar{e}_b \mid b \in \mathbf{B}_\lambda\}$ are primitive in $A_{\leq \lambda}$. Hence, $A_\lambda = \bigoplus_{a,b \in \mathbf{B}_\lambda} \bar{e}_a A_{\leq \lambda} \bar{e}_b$ is the basic algebra realizing the stratum \mathcal{R}_λ . It is immediate from the axioms (Ch1)–(Ch2) and the definition that A_λ is σ_λ -symmetric.

Finally to show that $L(b)^\circledast \cong L(b)$ for all $b \in \mathbf{B}$, suppose that $b \in \mathbf{B}_\lambda$. We have that $e_a L(b)^\circledast \cong (e_a L(b))^* = 0$ for all a with $\rho(a) \not\leq \lambda$, so $L(b)^\circledast \in \mathcal{R}_{\leq \lambda}$. Moreover, $e_b L(b)^\circledast \cong (e_b L(b))^*$ is one-dimensional. Since \bar{e}_b is primitive in $A_{\leq \lambda}$ this implies that $L(b)^\circledast \cong L(b)$. \square

Theorem 4.55 (Chevalley dualities commute with Ringel duality). *Let \mathcal{R} be a fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Assume that \mathcal{R} possesses a Chevalley duality $?^\vee$. Fix also a realization (A, σ) of $(\mathcal{R}, ?^\vee)$ in which σ is a Chevalley involution, and let $T(b)$ denote the left A -module corresponding to $T_+(b) \in \mathcal{R}$.*

- (1) *If \mathcal{R} is tilting-rigid and $\text{char } \mathbb{k} \neq 2$ then for each $b \in \mathbf{B}$ there exists a non-degenerate symmetric bilinear form $\langle \cdot, \cdot \rangle : T(b) \times T(b) \rightarrow \mathbb{k}$ satisfying the following σ -adjunction property:*

$$\langle xv, w \rangle = \langle v, \sigma(x)w \rangle \quad (4.47)$$

for $v, w \in T(b)$ and $x \in A$.

- (2) *Suppose that we are given objects of \mathcal{R} corresponding to A -modules $\{T_b \mid b \in \mathbf{B}\}$ such that each T_b is a direct sum of $T(b)$ and copies of $T(c)$ for $c \in \mathbf{B}$ with $\rho(c) < \rho(b)$. Assume moreover that each T_b is equipped with a non-degenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$ satisfying the σ -adjunction property. Then, \mathcal{R} is tilting-rigid with symmetric strata, and there is an induced Chevalley duality $?^\wedge$ on the Ringel dual \mathcal{R}' of \mathcal{R} satisfying (4.45).*

Proof. (1) Suppose that $b \in \mathbf{B}_\lambda$ for some $\lambda \in \Lambda$. For the purpose of proving (1) for $T(b)$, we can replace \mathcal{R} by $\mathcal{R}_{\leq \lambda}$ and the algebra A realizing \mathcal{R} by the corresponding quotient algebra to assume without loss of generality that $\mathcal{R} = \mathcal{R}_{\leq \lambda}$. So now \mathcal{R} is either finite or upper finite, and the chosen algebra A is either a finite-dimensional algebra or a locally finite-dimensional locally unital algebra, respectively. Let $\{e_a \mid a \in \mathbf{B}\}$ be the mutually orthogonal σ -invariant idempotents given by the axiom (Ch1). Let $e_\lambda := \sum_{b \in \mathbf{B}_\lambda} e_b$ and $A_\lambda := e_\lambda A e_\lambda$. By Lemma 4.54, this is the basic finite-dimensional algebra realizing the top stratum \mathcal{R}_λ , and $\{e_b \mid b \in \mathbf{B}_\lambda\}$ is a set of representatives for the conjugacy classes of primitive idempotents in A_λ . The anti-involution σ of A restricts to an anti-involution σ_λ of A_λ . Also $e_\lambda T(b)$ is isomorphic to the indecomposable projective A_λ -module $A_\lambda e_b$.

Claim 1: *Let $\psi : T(b) \rightarrow T(b)$ be an A -module homomorphism and $\bar{\psi} : e_\lambda T(b) \rightarrow e_\lambda T(b)$ be its restriction, which is an A_λ -module homomorphism. Then ψ is an isomorphism if and only if $\bar{\psi}$ is an isomorphism.* The forward implication is clear. For the converse, let $E := \text{End}_A(T(b))^{\text{op}}$ and $E_\lambda := \text{End}_{A_\lambda}(e_\lambda T(b))^{\text{op}}$. As $e_\lambda T(b)$ is an indecomposable A_λ -module, the algebra E_λ is a finite-dimensional local algebra, so its Jacobson radical

is of codimension one and any non-unit is nilpotent. The algebra E is also a finite-dimensional local algebra in the finite case, while in the upper finite case it is a pseudo-compact topological algebra with Jacobson radical $J(E)$ having codimension one. In either case, any element of E is either a unit or it belongs to $J(E)$. Let \bar{E} be the image of E under the homomorphism $E \rightarrow E_\lambda$ defined by restriction. The Jacobson radical of \bar{E} is the image of $J(E)$, so it is again of codimension one⁹ in \bar{E} . We are given $\psi \in E$ such that $\bar{\psi}$ is a unit in E_λ . This means that $\bar{\psi}$ is not nilpotent, hence, it is also a unit in \bar{E} . It follows that $\bar{\psi} \notin J(\bar{E})$ so $\psi \notin J(E)$. This shows that ψ is a unit in E , i.e., it is an isomorphism as required.

Claim 2: Let $\langle \cdot, \cdot \rangle$ be a bilinear form on $T(b)$ with the σ -adjunction property. Then $\langle \cdot, \cdot \rangle$ is non-degenerate if and only if its restriction $\langle \cdot, \cdot \rangle_\lambda$ to $e_\lambda T(b)$ is non-degenerate. To see this, observe that the form $\langle \cdot, \cdot \rangle$ induces an A -module homomorphism $\theta : T(b) \rightarrow T(b)^\circledast$ with $\theta(v)(w) = \langle v, w \rangle$, and the form is non-degenerate if and only if this induced homomorphism is an isomorphism. Similarly, the restriction $\langle \cdot, \cdot \rangle_\lambda$ induces an A_λ -module homomorphism $\bar{\theta} : e_\lambda T(b) \rightarrow (e_\lambda T(b))^\circledast$, and the restricted form is non-degenerate if and only if $\bar{\theta}$ is an isomorphism. If we identify $(e_\lambda T(b))^\circledast$ with $e_\lambda(T(b)^\circledast)$ in the natural way, we see that $\bar{\theta}$ is the restriction of θ . We are given that \mathcal{R} is tilting-rigid, and its strata are σ_λ -symmetric which implies that they are weakly symmetric, so there is an A -module isomorphism $\phi : T(b)^\circledast \xrightarrow{\sim} T(b)$ according to Theorem 4.50. This restricts to an A_λ -module isomorphism $\bar{\phi} : e_\lambda(T(b)^\circledast) \rightarrow e_\lambda T(b)$. Now Claim 2 is reduced to showing that the A -module homomorphism $\phi \circ \theta : T(b) \rightarrow T(b)$ is an isomorphism if and only if its restriction $\bar{\phi} \circ \bar{\theta} : e_\lambda T(b) \rightarrow e_\lambda T(b)$ is an isomorphism. This follows from Claim 1.

Claim 3: The socle of $A_\lambda e_b$ is irreducible, and any non-zero vector $z_b \in \text{soc}(A_\lambda e_b)$ satisfies $\sigma_\lambda(z_b) = z_b$. By (Ch2), there is a non-degenerate associative symmetric bilinear form $(\cdot, \cdot)_\lambda$ on A_λ with $(\sigma_\lambda(x), \sigma_\lambda(y))_\lambda = (x, y)_\lambda$ for all $x, y \in A_\lambda$. By the discussion before Lemma 4.52, $A_\lambda e_b$ is self-dual, so it has irreducible socle isomorphic to its head. Moreover, $(\cdot, \cdot)_\lambda$ restricts to a non-degenerate associative symmetric bilinear form on $e_b A_\lambda e_b$. This is a local symmetric algebra, so its Jacobson radical J is a two-sided ideal of codimension one and J^\perp is a two-sided ideal of dimension one. Let z_b be a non-zero vector in J^\perp . We must have that $(e_b, z_b)_\lambda \neq 0$ by the non-degeneracy of the form. Moreover, z_b also spans the socle of $A_\lambda e_b$. It remains to show that $\sigma_\lambda(z_b) = z_b$. Since σ_λ leaves J^\perp invariant we certainly have that $\sigma_\lambda(z_b) = cz_b$ for $c \in \mathbb{k}$. Now we use the σ_λ -symmetry of the form:

$$(e_b, z_b)_\lambda = (\sigma_\lambda(e_b), \sigma_\lambda(z_b))_\lambda = (e_b, cz_b)_\lambda.$$

Since $(e_b, z_b)_\lambda \neq 0$, this implies that $c = 1$.

Claim 4: Suppose that $\langle \cdot, \cdot \rangle_\lambda$ is a bilinear form on $A_\lambda e_b$ with the σ_λ -adjunction property, i.e., the analog of (4.47) with σ replaced by σ_λ holds for all $x \in A_\lambda$. This form is non-degenerate if and only if $\langle e_b, z_b \rangle_\lambda \neq 0$ for z_b as in Claim 3. Suppose first that $\langle e_b, z_b \rangle_\lambda \neq 0$. Take any $0 \neq x \in A_\lambda e_b$. Since the socle of $A_\lambda e_b$ is one-dimensional, there exists $y \in A_\lambda$ such that $yx = z_b$. Then $\langle e_b, yx \rangle_\lambda \neq 0$ so $\langle \sigma(y)e_b, x \rangle_\lambda \neq 0$. This shows that the function $A_\lambda e_b \rightarrow (A_\lambda e_b)^*, x \mapsto \langle \cdot, x \rangle_\lambda$ is injective, hence, the form is non-degenerate. Conversely, suppose that $\langle e_b, z_b \rangle_\lambda = 0$. Then the A_λ -submodule $\{x \in A_\lambda e_b \mid \langle x, z_b \rangle_\lambda = 0\}$ contains e_b , hence, it is all of $A_\lambda e_b$. So the form is degenerate.

Now we can complete the proof of (1). As noted in the proof of Claim 2, $T(b) \cong T(b)^\circledast$. Let $[\cdot, \cdot]$ be the bilinear form on $T(b)$ corresponding to such an isomorphism. This form is non-degenerate and has the σ -adjunction property. However, it is not necessarily symmetric, so we symmetrize by letting $\langle \cdot, \cdot \rangle$ be the form on $T(b)$ defined from

$$\langle v, w \rangle := [v, w] + [w, v].$$

⁹In fact, one can show that $\bar{E} = E_\lambda$ but we do not need to use this here.

Using that σ is an involution, it is easy to check that this new form still has the σ -adjunction property, and now it is symmetric, but we do not yet know that it is non-degenerate. To see this, let $\iota : A_\lambda e_b \xrightarrow{\sim} e_\lambda T(b)$ be an A_λ -module isomorphism. Let $[\cdot, \cdot]_\lambda$ and $\langle \cdot, \cdot \rangle_\lambda$ be the bilinear forms on $A_\lambda e_b$ defined from $[x, y]_\lambda := [\iota(x), \iota(y)]$ and $\langle x, y \rangle_\lambda := \langle \iota(x), \iota(y) \rangle$. Applying Claim 2, we see that the form $[\cdot, \cdot]_\lambda$ is non-degenerate, and the goal is to show that $\langle \cdot, \cdot \rangle_\lambda$ is non-degenerate. Applying Claim 4, we have that $[e_b, z_b]_\lambda \neq 0$ and we need to show that $\langle e_b, z_b \rangle_\lambda \neq 0$. This follows since

$$\langle e_b, z_b \rangle_\lambda = [e_b, z_b]_\lambda + [z_b, e_b]_\lambda = [e_b, z_b]_\lambda + [e_b, \sigma_\lambda(z_b)]_\lambda = 2[e_b, z_b]_\lambda \neq 0,$$

using that $\sigma_\lambda(z_b) = z_b$ by Claim 3 together with the hypothesis that $\text{char } \mathbb{k} \neq 2$.

(2) We are given non-degenerate symmetric bilinear forms $\langle \cdot, \cdot \rangle$ on each T_b satisfying the σ -adjunction property. It follows that $T_b \cong T_b^\circledast$. Since $T_+(b)^\vee \cong T_-(b)$ for each $b \in \mathbf{B}$, this is enough to deduce that \mathcal{R} is tilting-rigid. Also the assumption that $?^\vee$ is a Chevalley duality implies that the basic algebra A_λ realizing \mathcal{R}_λ is σ_λ -symmetric, hence, \mathcal{R}_λ is symmetric.

Now the argument proceeds in a similar way to the proof of Theorem 4.50. We just explain the details in the finite case; the other three cases are similar but there are slight notational differences. We may assume that the tilting generator used to define the Ringel dual category is $T = \bigoplus_{b \in \mathbf{B}} T_b$. Then $\mathcal{R}' = B\text{-mod}_{\text{fd}}$ for $B := \text{End}_A(T)^{\text{op}}$. The given forms on each T_b give us a non-degenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$ on T satisfying (4.47), with the summands T_b being mutually orthogonal. Define an anti-automorphism τ of B from the equation $\langle vy, w \rangle = \langle v, w\tau(y) \rangle$ for $v, w \in T$ and $y \in B$. This gives us a contravariant autoequivalence $?^\vee := ?^\circledast$ on \mathcal{R}' , and we get the isomorphisms (4.45) like in the proof of Theorem 4.50.

As $\langle \cdot, \cdot \rangle$ is symmetric and T is a faithful B -module, the following calculation implies that $\tau^2 = \text{id}$:

$$\langle vy, w \rangle = \langle v, w\tau(y) \rangle = \langle w\tau(y), v \rangle = \langle w, v\tau^2(y) \rangle = \langle v\tau^2(y), w \rangle.$$

For each $b \in \mathbf{B}$, let $f_b \in B$ be the idempotent projecting T onto the summand T_b . Using that the restriction of the form $\langle \cdot, \cdot \rangle$ to this summand is non-degenerate, it follows that $\tau(f_b) = f_b$. So $\{f_b \mid b \in \mathbf{B}\}$ is a set of mutually orthogonal τ -invariant idempotents in B . The idempotent f_b is equal to the primitive idempotent projecting T_b onto its summand $T(b)$ plus other orthogonal primitive idempotents which project onto summands $T(a)$ for $a \in \mathbf{B}_{<\rho(b)}$. Bearing in mind we are using the opposite ordering on Λ on the Ringel dual side, this is just what we need for the property (Ch1).

Finally, to see that property (Ch2) holds, let B_λ be the algebra obtained from B according to the construction of (Ch2) and $\tau_\lambda : B_\lambda \rightarrow B_\lambda$ be the anti-involution induced by τ . The pair $(B_\lambda, \tau_\lambda)$ is a realization of $(\mathcal{R}'_\lambda, ?^\wedge)$, where $?^\wedge$ here is the contravariant autoequivalence of \mathcal{R}'_λ induced by the one on \mathcal{R}' . We also have the pair $(A_\lambda, \sigma_\lambda)$ realizing \mathcal{R}_λ with its contravariant autoequivalence induced by $?^\vee$. We know already by Lemma 4.54 that A_λ is σ_λ -symmetric, and (Ch2) follows if we can show that B_λ is τ_λ -symmetric. This follows from Lemma 4.52 since the functor $F_\lambda : A_\lambda\text{-mod}_{\text{fd}} \rightarrow B_\lambda\text{-mod}_{\text{fd}}$ is an equivalence satisfying $F_\lambda \circ ?^\vee \cong ?^\wedge \circ F_\lambda$. Indeed, Theorem 4.42(2) gives that $F_\lambda \cong G_\lambda$, while (4.45) and the definitions (4.35)–(4.36) give that the dualities $?^\vee : \mathcal{R}_\lambda \rightarrow \mathcal{R}_\lambda$ and $?^\wedge : \mathcal{R}'_\lambda \rightarrow \mathcal{R}'_\lambda$ satisfy $G_\lambda \circ ?^\vee \cong ?^\wedge \circ F_\lambda$. \square

5. GENERALIZATIONS OF QUASI-HEREDITARY ALGEBRAS

Now we give some applications of semi-infinite Ringel duality. First, we use it to show that any upper finite highest weight category can be realized as $A\text{-mod}_{\text{fd}}$ for an upper finite based quasi-hereditary algebra A . The latter notion, which is Definition 5.1, already exists in the literature in some equivalent forms. When A is finite-dimensional, it gives an alternative algebraic characterization of the usual notion of quasi-hereditary algebra. Then, in §5.2, we introduce further notions of based ε -stratified algebras and based

ε -quasi-hereditary algebras, which correspond to ε -stratified categories and ε -highest weight categories, respectively. In §5.3, we introduce based stratified algebras and based properly stratified algebras corresponding to fully stratified and fibered highest weight categories, respectively. Finally, in §§5.4–5.5, we discuss the related notions of triangular bases and a triangular decompositions.

5.1. Based quasi-hereditary algebras. The following definition is a simplified version of [ELau, Def. 2.1] translated from the framework of \mathbb{k} -linear categories to that of locally unital algebras. Also, for finite-dimensional algebras, it is equivalent to [KM, Def. 2.4]. These assertions will be explained in more detail in Remarks 5.7–5.8 below.

Definition 5.1. By a *finite* (resp., *upper finite*, resp., *essentially finite*) *based quasi-hereditary algebra*, we mean a finite-dimensional (resp., locally finite-dimensional, resp., essentially finite-dimensional) locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$ with the following additional data:

- (QH1) A subset $\Lambda \subseteq I$ indexing *special idempotents* $\{e_\lambda \mid \lambda \in \Lambda\}$.
 - (QH2) A partial order \leq making the set Λ into a poset which is upper finite in the upper finite case and interval finite in the essentially finite case.
 - (QH3) Sets $Y(i, \lambda) \subset e_i A e_\lambda$, $X(\lambda, j) \subset e_\lambda A e_j$ for $\lambda \in \Lambda$, $i, j \in I$.
- Let $Y(\lambda) := \bigcup_{i \in I} Y(i, \lambda)$ and $X(\lambda) := \bigcup_{j \in I} X(\lambda, j)$. We impose the following axioms:
- (QH4) The products yx for $(y, x) \in \bigcup_{\lambda \in \Lambda} Y(\lambda) \times X(\lambda)$ are a basis for A .
 - (QH5) For $\lambda, \mu \in \Lambda$, the sets $Y(\mu, \lambda)$ and $X(\lambda, \mu)$ are empty unless $\mu \leq \lambda$.
 - (QH6) We have that $Y(\lambda, \lambda) = X(\lambda, \lambda) = \{e_\lambda\}$ for each $\lambda \in \Lambda$.

We say that A is *symmetrically based* if there is also some given algebra anti-involution $\sigma : A \rightarrow A$ with $\sigma(e_i) = e_i$ and $Y(i, \lambda) = \sigma(X(\lambda, i))$ for all $i \in I$, $\lambda \in \Lambda$.

We refer to the given basis for A from (QH4) as the *triangular basis*; it is certainly not unique since one can replace any $Y(i, \lambda)$ or $X(\lambda, j)$ by another basis that spans the same subspace up to “higher terms”. If A is symmetrically based rather than merely based, this basis is a *cellular basis* in the general sense of [GL], [Wes]. However, Definition 5.1 is considerably more restrictive than the general notions of cellular algebra or category introduced in *loc. cit.*. In fact, for finite-dimensional algebras, Definition 5.1 is equivalent to the usual notion of quasi-hereditary algebra, as we will explain more fully below.

Remark 5.2. It is clear from (QH4) that $A = \sum_{\lambda \in \Lambda} A e_\lambda A$. Hence, A is Morita equivalent to the idempotent truncation $\bigoplus_{\lambda, \mu \in \Lambda} e_\lambda A e_\mu$. This means that if one is prepared to pass to a Morita equivalent algebra then one can assume without loss of generality that the sets Λ and I in Definition 5.1 are actually equal, i.e., all distinguished idempotents are special. However, in naturally-occurring examples, one often encounters situations in which the set I is strictly larger than Λ .

Remark 5.3. A classical example of a finite symmetrically based quasi-hereditary algebra is the Schur algebra $S(n, r)$ with its basis of *codeterminants* $\xi_{i, \ell(\lambda)} \xi_{\ell(\lambda), j}$ as constructed by Green in [Gre]; one definitely needs $I \supsetneq \Lambda$ in this example.

Remark 5.4. For a well-known infinite-dimensional example, consider the path algebra A of the Temperley-Lieb category $\mathcal{TL}(\delta)$ for any value of its parameter $\delta \in \mathbb{k}$. The natural diagram basis gives a triangular basis making A into an upper finite symmetrically based quasi-hereditary algebra. For this, one takes $I = \Lambda = \mathbb{N}$ ordered by the opposite of the natural ordering. The set $Y(\lambda)$ (resp., $X(\lambda)$) consists of all cap-free Temperley-Lieb diagrams with λ strings at the bottom (resp., all cup-free Temperley-Lieb diagrams with λ strings at the top). The anti-automorphism σ is defined by reflecting diagrams in a horizontal axis.

Lemma 5.5. *Let A be a finite, essentially finite or upper finite based quasi-hereditary algebra. For $\lambda \in \Lambda$, any element f of the two-sided ideal $A e_\lambda A$ can be written as a linear combination of elements of the form yx for $y \in Y(\mu)$, $x \in X(\mu)$ and $\mu \geq \lambda$.*

Proof. We first consider the upper finite case. By considering the triangular basis, we may assume that $f = y_1 x_1 y_2 x_2$ for $y_1 \in Y(\mu_1), x_1 \in X(\mu_1, \lambda), y_2 \in Y(\lambda, \mu_2), x_2 \in X(\mu_2)$ and $\mu_1, \mu_2 \geq \lambda$. If $\mu_1 = \mu_2 = \lambda$ then $x_1 = e_\lambda = y_2$ and $f = y_1 x_2$, as required. This finished the proof for λ maximal. If $\mu_r > \lambda$ for some $r \in \{1, 2\}$, then we have that $f \in Ae_{\mu_r}A$ for this r , and are done by downward induction on the partial order on Λ .

The finite and essentially finite cases are similar. Now, assuming that $f \in e_i A e_j$ for $i, j \in I$, there are only finitely many $\mu \in \Lambda$ such that $e_i A e_\mu \neq 0$ or $e_\mu A e_j \neq 0$. Letting Λ' be the finite set of all such μ , we can then again proceed by downward induction on the partial order on Λ' . \square

Corollary 5.6. *Let Λ^\dagger be an upper set in Λ . The two-sided ideal J_{Λ^\dagger} of A generated by $\{e_\lambda \mid \lambda \in \Lambda^\dagger\}$ has basis $\{yx \mid (y, x) \in \bigcup_{\lambda \in \Lambda^\dagger} Y(\lambda) \times X(\lambda)\}$.*

Proof. Let J be the subspace of A with basis given by the products yx for $y \in Y(\lambda), x \in X(\lambda)$ and $\lambda \in \Lambda^\dagger$. For any such element $yx \in J$, we have that $yx = ye_\lambda x$, hence, $yx \in J_{\Lambda^\dagger}$. This shows that $J \subseteq J_{\Lambda^\dagger}$. Conversely, any element of J_{Λ^\dagger} is a linear combination of elements of $Ae_\lambda A$ for $\lambda \in \Lambda^\dagger$. In turn, Lemma 5.5 shows that any element of $Ae_\lambda A$ for $\lambda \in \Lambda^\dagger$ is a linear combination of elements yx for $y \in Y(\mu), x \in X(\mu)$ and $\mu \geq \lambda$. All of these elements yx belong to J because Λ^\dagger is an upper set; thus $J_{\Lambda^\dagger} \subseteq J$. \square

Remark 5.7. We have formulated Definition 5.1 only for algebras over our usual ground field \mathbb{k} , but the definition makes sense with \mathbb{k} replaced by some more general commutative ground ring R (“finite-dimensional” being interpreted as “free of finite rank”). Then, in the symmetrically based upper finite case, Definition 5.1 is equivalent to the notion of an object-adapted cellular category from [ELau, Def. 2.1]. This can be seen from Corollary 5.6 and [ELau, Lemmas 2.6-2.8]. Elias and Lauda also note in *loc. cit.* that the diagrammatic Hecke category $\mathcal{H}_{BS}(W, S)$ of [EW] associated to a Coxeter system (W, S) is an example of an object-adapted cellular category. In our language, the path algebra H of $\mathcal{H}_{BS}(W, S)$ is an upper finite symmetrically based quasi-hereditary algebra defined over the ground ring $R = \mathbb{Q}[\mathfrak{h}]$, that is, the ring of regular functions arising from a realization \mathfrak{h} of (W, S) . A cellular basis is given by the double light leaves basis. (One needs some assumptions on the realization as in [EW] for this basis to be defined.)

Remark 5.8. In the finite case, Definition 5.1 is equivalent to the notion of based quasi-hereditary algebra from [KM, Def. 2.4]. To see this, one takes the set Λ indexing the special idempotents in our setup to be the set I from *loc. cit.* (which indexes mutually orthogonal idempotents $e_i \in A$ according to [KM, Lem. 2.8]). Then we take our set I to be the set $\Lambda \sqcup \{0\}$, i.e., we add one more element indexing one more idempotent $e_0 := 1_A - \sum_{\lambda \in \Lambda} e_\lambda$. Kleshchev and Muth established the equivalence of their notion of based quasi-hereditary algebra with the original notion of quasi-hereditary algebra from [CPS1] (providing the partial order on Λ is actually a total order); for ground fields, we will reprove this equivalence in a different way below. See also [DuR] which established a similar result using a related notion of standardly based algebra.

Let A be a based quasi-hereditary algebra as in Definition 5.1. For $\lambda \in \Lambda$, let $A_{\leq \lambda}$ be the quotient of A by the two-sided ideal generated by the idempotents e_μ for $\mu \not\leq \lambda$. For $x \in A$, we often write simply \bar{x} for the image of x in $A_{\leq \lambda}$. Corollary 5.6 implies that

$$A_{\leq \lambda} = \bigoplus_{i, j \in I} \bar{e}_i A_{\leq \lambda} \bar{e}_j \quad (5.1)$$

is based quasi-hereditary in its own right, with special idempotents indexed by elements of the lower set $(-\infty, \lambda]$ and basis given by the products $\bar{y}\bar{x}$ for $y \in Y(\mu), x \in X(\mu)$ and $\mu \in (-\infty, \lambda]$. Define the *standard* and *costandard modules* associated to $\lambda \in \Lambda$ by

$$\Delta(\lambda) := A_{\leq \lambda} \bar{e}_\lambda \quad \nabla(\lambda) = (\bar{e}_\lambda A_{\leq \lambda})^{\otimes} \quad (5.2)$$

These are left A -modules which are projective and injective as $A_{\leq \lambda}$ -modules, respectively. In the finite or essentially finite case, $\bar{e}_\lambda A_{\leq \lambda}$ is finite-dimensional and one could

just take the full linear dual in (5.2), but in general in the upper finite case $\Delta(\lambda)$ and $\nabla(\lambda)$ are only locally finite-dimensional. The modules $\Delta(\lambda)$ may also be called *cell modules* and the modules $\nabla(\lambda)$ *dual cell modules*. The vectors $\{y\bar{e}_\lambda \mid y \in Y(\lambda)\}$ give the *standard basis* for $\Delta(\lambda)$. Similarly, the vectors $\{\bar{e}_\lambda x \mid x \in X(\lambda)\}$ are a basis for the right A -module $\bar{e}_\lambda A$; the dual basis to this is the *costandard basis* for $\nabla(\lambda)$.

Theorem 5.9 (Highest weight categories from based quasi-hereditary algebras). *Let A be a finite (resp., upper finite, resp., essentially finite) based quasi-hereditary algebra. The modules*

$$\{L(\lambda) := \text{hd } \Delta(\lambda) \cong \text{soc } \nabla(\lambda) \mid \lambda \in \Lambda\}$$

give a complete set of pairwise inequivalent irreducible left A -modules. Moreover, the category $\mathcal{R} := A\text{-mod}_{\text{fd}}$ (resp., $\mathcal{R} := A\text{-mod}_{\text{fd}}$, resp., $\mathcal{R} := A\text{-mod}_{\text{fd}}$) is a finite (resp., upper finite, resp., essentially finite) highest weight category with the given weight poset (Λ, \leq) . Its standard and costandard objects $\Delta(\lambda)$ and $\nabla(\lambda)$ are as defined by (5.2). If A is symmetrically based with anti-involution σ then ${}^? \circledast : \mathcal{R} \rightarrow \mathcal{R}$ is Chevalley duality of \mathcal{R} in the sense of Definition 4.53.

Proof. For $\lambda \in \Lambda$, let P_λ be the left ideal Ae_λ . We start by establishing the claim that P_λ has a Δ -flag with $\Delta(\lambda)$ at the top and other sections of the form $\Delta(\mu)$ for $\mu > \lambda$. To prove this, fix some λ and set $P := P_\lambda$ for short. This module has basis $\{yx \mid (y, x) \in \bigcup_{\mu \geq \lambda} Y(\mu) \times X(\mu, \lambda)\}$. Let $\{\mu_1, \dots, \mu_n\}$ be the finite set $\{\mu \in [\lambda, \infty) \mid X(\mu, \lambda) \neq \emptyset\}$ ordered so that $\mu_r \leq \mu_s \Rightarrow r \leq s$; in particular, $\mu_1 = \lambda$. For $1 \leq r \leq n$ let P_r be the subspace of P spanned by $\{yx \mid (y, x) \in \bigcup_{s=r+1}^n Y(\mu_s) \times X(\mu_s, \lambda)\}$. They define a filtration $P =: P_0 > P_1 > \dots > P_n = 0$, since each P_r is a A -submodule of P . Moreover, there is, for each $0 \leq r \leq n$ an A -module isomorphism

$$\theta_r : \bigoplus_{x \in X(\mu_r, \lambda)} \Delta(\mu_r) \xrightarrow{\sim} P_{r-1}/P_r \quad (5.3)$$

which in case $r \geq 1$ sends the basis vector $y\bar{e}_{\mu_r}$ ($y \in Y(\mu_r)$) in the x th copy of $\Delta(\mu_r)$ to $yx + P_r \in P_{r-1}/P_r$. This defines clearly a linear isomorphism, so we just need to check that it is an A -module homomorphism. For this take $y \in Y(j, \mu_r)$ and $u \in e_i Ae_j$. Expand uy in terms of the triangular basis as $\sum_p c_p y_p + \sum_q c'_q y'_q x'_q$ for scalars $c_p, c'_q, y_p \in Y(i, \mu_r)$, $y'_q \in Y(i, \nu_q)$, $x'_q \in X(\nu_q, \mu_r)$ and $\nu_q > \mu_r$. Then we have that $uy\bar{e}_{\mu_r} = \sum_p c_p y_p \bar{e}_{\mu_r}$ and $uyx + P_r = \sum_p c_p y_p x + P_r$, since the “higher terms” $y'_q x'_q$ act as zero on both \bar{e}_{μ_r} and $x + P_r$. This shows that θ_r intertwines the actions of u and so the claim follows, since $P_0/P_1 \cong \Delta(\lambda)$ by construction.

Now we can classify the irreducible A -modules. The first step for this is to show that $\Delta(\lambda)$ has a unique irreducible quotient. To see this, note that the “weight space” $e_\lambda \Delta(\lambda)$ is one-dimensional with basis \bar{e}_λ , due to the fact that $Y(\lambda, \lambda) = \{e_\lambda\}$. This is a cyclic vector, so any proper submodule of $\Delta(\lambda)$ must intersect $e_\lambda \Delta(\lambda)$ trivially. It follows that the sum of all proper submodules is proper, so $\Delta(\lambda)$ has a unique irreducible quotient $L(\lambda)$. Since $e_\lambda L(\lambda)$ is one-dimensional and all other μ with $e_\mu L(\lambda) \neq 0$ satisfy $\mu < \lambda$, the modules $\{L(\lambda) \mid \lambda \in \Lambda\}$ are pairwise inequivalent. To see that they give a full set of irreducible A -modules, let L be any irreducible A -module. In view of Remark 5.2, there exists $\lambda \in \Lambda$ such that $e_\lambda L \neq 0$. Then L is a quotient of $P_\lambda = Ae_\lambda$. By the claim we proved already, it follows that L is a quotient of $\Delta(\mu)$ for some $\mu \geq \lambda$, i.e., $L \cong L(\mu)$.

Thus, we have shown that the modules $\{L(\lambda) \mid \lambda \in \Lambda\}$ give a full set of pairwise inequivalent irreducible left A -modules. Now consider the stratification of \mathcal{R} arising from the given partial order on the index set Λ . In the recollement situation of (3.4), the Serre subcategory $\mathcal{R}_{\leq \lambda}$ (resp., $\mathcal{R}_{< \lambda}$) may be identified with $A_{\leq \lambda}\text{-mod}_{\text{fd}}$ (resp., $A_{< \lambda}\text{-mod}_{\text{fd}}$), and the Serre quotient $\mathcal{R}_\lambda = \mathcal{R}_{\leq \lambda}/\mathcal{R}_{< \lambda}$ is $A_\lambda\text{-mod}_{\text{fd}}$ where $A_\lambda := \bar{e}_\lambda A_{\leq \lambda} \bar{e}_\lambda$. The algebra A_λ has basis \bar{e}_λ , i.e., it is a copy of the ground field \mathbb{k} . This shows that all strata are simple in the sense of Lemma 3.4. Moreover, the standard and costandard objects in the general sense of (1.1) are obtained by applying the standardization functor

$j_i^\lambda := A_{\leq \lambda} \bar{e}_\lambda \otimes_{A_\lambda} ?$ and the costandardization functor $j_*^\lambda := \bigoplus_{i \in I} \text{Hom}_{A_\lambda}(\bar{e}_\lambda A_{\leq \lambda} \bar{e}_i, ?)$ to the irreducible A_λ -module $A_\lambda = \mathbb{k} \bar{e}_\lambda$. Clearly, the resulting modules are isomorphic to $\Delta(\lambda)$ and $\nabla(\lambda)$ as defined by (5.2). The axiom $(\widehat{P}\Delta)$ follows from the claim.

For the final statement about Chevalley duality, the observations made earlier in the proof establish property (Ch1) from Definition 4.53, and (Ch2) is vacuous as we are in the highest weight setting. Hence, σ is a Chevalley anti-involution. \square

Finally in this subsection we are going to prove a converse to Theorem 5.9. This will be deduced from the next theorem together with an application of Ringel duality. In fact, the next theorem is a reformulation of the main result of [AST].

Theorem 5.10 (Based quasi-hereditary algebras from highest weight categories). *Let \mathcal{R} be a finite (resp., lower finite, resp., tilting-bounded essentially finite) highest weight category with weight poset (Λ, \leq) and labelling function L . Suppose we are given $\Lambda \subseteq I$ and a tilting generator $T = \bigoplus_{i \in I} T_i$ for \mathcal{R} such that each T_λ ($\lambda \in \Lambda$) is a direct sum of $T(\lambda)$ and other $T(\mu)$ for $\mu < \lambda$. Let*

$$A := \left(\bigoplus_{i, j \in I} \text{Hom}_{\mathcal{R}}(T_i, T_j) \right)^{\text{op}}.$$

(1) *For $i, j \in I$ and $\lambda \in \Lambda$, pick morphisms*

$$Y(i, \lambda) \subset \text{Hom}_{\mathcal{R}}(T_i, T_\lambda), \quad X(\lambda, j) \subset \text{Hom}_{\mathcal{R}}(T_\lambda, T_j)$$

lifting bases for $\text{Hom}_{\mathcal{R}}(T_i, \nabla(\lambda))$ and $\text{Hom}_{\mathcal{R}}(\Delta(\lambda), T_j)$ as in Corollary 4.44, such that $Y(\lambda, \lambda) = X(\lambda, \lambda) = \{\text{id}_{T_\lambda}\}$. Then $\{yx \mid (y, x) \in \bigcup_{i, j \in I} \bigcup_{\lambda \in \Lambda} Y(i, \lambda) \times X(\lambda, j)\}$ is a triangular basis making A into a finite (resp., upper finite, resp., essentially finite) based quasi-hereditary algebra with respect to the opposite poset (Λ, \geq) .

(2) *If in addition \mathcal{R} has a Chevalley duality $?^\vee$ and, in a suitable realization, the modules corresponding to each T_i possess non-degenerate symmetric bilinear forms satisfying the adjunction property as in (4.47), then the triangular basis in (1) can be chosen so that A is symmetrically based.*

Proof. (1) We have all of the necessary data in place to have a based quasi-hereditary algebra, taking $e_i := \text{id}_{T_i}$ in the obvious way. To check the axioms, Corollary 4.44 checks (QH4), and we have chosen the lifts so that $Y(\lambda, \lambda) = \{e_\lambda\} = X(\lambda, \lambda)$ as in (QH6). For (QH5), note that $Y(\mu, \lambda)$ and $X(\lambda, \mu)$ are empty unless $\mu \geq \lambda$ because $\text{Hom}_{\mathcal{R}}(T_\mu, \nabla(\lambda))$ and $\text{Hom}_{\mathcal{R}}(\Delta(\lambda), T_\mu)$ are zero unless $\lambda \leq \mu$.

(2) Suppose that we are working in a particular algebra realization (B, τ) of $(\mathcal{R}, ?^\vee)$ in which τ is a Chevalley anti-involution and each T_i admits a non-degenerate symmetric bilinear form with the τ -adjunction property. Let $T := \bigoplus_{i \in I} T_i$ and $\langle \cdot, \cdot \rangle : T \times T \rightarrow \mathbb{k}$ be the orthogonal sum of the given forms. Then we obtain an algebra anti-involution $\sigma : A \rightarrow A$ such that $\langle vx, w \rangle = \langle c, w\sigma(x) \rangle$ for all $v, w \in T$, $x \in A$; cf. the proof of Theorem 4.55(2). This fixes each of the idempotents $e_i \in A$. The bilinear form on T_i induces a B -module isomorphism $\phi_i : T_i \xrightarrow{\sim} T_i^\oplus$. Also let $\pi_\lambda : T_\lambda \twoheadrightarrow \nabla(\lambda)$ be some choice of epimorphism for each $\lambda \in \Lambda$ as needed for Corollary 4.44. Then define the embeddings $\iota_\lambda : \Delta(\lambda) \hookrightarrow T_\lambda$ there so that there are induced isomorphisms $\Delta(\lambda) \xrightarrow{\sim} \nabla(\lambda)^\oplus$ making the following diagrams commute for all $\lambda \in \Lambda$:

$$\begin{array}{ccc} \Delta(\lambda) & \xrightarrow{\sim} & \nabla(\lambda)^\oplus \\ \downarrow \iota_\lambda & & \downarrow \pi_\lambda^* \\ T_\lambda & \xrightarrow{\phi_\lambda} & T_\lambda^\oplus \end{array}$$

Now we pick the sets $X(\lambda, i)$ lifting bases for $\text{Hom}_B(\Delta(\lambda), T_j)$ as in Corollary 4.44. Then define $Y(i, \lambda) := \{\phi_\lambda^{-1} \circ x^* \circ \phi_i \mid x \in X(\lambda, i)\}$. This set lifts a basis for $\text{Hom}_B(T_i, \nabla(\lambda))$

as stipulated in Corollary 4.44. Using these choices, the construction from the previous paragraph makes A into a based quasi-hereditary algebra. Moreover, we now have that $Y(i, \lambda) = \sigma(X(\lambda, i))$ for all i, λ , so A is symmetrically based with the underlying anti-involution σ . \square

Corollary 5.11 (Quasi-hereditary algebras are based quasi-hereditary). *Let*

$$A = \bigoplus_{i,j \in I} e_i A e_j$$

be an algebra realization of a finite (resp., upper finite, resp., tilting-bounded essentially finite) highest weight category \mathcal{R} , with weight poset (Λ, \leq) and labelling function L .

- (1) *There is an idempotent expansion $A = \bigoplus_{i,j \in \hat{I}} \hat{e}_i A \hat{e}_j$ of A with $\Lambda \subseteq \hat{I}$, and subsets*

$$Y(i, \lambda) \subset \hat{e}_i A \hat{e}_\lambda, \quad X(\lambda, j) \subset \hat{e}_\lambda A \hat{e}_j$$

for all $\lambda \in \Lambda$ and $i, j \in \hat{I}$ making A into a finite (resp., upper finite, resp., essentially finite) based quasi-hereditary algebra with respect to the given ordering on Λ .

- (2) *If $\text{char } \mathbb{k} \neq 2$ and \mathcal{R} has a Chevalley duality $?^\vee$ then the choices in (1) can be made so that A is symmetrically based with anti-involution σ realizing $?^\vee$.*

Proof. (1) Let $A = \bigoplus_{i,j \in \hat{I}} \hat{e}_i A \hat{e}_j$ be an idempotent expansion indexed by a set \hat{I} chosen so that $\Lambda \subseteq \hat{I}$ and $\text{hd}(A \hat{e}_\lambda) \cong L(\lambda)$ for each $\lambda \in \Lambda$. We are going to apply the Ringel duality from Definition 4.9 (resp., Definition 4.26, resp., Definition 4.33). In the finite or upper finite cases, we fix a choice of tilting generator T for \mathcal{R} and let $B := \text{End}_{\mathcal{R}}(T)^{\text{op}}$. In the essentially finite case, we fix a tilting generator $T = \bigoplus_{j \in J} T_j$ for \mathcal{R} then let $B := \left(\bigoplus_{i,j \in J} \text{Hom}_{\mathcal{R}}(T_i, T_j) \right)^{\text{op}}$. Then in all cases the category $\mathcal{R}' := B\text{-mod}_{\text{fd}}$ is the Ringel dual of the original category. It is a finite (resp., lower finite, resp., tilting-bounded essentially finite) highest weight category with irreducible objects denoted $\{L'(\lambda) | \lambda \in \Lambda\}$ and weight poset (Λ, \geq) . Let $T'_i := (\hat{e}_i T)^* \in \mathcal{R}'$. By Corollary 4.11 (resp., Corollary 4.30, resp., Corollary 4.35), $T' = \bigoplus_{i \in \hat{I}} T'_i$ is a tilting generator for \mathcal{R}' such that the original algebra $A = \bigoplus_{i,j \in \hat{I}} \hat{e}_i A \hat{e}_j$ is isomorphic as a locally unital algebra to $\left(\bigoplus_{i,j \in \hat{I}} \text{Hom}_{\mathcal{R}'}(T'_i, T'_j) \right)^{\text{op}}$. Moreover, T'_λ is the indecomposable tilting module $T'(\lambda)$ for each $\lambda \in \Lambda$. To make A into a based quasi-hereditary algebra, it remains to apply Theorem 5.10(1) with \mathcal{R} , (Λ, \leq) and T_i replaced by \mathcal{R}' , (Λ, \geq) and T'_i in the present setup.

(2) Assume that \mathcal{R} has a Chevalley duality $?^\vee$. Then the category \mathcal{R}' admits a Chevalley duality $?^\wedge$ such that the Ringel duality functors intertwine $?^\vee$ and $?^\wedge$ as in (4.45). This follows by Theorem 4.55, using the assumption that $\text{char } \mathbb{k} \neq 2$ and part (1) of the theorem to establish the existence of suitable bilinear forms as in part (2). Hence, \mathcal{R}' has a realization (B, τ) with τ being a Chevalley involution realizing $?^\wedge$. Then we can appeal to Theorem 5.10(2), again using Theorem 4.55(1) to obtain suitable bilinear forms on each T'_i , to deduce that the triangular basis can be chosen so that A is symmetrically based. In particular, this gives an anti-involution $\sigma : A \rightarrow A$ fixing each \hat{e}_i . It remains to note that $?^\circledast$ realizes $?^\vee$. It suffices to check this on finitely generated projectives when it follows from (4.45) (applied twice since we have used Ringel duality twice). \square

In the finite case, Corollary 5.11 recovers [KM, Prop. 3.5] (but note that the result in *loc. cit.* is also valid over more general ground rings).

5.2. Based ε -stratified and ε -quasi-hereditary algebras. In this subsection, we upgrade the results of §5.1 (excluding any that involve Chevalley duality) to ε -stratified and ε -highest weight categories. The main new definition is as follows.

Definition 5.12. By a *finite* (resp., *upper finite*, resp., *essentially finite*) based ε -stratified algebra, we mean a finite-dimensional (resp., locally finite-dimensional, resp., essentially finite-dimensional) locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$ with the following additional data:

- (ε S1) A subset $\mathbf{B} \subseteq I$ indexing the *special idempotents* $\{e_b \mid b \in \mathbf{B}\}$.
- (ε S2) A poset (Λ, \leq) which is upper finite in the upper finite case and interval finite in the essentially finite case, such that $\Lambda \cap I = \emptyset$.
- (ε S3) A sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$.
- (ε S4) A function $\rho : \mathbf{B} \rightarrow \Lambda$ with finite fibers $\mathbf{B}_\lambda := \rho^{-1}(\lambda)$.
- (ε S5) Sets $Y(i, b) \subset e_i A e_b$ and $X(b, j) \subset e_b A e_j$ for all $b \in \mathbf{B}$ and $i, j \in I$.

Let $Y(b) := \bigcup_{i \in I} Y(i, b)$ and $X(b) := \bigcup_{j \in I} X(b, j)$. There are then four axioms, the first three of which are as follows:

- (ε S6) The products yx for $(y, x) \in \bigcup_{b \in \mathbf{B}} Y(b) \times X(b)$ are a basis for A .
- (ε S7) For $a, b \in \mathbf{B}$, the sets $Y(b, a)$ and $X(a, b)$ are empty unless $\rho(b) \leq \rho(a)$.
- (ε S8) The following hold for all $\lambda \in \Lambda$ and $a, b \in \mathbf{B}_\lambda$:
 - if $\varepsilon(\lambda) = -$ then $Y(a, a) = \{e_a\}$ and $Y(a, b) = \emptyset$ for $a \neq b$;
 - if $\varepsilon(\lambda) = +$ then $X(a, a) = \{e_a\}$ and $X(a, b) = \emptyset$ for $a \neq b$.

To formulate the fourth axiom, let $e_\lambda := \sum_{b \in \mathbf{B}_\lambda} e_b$ for short¹⁰ let $A_{\leq \lambda}$ be the quotient of A by the two-sided ideal generated by $\{e_\mu \mid \mu \not\leq \lambda\}$, and set $A_\lambda := \bar{e}_\lambda A_{\leq \lambda} \bar{e}_\lambda$ (where $\bar{x} \in A_{\leq \lambda}$ denotes the image of $x \in A$ as usual). Then:

- (ε S9) For each $\lambda \in \Lambda$, the finite-dimensional algebra A_λ is basic and $\bar{e}_\lambda = \sum_{b \in \mathbf{B}_\lambda} \bar{e}_b$ is a decomposition of its identity element into mutually orthogonal primitive idempotents.

Definition 5.12 in the special case that the stratification function ρ is a bijection deserves its own name:

Definition 5.13. A *finite* (resp., *upper finite*, resp., *essentially finite*) based ε -quasi-hereditary algebra is a finite-dimensional (resp., locally finite-dimensional, resp., essentially finite-dimensional) locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$ with the following additional data:

- (ε QH1) A subset $\Lambda \subseteq I$ indexing the *special idempotents* $\{e_\lambda \mid \lambda \in \Lambda\}$.
 - (ε QH2) A partial order \leq making the set Λ into a poset which is interval finite in the essentially finite case and upper finite in the upper finite case.
 - (ε QH3) A sign function $\varepsilon : \Lambda \rightarrow \{\pm\}$.
 - (ε QH4) Sets $Y(i, \lambda) \subset e_i A e_\lambda$, $X(\lambda, j) \subset e_\lambda A e_j$ for $i, j \in I$ and $\lambda \in \Lambda$.
- Let $Y(\lambda) := \bigcup_{i \in I} Y(i, \lambda)$ and $X(\lambda) := \bigcup_{j \in I} X(\lambda, j)$. The axioms are as follows:
- (ε QH5) The products yx for $(y, x) \in \bigcup_{\lambda \in \Lambda} Y(\lambda) \times X(\lambda)$ are a basis for A .
 - (ε QH6) For $\lambda, \mu \in \Lambda$, the sets $Y(\mu, \lambda)$ and $X(\lambda, \mu)$ are empty unless $\mu \leq \lambda$.
 - (ε QH7) If $\varepsilon(\lambda) = -$ then $Y(\lambda, \lambda) = \{e_\lambda\}$, and if $\varepsilon(\lambda) = +$ then $X(\lambda, \lambda) = \{e_\lambda\}$.
 - (ε QH8) For each $\lambda \in \Lambda$, the finite-dimensional algebra A_λ as defined in Definition 5.12 is basic and local.

From now on, we just formulate the results for based ε -stratified algebras, since based ε -quasi-hereditary algebras are a special case. The development below parallels the treatment in the previous subsection, but there are some additional subtleties.

Remark 5.2 remains true: one can always pass to a Morita equivalent algebra in which all of the distinguished idempotents are special. The analog of Lemma 5.5 is as follows.

Lemma 5.14. *Let A be a finite, essentially finite or upper finite based ε -stratified algebra. For $\lambda \in \Lambda$, any element f of the two-sided ideal $A e_\lambda A$ can be written as a linear combination of elements of the form yx for $y \in Y(a)$, $x \in X(a)$ and $a \in \mathbf{B}_{\geq \lambda}$.*

¹⁰This notation is unambiguous due to the assumption $\Lambda \cap I = \emptyset$.

Proof. This is similar to the proof of Lemma 5.5. We just explain in the upper finite case. We may assume that $f = y_1 x_1 y_2 x_2$ for $y_1 \in Y(a_1), x_1 \in X(a_1, b), y_2 \in Y(b, a_2), x_2 \in X(a_2), b \in \mathbf{B}_\lambda$ and $a_1, a_2 \in \mathbf{B}_{\geq \lambda}$. If $a_1 \in \mathbf{B}_{> \lambda}$ or $a_2 \in \mathbf{B}_{> \lambda}$, we are done by induction. If $a_1, a_2 \in \mathbf{B}_\lambda$, there are two cases according to whether $\varepsilon(\lambda) = +$ or $\varepsilon(\lambda) = -$. The arguments for these are similar, so we just go through the former case when $\varepsilon(\lambda) = +$. Then we have that $a_1 = b$ and $x_1 = e_b$. Hence $f = y_1 y_2 x_2$. Then we use the basis again to expand $y_1 y_2$ as a linear combination of terms $y_3 x_3$ for $y_3 \in Y(a_3), x_3 \in X(a_3, a_2)$ and $a_3 \in \mathbf{B}_{\geq \lambda}$. If $a_3 \in \mathbf{B}_\lambda$ then we get that $a_3 = a_2$ and $x_3 = e_{a_2}$, so $y_3 x_3 x_2 = y_3 x_2$ as required. If $a_3 \in \mathbf{B}_{> \lambda}$, we can then rewrite $y_3 x_3 x_2$ in the desired form by induction. \square

Corollary 5.15. *Let Λ^\dagger be an upper set in Λ and $\mathbf{B}^\dagger := \rho^{-1}(\Lambda^\dagger)$. The two-sided ideal J_{Λ^\dagger} of A generated by $\{e_\lambda \mid \lambda \in \Lambda^\dagger\}$ has basis $\{yx \mid (y, x) \in \bigcup_{b \in \mathbf{B}^\dagger} Y(b) \times X(b)\}$.*

Let A be a based ε -stratified algebra as in Definition 5.12. For $\lambda \in \Lambda$, Corollary 5.15 implies that $A_{\leq \lambda}$ has basis $\{\bar{y}\bar{x} \mid y \in Y(b), x \in X(b) \text{ and } b \in \mathbf{B}_{\leq \lambda}\}$. Hence, the basic algebra $A_\lambda = \bar{e}_\lambda A_{\leq \lambda} \bar{e}_\lambda$ has basis

$$\{\bar{y} \mid y \in \bigcup_{a, b \in \mathbf{B}_\lambda} Y(a, b)\} \quad \text{if } \varepsilon(\lambda) = +, \quad \{\bar{x} \mid x \in \bigcup_{a, b \in \mathbf{B}_\lambda} X(a, b)\} \quad \text{if } \varepsilon(\lambda) = -.$$

Let $j^\lambda : A_{\leq \lambda}\text{-mod}_{\text{fd}} \rightarrow A_\lambda\text{-mod}_{\text{fd}}, V \mapsto \bar{e}_\lambda V$ be the quotient functor $V \mapsto \bar{e}_\lambda V$, then define the standardization and costandardization functors

$$j_!^\lambda := A_{\leq \lambda} \bar{e}_\lambda \otimes_{A_\lambda} ?, \quad j_*^\lambda := \bigoplus_{i \in I} \text{Hom}_{A_\lambda}(\bar{e}_\lambda A_{\leq \lambda} \bar{e}_i, ?), \quad (5.4)$$

which are left and right adjoints of j^λ , respectively.

Lemma 5.16. *If $\lambda \in \Lambda$ has $\varepsilon(\lambda) = -$ then the standardization functor $j_!^\lambda$ is exact.*

Proof. There is an isomorphism of right A_λ -modules $\bigoplus_{a \in \mathbf{B}_\lambda} \bigoplus_{y \in Y(a)} \bar{e}_a A_\lambda \xrightarrow{\sim} A_{\leq \lambda} \bar{e}_\lambda$ sending the vector \bar{e}_a in the y th copy of $\bar{e}_a A_\lambda$ to $\bar{y} \in A_{\leq \lambda} \bar{e}_\lambda$. To see this, note as $\varepsilon(\lambda) = -$ that the projective A_λ -module $\bar{e}_a A_\lambda$ has basis $\{\bar{x} \mid x \in \bigcup_{b \in \mathbf{B}_\lambda} X(a, b)\}$, and $A_{\leq \lambda} \bar{e}_\lambda$ has basis $\{\bar{y}\bar{x} \mid (y, x) \in \bigcup_{a, b \in \mathbf{B}_\lambda} Y(a) \times X(a, b)\}$. Hence, $A_{\leq \lambda} \bar{e}_\lambda$ is a projective right A_λ -module, and the exactness follows. \square

Continuing with A being a based ε -stratified algebra, we let

$$P_\lambda(b) := A_\lambda \bar{e}_b, \quad I_\lambda(b) := (\bar{e}_b A_\lambda)^\otimes, \quad L_\lambda(b) := \text{hd } P_\lambda(b) \cong \text{soc } I_\lambda(b) \quad (5.5)$$

for $b \in \mathbf{B}_\lambda$. These give full sets of indecomposable projective, indecomposable injective, and irreducible A_λ -modules, respectively. Then we define standard, proper standard, costandard and proper costandard modules

$$\Delta(b) := A_{\leq \lambda} \bar{e}_b \cong j_!^\lambda P_\lambda(b), \quad \bar{\Delta}(b) := j_!^\lambda L_\lambda(b), \quad (5.6)$$

$$\nabla(b) := (\bar{e}_b A_{\leq \lambda})^\otimes \cong j_*^\lambda I_\lambda(b), \quad \bar{\nabla}(b) := j_*^\lambda L_\lambda(b), \quad (5.7)$$

cf. (1.1). Adopt the shorthands $\Delta_\varepsilon(b)$ and $\nabla_\varepsilon(b)$ from (1.2) too. The module $\Delta_\varepsilon(b)$ has a standard basis indexed by the set $Y(b)$. In the case that $\varepsilon(\lambda) = +$, when $\Delta_\varepsilon(b) = \Delta(b)$, this basis is $\{y\bar{e}_b \mid y \in Y(b)\}$. In the case that $\varepsilon(\lambda) = -$, when $\Delta_\varepsilon(b) = \bar{\Delta}(b)$, let \tilde{e}_b be the canonical image of \bar{e}_b under the natural quotient map $\Delta(b) \twoheadrightarrow \bar{\Delta}(b)$. Then the basis is $\{y\tilde{e}_b \mid y \in Y(b)\}$. (One can also construct a costandard basis for $\nabla_\varepsilon(b)$ indexed by $X(b)$ by taking a certain dual basis, but we will not need this here.)

Theorem 5.17 (ε -Highest weight categories from based ε -stratified algebras). *Let A be a finite (resp., upper finite, resp., essentially finite) based ε -stratified algebra as above. The modules*

$$\{L(b) := \text{hd } \Delta_\varepsilon(b) \cong \text{soc } \nabla_\varepsilon(b) \mid b \in \mathbf{B}\}$$

give a complete set of pairwise inequivalent irreducible left A -modules. Moreover, $\mathcal{R} := A\text{-mod}_{\text{fd}}$ (resp., $\mathcal{R} := A\text{-mod}_{\text{fd}}$, resp., $\mathcal{R} := A\text{-mod}_{\text{fd}}$) is a finite (resp., upper finite, resp., essentially finite) ε -stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Its strata

may be identified with the categories $\mathcal{R}_\lambda := A_\lambda\text{-mod}_{\text{fd}}$ with standardization and costandardization functors as in (5.4).

Proof. For $b \in \mathbf{B}$, let P_b be the left ideal Ae_b . We claim that P_b has a Δ_ε -flag with $\Delta_\varepsilon(b)$ at the top and other sections of the form $\Delta_\varepsilon(a)$ for $a \in \mathbf{B}$ with $\rho(a) \geq \rho(b)$. To prove this, suppose that $b \in \mathbf{B}_\lambda$ and set $P := P_b$ for short. Note P has basis $\{yx \mid (y, x) \in \bigcup_{a \in \mathbf{B}_{\geq \lambda}} Y(a) \times X(a, b)\}$. Let $\{\mu_1, \dots, \mu_n\}$ be the finite set

$$\{\mu \in [\lambda, \infty) \mid \bigcup_{a \in \mathbf{B}_\mu} X(a, b) \neq \emptyset\}$$

ordered so that $\mu_r \leq \mu_s \Rightarrow r \leq s$; in particular, $\mu_1 = \lambda$. Let P_r be the subspace of P spanned by $\{yx \mid (y, x) \in \bigcup_{s=r+1}^n \bigcup_{a \in \mathbf{B}_{\mu_s}} Y(a) \times X(a, b)\}$. This defines a filtration $P = P_0 > P_1 > \dots > P_n = 0$ in which the section P_{r-1}/P_r has basis $\{yx + P_r \mid (y, x) \in \bigcup_{a \in \mathbf{B}_{\mu_r}} Y(a) \times X(a, b)\}$. Now we show that each P_{r-1}/P_r has a Δ_ε -flag with sections of the form $\Delta_\varepsilon(a)$ for $a \in \mathbf{B}_{\mu_r}$. There are two cases:

Case 1: $\varepsilon(\mu_r) = +$. In this case, there is an A -module isomorphism

$$\theta : \bigoplus_{a \in \mathbf{B}_{\mu_r}} \bigoplus_{x \in X(a, b)} \Delta(a) \xrightarrow{\sim} P_{r-1}/P_r$$

sending the basis vector $y\bar{e}_a$ ($y \in Y(a)$) in the x th copy of $\Delta(a)$ to $yx + P_r \in P_{r-1}/P_r$. This follows from properties of the basis and is similar to the proof of (5.3).

Case 2: $\varepsilon(\mu_r) = -$. Note that P_{r-1}/P_r is naturally an $A_{\leq \mu_r}$ -module. Let $Q_r := \bar{e}_{\mu_r}(P_{r-1}/P_r)$. This is an A_{μ_r} -module with basis $\{x + P_r \mid x \in X(a, b), a \in \mathbf{B}_{\mu_r}\}$. We claim that the natural multiplication map

$$A_{\leq \mu_r} \bar{e}_{\mu_r} \otimes_{A_{\mu_r}} Q_r \rightarrow P_{r-1}/P_r, \quad y\bar{e}_{\mu_r} \otimes (x + P_r) \mapsto yx + P_r$$

is an isomorphism. This follows because the module on the left hand side is spanned by the vectors $\{y\bar{e}_{\mu_r} \otimes (x + P_r) \mid (y, x) \in \bigcup_{a \in \mathbf{B}_{\mu_r}} Y(a) \times X(a, b)\}$, and the images of these vectors under multiplication are a basis for the module on the right. Hence, $P_{r-1}/P_r \cong j_!^{\mu_r} Q_r$. We deduce that it has a Δ_ε -flag with sections of the form $\bar{\Delta}(a)$ ($a \in \mathbf{B}_{\mu_r}$) on applying the standardization functor to a composition series for Q_r , using the exactness from Lemma 5.16.

We can now complete the proof of the claim. The only thing left is to check that the top section of the Δ_ε -flag we have constructed so far is isomorphic to $\Delta_\varepsilon(b)$. This follows from the constructions just explained: in the case $\varepsilon(\lambda) = +$ we showed that $P_0/P_1 \cong \Delta(b) = \Delta_\varepsilon(b)$, while if $\varepsilon(\lambda) = -$ then the top section is $j_!^\lambda L_\lambda(b) = \Delta_\varepsilon(b)$.

Using the claim just established, we can now classify the irreducible A -modules. For $b \in \mathbf{B}_\lambda$, the proper standard module $\Delta_\varepsilon(b)$ has irreducible head denoted $L(b)$. This follows by the usual properties of adjunctions and the quotient functor $j^\lambda : A_{\leq \lambda}\text{-mod}_{\text{fd}} \rightarrow A_\lambda\text{-mod}_{\text{fd}}, V \mapsto \bar{e}_\lambda V$. Moreover, $L(b)$ is the unique (up to isomorphism) irreducible $A_{\leq \lambda}$ -module such that $j^\lambda L(b) \cong L_\lambda(b)$, hence, the modules $\{L(b) \mid b \in \mathbf{B}\}$ are pairwise inequivalent. To see that they give a full set of irreducible A -modules, let L be any irreducible A -module. By the analog of Remark 5.2, there exists $b \in \mathbf{B}$ such that $e_b L \neq 0$. Then L is a quotient of $P_b = Ae_b$. Finally, using the claim, we deduce that L is a quotient of $\Delta_\varepsilon(a)$ for some $a \in \mathbf{B}$ with $\rho(a) \geq \rho(b)$ and thus L is isomorphic to $L(a)$.

Having classified the irreducible A -modules $\{L(b) \mid b \in \mathbf{B}\}$, $(\mathbf{B}, L, \rho, \Lambda, \leq)$ defines a stratification of \mathcal{R} . We are in the recollement situation of (3.4), with \mathcal{R}_λ identified with $A_\lambda\text{-mod}_{\text{fd}}$. Since (5.6)–(5.7) agrees with (1.1), the standard, proper standard, costandard and proper costandard modules are the correct objects. Moreover, the claim established at the start of the proof verifies the property $(\widehat{P}\Delta_\varepsilon)$. \square

The goal in the remainder of the subsection is to prove a converse to Theorem 5.17.

Theorem 5.18 (Based ε -stratified algebras from ε -highest weight categories). *Let \mathcal{R} be a finite (resp., lower finite, resp., tilting-bounded essentially finite) ε -stratified category*

with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Suppose we are given $\mathbf{B} \subseteq I$ disjoint from Λ and an ε -tilting generator $T = \bigoplus_{i \in I} T_i$ such that each T_b ($b \in \mathbf{B}$) is a direct sum of $T_\varepsilon(b)$ and other $T_\varepsilon(c)$ for c with $\rho(c) < \rho(b)$. Let

$$A := \left(\bigoplus_{i, j \in I} \text{Hom}_{\mathcal{R}}(T_i, T_j) \right)^{\text{op}}$$

For $i, j \in I$ and $b \in \mathbf{B}$, pick morphisms

$$Y(i, b) \subset \text{Hom}_{\mathcal{R}}(T_i, T_b), \quad X(b, j) \subset \text{Hom}_{\mathcal{R}}(T_b, T_j)$$

lifting bases for $\text{Hom}_{\mathcal{R}}(T_i, \nabla_\varepsilon(b))$ and $\text{Hom}_{\mathcal{R}}(\Delta_\varepsilon(b), T_j)$ as in Theorem 4.43 such that $Y(b, b) = \{\text{id}_{T_b}\}$ when $\varepsilon(b) = +$ and $X(b, b) = \{\text{id}_{T_b}\}$ when $\varepsilon(b) = -$. These choices make A into a finite (resp., upper finite, resp., essentially finite) based $(-\varepsilon)$ -stratified algebra with respect to the poset (Λ, \geq) (the opposite ordering on Λ compared to \mathcal{R}).

Proof. We need to check the axioms (εS6) – (εS9) . Theorem 4.43 checks the first one. The axioms (εS7) – (εS8) also hold. For example, if $\varepsilon(\lambda) = +$ and $b \in \mathbf{B}_\lambda$, we have that $Y(b, b) = \{e_b\}$ by the choice of lifts, and $\text{Hom}_{\mathcal{R}}(T_b, \nabla_\varepsilon(a))$ is zero unless $a = b$ or $\rho(a) < \rho(b)$ (remembering we are checking these axioms for $-\varepsilon$ not ε). It remains to check the final axiom (εS9) . The algebra A_λ in the statement of the axiom (remembering that we are working now with the opposite ordering) is the same as the algebra A_λ in Lemma 4.41. By that lemma, there is an algebra isomorphism

$$\phi_\lambda : A_\lambda \xrightarrow{\sim} \text{End}_{\mathcal{R}_\lambda}(j^\lambda T_\lambda)^{\text{op}}, \quad (5.8)$$

where $T_\lambda := \bigoplus_{b \in \mathbf{B}_\lambda} T_b$. If $\varepsilon(\lambda) = +$ then $j^\lambda T_\lambda$ is a minimal projective generator for \mathcal{R}_λ thanks to Theorem 4.2(3), so the algebra on the right hand side of (5.8) is basic and $\bar{e}_\lambda = \sum_{b \in \mathbf{B}_\lambda} \bar{e}_b$ is a decomposition of its identity element as a sum of mutually orthogonal primitive idempotents. If $\varepsilon(\lambda) = -$, we have instead that $j^\lambda T_\lambda$ is a minimal injective cogenerator for \mathcal{R}_λ and the conclusion follows similarly. \square

Corollary 5.19. *Let \mathcal{R} be a finite (resp., upper finite, resp., tilting-bounded essentially finite) ε -stratified category with the usual stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Let $A = \bigoplus_{i, j \in I} e_i A e_j$ be an algebra realization of \mathcal{R} . There is an idempotent expansion $A = \bigoplus_{i, j \in \hat{I}} \hat{e}_i A \hat{e}_j$ with $\mathbf{B} \subseteq \hat{I}$, and finite sets $Y(i, b) \subset \hat{e}_i A \hat{e}_b$, $X(b, j) \subset \hat{e}_b A \hat{e}_j$ for all $i, j \in \hat{I}$ and $b \in \mathbf{B}$, making A into a finite (resp., upper finite, resp., essentially finite) based ε -stratified algebra with ρ as its stratification function.*

Proof. This follows from Theorem 5.18 in the same way as Corollary 5.11 was deduced from Theorem 5.10. \square

5.3. Based stratified and properly stratified algebras. In this subsection, we consider modified versions of Definitions 5.12 and 5.13 which involve bases which do not depend on the sign function ε . These definitions, which were inspired in part by [ELau, Def. 2.17], are relevant when studying fully stratified rather than merely ε -stratified categories.

Definition 5.20. A finite (resp., upper finite, resp., essentially finite) based stratified algebra is a finite-dimensional (resp., locally finite-dimensional, resp., essentially finite-dimensional) locally unital algebra $A = \bigoplus_{i, j \in I} e_i A e_j$ with the following additional data:

- (S1) A subset $\mathbf{B} \subseteq I$ indexing special idempotents $\{e_b \mid b \in \mathbf{B}\}$.
- (S2) A poset (Λ, \leq) which is upper finite in the upper finite case and interval finite in the essentially finite case, such that $\Lambda \cap I = \emptyset$.
- (S3) A function $\rho : \mathbf{B} \rightarrow \Lambda$ with finite fibers $\mathbf{B}_\lambda := \rho^{-1}(\lambda)$.
- (S4) Sets $Y(i, a) \subset e_i A e_a$, $H(a, b) \subset e_a A e_b$, $X(b, j) \subset e_b A e_j$ for $i, j \in I$ and $a, b \in \mathbf{B}$.

Let $Y(a) := \bigcup_{i \in I} Y(i, a)$ and $X(b) := \bigcup_{j \in I} X(b, j)$. The axioms are as follows:

- (S5) The products yhx for $(y, h, x) \in \bigcup_{a, b \in \mathbf{B}} Y(a) \times H(a, b) \times X(b)$ are a basis for A .

- (S6) For $a, b \in \mathbf{B}$ with $a \neq b$, the set $H(a, b)$ is empty unless $\rho(a) = \rho(b)$, the sets $Y(b, a)$ and $X(a, b)$ are empty unless $\rho(b) < \rho(a)$, and $Y(a, a) = X(a, a) = \{e_a\}$.
- (S7) The finite-dimensional algebra A_λ defined as in Definition 5.12 is basic and $\bar{e}_\lambda = \sum_{b \in \mathbf{B}_\lambda} \bar{e}_b$ is a decomposition of its identity element as a sum of mutually orthogonal primitive idempotents.

We say that A is *symmetrically based* if there is also some given algebra anti-involution $\sigma : A \rightarrow A$ with $\sigma(e_i) = e_i$ and $Y(i, b) = \sigma(X(b, i))$ for all $i \in I, b \in \mathbf{B}$, such that each of the algebras A_λ ($\lambda \in \Lambda$) is σ_λ -symmetric in the sense of Definition 4.51, where σ_λ here is the anti-involution of A_λ induced by σ .

Here is the same definition rewritten in the special case that the stratification function ρ is a bijection.

Definition 5.21. A *finite* (resp., *upper finite*, resp., *essentially finite*) *based properly stratified algebra* is a finite-dimensional (resp., locally finite-dimensional, resp., essentially finite-dimensional) locally unital algebra $A = \bigoplus_{i, j \in I} e_i A e_j$ with the following additional data:

- (PS1) A subset $\Lambda \subseteq I$ indexing *special idempotents* $\{e_\lambda \mid \lambda \in \Lambda\}$.
- (PS2) A poset (Λ, \leq) upper finite in the upper finite case and interval finite in the essentially finite case.
- (PS3) Sets $Y(i, \lambda) \subset e_i A e_\lambda$, $H(\lambda) \subset e_\lambda A e_\lambda$, $X(\lambda, i) \subset e_\lambda A e_i$ for $\lambda \in \Lambda, i \in I$.

Let $Y(\lambda) := \bigcup_{i \in I} Y(i, \lambda)$ and $X(\lambda) := \bigcup_{i \in I} X(\lambda, i)$. The axioms are as follows.

- (PS4) The products yxh for $(y, h, x) \in \bigcup_{\lambda \in \Lambda} Y(\lambda) \times H(\lambda) \times X(\lambda)$ are a basis for A .
- (PS5) For $\lambda, \mu \in \Lambda$, the sets $Y(\mu, \lambda)$ and $X(\lambda, \mu)$ are empty unless $\mu \leq \lambda$, and $Y(\lambda, \lambda) = X(\lambda, \lambda) = \{e_\lambda\}$.
- (PS6) The finite-dimensional algebra A_λ defined as in Definition 5.13 is basic and local.

We say that A is *symmetrically based* if there is also some given algebra anti-involution $\sigma : A \rightarrow A$ with $\sigma(e_i) = e_i$ and $Y(i, \lambda) = \sigma(X(\lambda, i))$ for all $i \in I, \lambda \in \Lambda$, such that each of the algebras A_λ ($\lambda \in \Lambda$) is σ_λ -symmetric, where σ_λ here is the anti-involution of A_λ induced by σ .

In the remainder of the subsection, we just explain the results for based stratified algebras, since based properly stratified algebras are a special case. For the next lemma, we adopt the shorthands

$$YH(i, b) := \{yh \mid (y, h) \in \bigcup_{a \in \mathbf{B}} Y(i, a) \times H(a, b)\}, \quad (5.9)$$

$$HX(b, j) := \{hx \mid (h, x) \in \bigcup_{a \in \mathbf{B}} H(b, a) \times X(a, j)\}. \quad (5.10)$$

Also set $YH(b) := \bigcup_{i \in I} YH(i, b)$ and $HX(b) := \bigcup_{j \in I} HX(b, j)$.

Lemma 5.22. *Suppose that A is a based stratified algebra as in Definition 5.20. Also let $\varepsilon : \Lambda \rightarrow \{\pm\}$ be any choice of sign function. Then A is a based ε -stratified algebra in the sense of Definition 5.12 with the required sets $Y(i, b)$ and $X(b, j)$ for that being the sets $YH(i, b)$ and $X(b, j)$ in the present setup if $\varepsilon(\rho(b)) = +$, or the sets $Y(i, b)$ and $HX(b, j)$ in the present setup if $\varepsilon(\rho(b)) = -$.*

Proof. This follows on comparing Definitions 5.12 and 5.20. \square

This means that the results from the previous subsection apply to based stratified algebras too. In particular, we define the standard, proper standard, costandard and proper costandard modules as in (5.6)–(5.7). The modules $\Delta(b)$ and $\bar{\Delta}(b)$ have standard bases $\{y\bar{e}_b \mid y \in YH(b)\}$ and $\{y\tilde{e}_b \mid y \in Y(b)\}$, respectively. Similarly, one can introduce costandard bases for $\nabla(b)$ and $\bar{\nabla}(b)$ indexed by the sets $HX(b)$ and $X(b)$, respectively. Note also that the basic algebra

$$A_\lambda = \bigoplus_{a, b \in \mathbf{B}_\lambda} \bar{e}_a A_\lambda \bar{e}_b$$

has basis $\{\bar{h} \mid h \in \bigcup_{a,b \in \mathbf{B}_\lambda} H(a,b)\}$.

Theorem 5.23 (Fully stratified categories from based stratified algebras). *Let A be a finite (resp., upper finite, resp., essentially finite) based stratified algebra as above. The modules*

$$\{L(b) := \text{hd } \Delta(b) \cong \text{hd } \bar{\Delta}(b) \cong \text{soc } \bar{\nabla}(b) \cong \text{soc } \nabla(b) \mid b \in \mathbf{B}\}$$

give a full set of pairwise inequivalent irreducible left A -modules. Moreover, $\mathcal{R} := A\text{-mod}_{\text{fd}}$ (resp., $\mathcal{R} := A\text{-mod}_{\text{fd}}$, resp., $\mathcal{R} := A\text{-mod}_{\text{fd}}$) is a finite (resp., upper finite, resp., essentially finite) fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ with strata $\mathcal{R}_\lambda := A_\lambda\text{-mod}_{\text{fd}}$. If A is symmetrically based with anti-involution σ then $?^\circ : \mathcal{R} \rightarrow \mathcal{R}$ is a Chevalley duality of \mathcal{R} in the sense of Definition 4.53.

Proof. Using Lemma 5.22, the first part follows from Theorem 5.17 applied twice, once with $\varepsilon = +$ and once with $\varepsilon = -$. For the final part about Chevalley duality, axiom (Ch1) from Definition 4.53 is established in the course of the proof of Theorem 5.17, and (Ch2) follows from the definition of symmetrically based stratified algebra. \square

For the converse recall the definition of tilting-rigid from Definition 4.36.

Theorem 5.24 (Based stratified algebras from fully stratified categories). *Let \mathcal{R} be a finite (resp., lower finite, resp., essentially finite) fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Assume that \mathcal{R} is tilting-rigid with weakly symmetric strata. Suppose we are given $\mathbf{B} \subseteq I$ disjoint from Λ and a tilting generator $T = \bigoplus_{i \in I} T_i$ such that each T_b ($b \in \mathbf{B}$) is a direct sum of $T(b)$ and other $T(c)$ for c with $\rho(c) < \rho(b)$. Let*

$$A := \left(\bigoplus_{i,j \in I} \text{Hom}_{\mathcal{R}}(T_i, T_j) \right)^{\text{op}}$$

(1) *For $i, j \in I$ and $a, b \in \mathbf{B}$, pick morphisms*

$$Y(i, a) \subset \text{Hom}_{\mathcal{R}}(T_i, T_a), \quad H(a, b) \subset \text{Hom}_{\mathcal{R}}(T_a, T_b), \quad X(b, j) \subset \text{Hom}_{\mathcal{R}}(T_b, T_j)$$

lifting bases for $\text{Hom}_{\mathcal{R}}(T_i, \bar{\nabla}(a))$, $\text{Hom}_{\mathcal{R}}(\Delta(a), \nabla(b))$ and $\text{Hom}_{\mathcal{R}}(\bar{\Delta}(b), T_j)$ as in Theorem 4.45 such that $Y(b, b) = X(b, b) = \{\text{id}_{T_b}\}$. These choices give a triangular basis making into a finite (resp., upper finite, resp., essentially finite) based stratified algebra with respect to the poset (Λ, \geq) (the opposite ordering on Λ compared to \mathcal{R}).

(2) *If in addition \mathcal{R} has a Chevalley duality $?^\vee$ and, in a suitable realization, the modules corresponding to each T_i possess non-degenerate symmetric bilinear forms satisfying the adjunction property as in (4.47), then the triangular basis in (1) can be chosen so that A is symmetrically based.*

Proof. Part (1) is similar to the proof of Theorem 5.18, using Theorem 4.45 in place of Theorem 4.43. Part (2) follows in the same way as in the proof of Theorem 5.11(2). \square

Corollary 5.25. *Let \mathcal{R} be a finite (resp., upper finite, resp., essentially finite) fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Let $A = \bigoplus_{i,j \in I} e_i A e_j$ be an algebra realization of \mathcal{R} .*

(1) *Assume that \mathcal{R} is tilting-rigid with weakly symmetric strata. Then there is an idempotent expansion $A = \bigoplus_{i,j \in \hat{I}} \hat{e}_i A \hat{e}_j$ with $\mathbf{B} \subseteq \hat{I}$, and finite sets*

$$Y(i, a) \subset \hat{e}_i A \hat{e}_a, \quad H(a, b) \subset \hat{e}_a A \hat{e}_b, \quad X(b, j) \subset \hat{e}_b A \hat{e}_j$$

for all $i, j \in \hat{I}$ and $a, b \in \mathbf{B}$, making A into an upper finite (resp., essentially finite) based stratified algebra.

(2) *Assume that \mathcal{R} is tilting-rigid with a Chevalley duality $?^\vee$ and that $\text{char } \mathbb{k} \neq 2$. Then the choices in (1) can be made so that A is symmetrically based with anti-involution σ realizing $?^\vee$.*

Proof. This follows from Theorem 5.24 in the same way as Corollary 5.11 was deduced from Theorem 5.10. One also needs to use the fact that the Ringel dual \mathcal{R}' of \mathcal{R} is tilting-rigid by Theorem 4.42. \square

5.4. Algebras with a triangular basis. The final axiom (S7) of Definition 5.20, namely, that the algebra A_λ is basic, is quite restrictive. However, this assumption is not essential, as we will explain in this subsection. The following simply repeats Definition 5.20 with the final axiom dropped, but at the same time we switch to using the notation $\partial : \mathbf{S} \rightarrow \Lambda$ where we had $\rho : \mathbf{B} \rightarrow \Lambda$ before.

Definition 5.26. Let $A = \bigoplus_{i,j \in I} e_i A e_j$ be a finite-dimensional (resp., locally finite-dimensional, resp., essentially finite-dimensional) locally unital algebra. We say that A has a *triangular basis* if we are given the following additional data:

- (TB1) A subset $\mathbf{S} \subseteq I$ indexing *special idempotents* $\{e_s \mid s \in \mathbf{S}\}$.
- (TB2) A poset (Λ, \leq) which is upper finite in the locally finite-dimensional case and interval finite in the essentially finite-dimensional case, such that $\Lambda \cap I = \emptyset$.
- (TB3) A function $\partial : \mathbf{S} \rightarrow \Lambda$ with finite fibers $\mathbf{S}_\lambda := \partial^{-1}(\lambda)$.
- (TB4) Sets $Y(i, s) \subset e_i A e_s$, $H(s, t) \subset e_s A e_t$, $X(t, j) \subset e_t A e_j$ for $i, j \in I$ and $s, t \in \mathbf{S}$.

Let $Y(s) := \bigcup_{i \in I} Y(i, s)$ and $X(t) := \bigcup_{j \in I} X(t, j)$. The axioms are as follows:

- (TB5) The products yhx for $(y, h, x) \in \bigcup_{s,t \in \mathbf{S}} Y(s) \times H(s, t) \times X(t)$ are a basis for A .
- (TB6) For $s, t \in \mathbf{S}$ with $s \neq t$, the set $H(s, t)$ is empty unless $\partial(s) = \partial(t)$, the sets $Y(t, s)$ and $X(s, t)$ are empty unless $\partial(t) < \partial(s)$, and $Y(s, s) = X(s, s) = \{e_s\}$.

Suppose that A has a triangular basis as in Definition 5.26. We define algebras $A_\lambda = \bar{e}_\lambda A_{\leq \lambda} \bar{e}_\lambda$ for each $\lambda \in \Lambda$ like at the end of Definition 5.12. Thus, we let $e_\lambda := \sum_{s \in \mathbf{S}_\lambda} e_s$, then set $A_\lambda := \bar{e}_\lambda A_{\leq \lambda} \bar{e}_\lambda$ where $A_{\leq \lambda}$ is the quotient of A by the two-sided ideal generated by $\{e_\mu \mid \mu \not\leq \lambda\}$. Corollary 5.15 carries over to show that $A_{\leq \lambda}$ has basis $\bar{y}\bar{h}\bar{x}$ for all $y \in Y(s), h \in H(s, t), x \in X(t)$ and $s, t \in \mathbf{S}$ with $\partial(s), \partial(t) \leq \lambda$. Hence, A_λ has basis $\{\bar{h} \mid h \in \bigcup_{s,t \in \mathbf{S}_\lambda} H(s, t)\}$. Let $j^\lambda : A_{\leq \lambda}\text{-mod} \rightarrow A_\lambda\text{-mod}, V \mapsto \bar{e}_\lambda V$ be the quotient functor and define $j_!^\lambda$ and j_*^λ analogously to (5.4).

Lemma 5.27. *The functors $j_!^\lambda$ and j_*^λ are exact.*

Proof. By the argument from the proof of Lemma 5.16, there is an isomorphism of right A_λ -modules $\bigoplus_{s \in \mathbf{S}_\lambda} \bigoplus_{y \in Y(s)} \bar{e}_s A_\lambda \xrightarrow{\sim} A_{\leq \lambda} \bar{e}_\lambda$ sending the vector \bar{e}_s in the y th copy of $\bar{e}_s A_\lambda$ to $\bar{y} \in A_{\leq \lambda} \bar{e}_\lambda$. So the right A_λ -module $A_{\leq \lambda} \bar{e}_\lambda$ is projective, which implies the exactness of $j_!^\lambda$. Similarly, the left A_λ -module $\bar{e}_\lambda A_{\leq \lambda}$ is projective, which implies the exactness of j_*^λ . \square

The following theorem is essentially [GRS, Th. 3.5], although we give a self-contained proof since our notation is different enough. See Remark 5.30 for further historical discussion.

Theorem 5.28 (Fully stratified categories from triangular bases). *Let A be a finite-dimensional (resp., locally finite-dimensional, resp. essentially finite-dimensional algebra with a triangular basis as above. Let $\rho : \mathbf{B} \rightarrow \Lambda$ be a function whose fibers $\mathbf{B}_\lambda := \rho^{-1}(\lambda)$ label a full set $\{L_\lambda(b) \mid b \in \mathbf{B}_\lambda\}$ of pairwise inequivalent irreducible left A_λ -modules. Let $\bar{\Delta}(b) := j_!^\lambda L_\lambda(b)$ and $\bar{\nabla}(b) := j_*^\lambda L_\lambda(b)$ for $b \in \mathbf{B}_\lambda$. Then the modules*

$$\{L(b) := \text{hd } \bar{\Delta}(b) \cong \text{soc } \bar{\nabla}(b) \mid b \in \mathbf{B}\}$$

give a full set of pairwise inequivalent irreducible left A -modules. Moreover, $\mathcal{R} := A\text{-mod}_{\text{fd}}$ (resp., $\mathcal{R} := A\text{-mod}_{\text{fd}}$, resp., $\mathcal{R} := A\text{-mod}_{\text{fd}}$) is a finite (resp., upper finite, resp., essentially finite) fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Its strata are the categories $\mathcal{R}_\lambda := A_\lambda\text{-mod}_{\text{fd}}$, with standardization and costandardization functors as in (5.4).

Proof. Take $u \in \mathbf{S}_\lambda$ and any $b \in \mathbf{B}_\lambda$ such that $\bar{e}_u L_\lambda(b) \neq 0$. We claim that Ae_u has a $\bar{\Delta}$ -flag with $\bar{\Delta}(b)$ at the top and other sections of the form $\bar{\Delta}(c)$ for c with $\rho(c) \geq \lambda$. To see this, let $P := Ae_u$ for short. Note P has basis

$$\{yhx \mid (y, h, x) \in \bigcup_{\mu \geq \lambda} \bigcup_{s, t \in \mathbf{S}_\mu} Y(s) \times H(s, t) \times X(t, u)\}.$$

Let $\{\mu_1, \dots, \mu_n\}$ be the finite set $\{\mu \in [\lambda, \infty) \mid \bigcup_{t \in \mathbf{S}_\mu} X(t, u) \neq \emptyset\}$ enumerated in some order refining \leq . There is a filtration $P = P_0 > P_1 > \dots > P_n = 0$ in which the section P_{r-1}/P_r has basis $\{yhx + P_r \mid (y, h, x) \in \bigcup_{s, t \in \mathbf{S}_{\mu_r}} Y(s) \times H(s, t) \times X(t, u)\}$. Moreover, $P_{r-1}/P_r \cong j_1^{\mu_r} Q_r$ where $Q_r := \bar{e}_{\mu_r}(P_{r-1}/P_r)$. This follows by a similar argument to the Case 2 in the proof of Theorem 5.17. Since $j_1^{\mu_r}$ is exact by Lemma 5.27, it follows that P_{r-1}/P_r has a $\bar{\Delta}$ -flag with sections $\bar{\Delta}(c)$ for $c \in \mathbf{B}_{\mu_r}$. So we have proved that P has a $\bar{\Delta}$ -flag with sections $\bar{\Delta}(c)$ for $c \in \mathbf{B}$ with $\rho(c) \geq \lambda$. Moreover, $P_0/P_1 \cong j_1^\lambda(A_\lambda \bar{e}_u)$. Since $A_\lambda \bar{e}_u$ has $L_\lambda(b)$ in its head, it follows that the $\bar{\Delta}$ -flag can be chosen so that it has $\bar{\Delta}(b)$ at its top.

Now we can classify the irreducible left A -modules. As in the penultimate paragraph of the proof of Theorem 5.17, the modules $\{L(b) := \text{hd } \bar{\Delta}(b) \mid b \in \mathbf{B}\}$ are pairwise inequivalent irreducible A -modules. It remains to show that any irreducible left A -module L is isomorphic to some such module. There exists $u \in \mathbf{S}$ such that $e_u L \neq 0$. Hence, L is a quotient of Ae_u . By considering the filtration of Ae_u from the previous paragraph we deduce that L is a quotient of $\bar{\Delta}(c)$ for some $c \in \mathbf{B}$, i.e., $L \cong L(c)$.

At this point, we have in hand the data of a stratification of \mathcal{R} with strata $\mathcal{R}_\lambda := A_\lambda\text{-mod}_{\text{id}}$ and standardization and costandardization functors as in (5.4). For each $b \in \mathbf{B}_\lambda$, choose $u \in \mathbf{S}_\lambda$ such that $\bar{e}_u L_\lambda(b) \neq 0$ then set $P_b := Ae_u$. The claim established in the first paragraph of the proof checks that these modules satisfy the property $(\widehat{P\Delta}_-)$, hence, \mathcal{R} is an upper finite (resp., essentially finite) $--$ -stratified category. Finally we deduce that it is fully stratified using the criterion from Lemma 3.20(iv) plus Lemma 5.27. \square

Corollary 5.29. *Let A be as above. If each of the finite-dimensional algebras A_λ is quasi-hereditary (e.g., they could all be semisimple), then the stratification can be refined to make the category \mathcal{R} from Theorem 5.28 into a highest weight category.*

Proof. Combine Theorem 5.28 and Corollary 3.67. \square

Remark 5.30. We did not fully appreciate the utility of Definition 5.26 before seeing [GRS], in which Gao, Rui and Song introduce a notion of an algebra with a *weak triangular decomposition* and give a (slightly different) proof of Theorem 5.28 for such algebras. They justify their definition by constructing several interesting families of examples, namely, cyclotomic quotients of the affine oriented Brauer and HOMFLY-PT skein categories and of the affine Brauer and Kauffman skein categories. In the special case that $I = \mathbf{S}$, i.e., all distinguished idempotents are special, our notion of an algebra with a triangular basis is exactly equivalent to the notion of an algebra with a weak triangular decomposition. More precisely, a weak triangular decomposition is the data of subspaces $A^- = \bigoplus_{i, j \in I} e_i A^- e_j$, $A^\circ = \bigoplus_{i, j \in I} e_i A^\circ e_j$, $A^+ = \bigoplus_{i, j \in I} e_i A^+ e_j$ for $i, j \in I$ subject to certain axioms. Picking homogeneous bases $Y(i, j)$, $H(i, j)$ and $X(i, j)$ for $e_i A^- e_j$, $e_i A^\circ e_j$ and $e_i A^+ e_j$, respectively, produces a triangular basis in the sense of Definition 5.26. Conversely given a triangular basis one obtains a weak triangular decomposition by replacing the bases by the subspaces that they span.

5.5. Algebras with a triangular decomposition. Let A be an algebra with a triangular basis as in Definition 5.26 and assume in addition that $I = \mathbf{S}$, i.e., all of the distinguished idempotents are special. Let A^\flat and A^\sharp be the subspaces spanned by $\{yh \mid (y, h) \in \bigcup_{i, j \in I} Y(i) \times H(i, j)\}$ and $\{hx \mid (h, x) \in \bigcup_{i, j \in I} H(i, j) \times X(j)\}$, respectively.

If it happens that these subspaces are locally unital subalgebras¹¹ of A then A has a triangular decomposition in the following sense.

Definition 5.31. Let $A = \bigoplus_{i,j \in I} e_i A e_j$ be a finite-dimensional (resp., locally finite-dimensional, resp., essentially finite-dimensional) locally unital algebra. A *triangular decomposition* of A is the following additional data:

- (TD1) A poset (Λ, \leq) which is upper finite in the locally finite-dimensional case or interval finite in the essentially finite-dimensional case.
- (TD2) A function $\partial : I \rightarrow \Lambda$ with finite fibers $I_\lambda := \partial^{-1}(\lambda)$.
- (TD3) Locally unital subalgebras A^\flat and A^\sharp .

We call A^\flat and A^\sharp the *negative* and *positive Borel subalgebras*. Let $A^\circ := A^\flat \cap A^\sharp$. This is also a locally unital subalgebra called the *Cartan subalgebra*. The following axioms are required to hold:

- (TD4) A^\flat is a projective right A° -module and A^\sharp is a projective left A° -module.
- (TD5) The natural multiplication map $A^\flat \otimes_{A^\circ} A^\sharp \rightarrow A$ is a linear isomorphism.
- (TD6) For $i, j \in I$, $e_j A^\flat e_i$ and $e_i A^\sharp e_j$ are zero unless $\partial(j) \leq \partial(i)$, and $e_i A^\flat e_j = e_i A^\sharp e_j$ when $\partial(i) = \partial(j)$.

Remark 5.32. Our formulation of Definition 5.31 has been influenced by the definition of a *triangular category* from a recent preprint of Sam and Snowden [SS]; these are finite-dimensional categories satisfying equivalent axioms to algebras with an upper finite triangular decomposition in the above sense in which the Cartan subalgebra is semisimple. In an earlier draft, we had formulated a slightly more restrictive notion which we now refer to a *split triangular decomposition*, as follows. Let $A = \bigoplus_{i,j \in I} e_i A e_j$ be a finite-dimensional (resp., locally finite-dimensional, resp., essentially finite-dimensional) locally unital algebra. We say that A has a *split triangular decomposition* if we have the additional data:

- (STD1) A poset (Λ, \leq) which is upper finite in the locally finite-dimensional case and interval finite in the essentially finite-dimensional case.
- (STD2) A function $\partial : I \rightarrow \Lambda$ with finite fibers $I_\lambda := \partial^{-1}(\lambda)$.
- (STD3) Locally unital subalgebras A^- , A° and A^+ .

Letting $\mathbb{K} := \bigoplus_{i \in I} \mathbb{k} e_i$, the axioms are:

- (STD4) The subspaces $A^\flat := A^- A^\circ$ and $A^\sharp := A^\circ A^+$ are subalgebras.
- (STD5) The natural multiplication map $A^- \otimes_{\mathbb{K}} A^\circ \otimes_{\mathbb{K}} A^+ \rightarrow A$ is a linear isomorphism.
- (STD6) For $i, j \in I$ with $i \neq j$, $e_i A^\circ e_j$ is zero unless $\partial(i) = \partial(j)$, $e_j A^- e_i$ and $e_i A^+ e_j$ are zero unless $\partial(j) < \partial(i)$, and $e_i A^\flat e_i = e_i A^\sharp e_i = \mathbb{k} e_i$ for all $i \in I$.

The axiom (STD5) implies that $A^\flat \cong A^- \otimes_{\mathbb{K}} A^\circ$ and $A^\sharp \cong A^\circ \otimes_{\mathbb{K}} A^+$. Hence, by associativity of tensor product we have that

$$A^\flat \otimes_{A^\circ} A^\sharp \cong A^- \otimes_{\mathbb{K}} A^\circ \otimes_{A^\circ} A^\circ \otimes_{\mathbb{K}} A^+ \cong A^- \otimes_{\mathbb{K}} A^\circ \otimes_{\mathbb{K}} A^+ \cong A,$$

proving (TD5). Moreover, the isomorphisms $A^\flat \cong A^- \otimes_{\mathbb{K}} A^\circ$ and $A^\sharp \cong A^\circ \otimes_{\mathbb{K}} A^+$ show that A^\flat and A^\sharp are I -free in the sense of Definition 2.17 as right and left A° -modules, respectively, which implies (TD4). Axiom (TD6) is also easily deduced from (STD6). When they hold, the axioms (STD4)–(STD6) are easier to check than (TD4)–(TD6), so this gives a practical way to obtain triangular decompositions. In fact, most of the examples of triangular decompositions arising from diagrammatic monoidal categories considered in [SS] and elsewhere are split triangular decompositions, so the split formulation is useful.

Remark 5.33. In [HN], Holmes and Nakano introduced a notion of a \mathbb{Z} -graded algebra with a triangular decomposition. To explain the connection to our setup, suppose we

¹¹Locally unital subalgebra means subspace closed under multiplication and containing all of the distinguished idempotents.

are given a unital \mathbb{Z} -graded algebra $\tilde{A} = \bigoplus_{\lambda \in \mathbb{Z}} \tilde{A}_\lambda$. There is an associated locally unital algebra $A = \bigoplus_{\lambda, \mu \in \mathbb{Z}} e_\lambda A e_\mu$ with $e_\lambda A e_\mu := \tilde{A}_{\lambda-\mu}$ and multiplication induced by multiplication in \tilde{A} in the natural way. Moreover, any \mathbb{Z} -graded left \tilde{A} -module $V = \bigoplus_{\lambda \in \mathbb{Z}} V_\lambda$ can be viewed as a left A -module with $e_\lambda V := V_\lambda$; this defines an isomorphism from the usual category $\tilde{A}\text{-grmod}$ of \mathbb{Z} -graded \tilde{A} -modules and degree-preserving morphisms to the category $A\text{-mod}$ of locally unital A -modules. If we start with \tilde{A} that is a finite-dimensional \mathbb{Z} -graded algebra with a triangular decomposition $(\tilde{A}^-, \tilde{A}^\circ, \tilde{A}^+)$ as in [HN] (see also [BT, Def. 3.1]) then the essentially finite-dimensional locally unital algebra A and the subalgebras A°, A^- and A^+ obtained via this construction has a split triangular decomposition, with $I = \Lambda = \mathbb{Z}$ ordered in the natural way.

To make the connection with Definition 5.26, suppose that A has a triangular decomposition. For $\lambda \in \Lambda$, let $1_\lambda := \sum_{i \in I_\lambda} e_i$. The axioms imply that $e_i A^\circ e_j = 0$ unless $\partial(i) = \partial(j)$, so $1_\lambda A^\circ 1_\mu = 0$ for $\lambda \neq \mu$. It follows that $\{1_\lambda \mid \lambda \in \Lambda\}$ are mutually orthogonal central idempotents in A° , and the Cartan subalgebra has the ‘‘block’’ decomposition

$$A^\circ = \bigoplus_{\lambda \in \Lambda} A_\lambda^\circ \quad \text{where} \quad A_\lambda^\circ := 1_\lambda A^\circ = A^\circ 1_\lambda. \quad (5.11)$$

Lemma 5.34. *Let A be as in Definition 5.31 with $\Lambda \cap I = \emptyset$. Suppose we are given $\mathbf{S} \subseteq I$ such that all $e_i A^\flat$ and $A^\sharp e_j$ are \mathbf{S} -free as right and left A° -modules, respectively. For $i, j \in I, s, t \in \mathbf{S}$, one can choose subsets $Y(i, s) \subset e_i A^\flat e_s, X(t, j) \subset e_t A^\sharp e_j$ so that*

- (i) $e_i A^\flat = \bigoplus_{s \in \mathbf{S}} \bigoplus_{y \in Y(i, s)} y A^\circ$ with $y A^\circ \cong e_s A^\circ$ for $y \in Y(i, s)$;
- (ii) $A^\sharp e_j = \bigoplus_{t \in \mathbf{S}} \bigoplus_{x \in X(t, j)} A^\circ x$ with $A^\circ x \cong A^\circ e_t$ for $x \in X(t, j)$;
- (iii) $Y(t, t) = X(t, t) = \{e_t\}$ for all $t \in \mathbf{S}$.

Also let $H(s, t)$ be a basis for $e_s A^\circ e_t$. This makes $A = \bigoplus_{i, j \in I} e_i A e_j$ into an algebra with a triangular basis in the sense of Definition 5.26 with $\partial : \mathbf{S} \rightarrow \Lambda$ being the restriction of the given function $\partial : I \rightarrow \Lambda$. For $\lambda \in \Lambda$ and $e_\lambda := \sum_{s \in \mathbf{S}_\lambda} e_s$, the subquotient $A_\lambda = \bar{e}_\lambda A_{\leq \lambda} \bar{e}_\lambda$ defined after Definition 5.26 is isomorphic to the subalgebra $e_\lambda A_\lambda^\circ e_\lambda$ of A_λ° . Moreover, we have that $A_\lambda^\circ = A_\lambda^\circ e_\lambda A_\lambda^\circ$ so A_λ is Morita equivalent to A_λ° .

Proof. By the definition of \mathbf{S} -free, there are subsets $Y(i, s) \subset e_i A^\flat e_s$ as in (i). Since $e_i A^\flat e_s$ is zero unless $\partial(i) \leq \partial(s)$, we have that $Y(i, s) = \emptyset$ unless $\partial(i) \leq \partial(s)$. Suppose that $t \in \mathbf{S}_\lambda := \mathbf{S} \cap I_\lambda$. By (TD6), we have that

$$e_t A^\flat 1_\lambda = e_t A_\lambda^\circ = \bigoplus_{s \in \mathbf{S}_\lambda} \bigoplus_{y \in Y(t, s)} y A_\lambda^\circ,$$

i.e., the sets $Y(t, s)$ for $s \in \mathbf{S}_\lambda$ come from an \mathbf{S} -free decomposition of $e_t A_\lambda^\circ$. This means we can choose them so that $Y(t, t) = \{e_t\}$ as in (iii), in which case $Y(t, s) = \emptyset$ for $s \in \mathbf{S}_\lambda$ with $s \neq t$. Hence, for $s, t \in \mathbf{S}$ with $s \neq t$, we have that $Y(t, s) = \emptyset$ unless $\partial(t) < \partial(s)$. Similarly, we choose subsets $X(t, j) \subset e_t A^\sharp e_j$ according to (ii) and (iii), and then for $s, t \in \mathbf{S}$ with $s \neq t$ we have that $X(s, t) = \emptyset$ unless $\partial(t) < \partial(s)$. Note also that $H(s, t) = \emptyset$ unless $\partial(s) = \partial(t)$ due to (5.11). Thus we have the required data from (TB1)–(TB4), and the conditions of (TB6) are satisfied.

In this paragraph, we check (TB5). Let $Y(s) = \bigcup_{i \in I} Y(i, s)$ and $X(t) = \bigcup_{j \in I} X(t, j)$. We have seen already that $A^\flat = \bigoplus_{s \in \mathbf{S}} \bigoplus_{y \in Y(s)} y A^\circ$ and $A^\sharp = \bigoplus_{t \in \mathbf{S}} \bigoplus_{x \in X(t)} A^\circ x$. Tensoring these together, we deduce that

$$A^\flat \otimes_{A^\circ} A^\sharp = \bigoplus_{s, t \in \mathbf{S}} \bigoplus_{y \in Y(s), x \in X(t)} y A^\circ \otimes_{A^\circ} A^\circ x.$$

Each summand $y A^\circ \otimes_{A^\circ} A^\circ x$ here is isomorphic to $e_s A^\circ \otimes_{A^\circ} A^\circ e_t \cong e_s A^\circ e_t$. We deduce that $A^\flat \otimes_{A^\circ} A^\sharp$ has basis $\{y h \otimes x = y \otimes h x \mid (y, h, x) \in \bigcup_{s, t \in \mathbf{S}} Y(s) \times H(s, t) \times X(t)\}$. Then we use (TD5) to see that the axiom (TB5) is satisfied.

Finally we must identify the algebra A_λ . The quotient map $A \rightarrow A_{\leq \lambda}$ restricts to a homomorphism $\phi : A^\circ \rightarrow A_{\leq \lambda}$ which further restricts to

$$\phi_\lambda : e_\lambda A_\lambda^\circ e_\lambda \xrightarrow{\sim} A_\lambda. \quad (5.12)$$

The subalgebra A_λ° has basis $\{yhx \mid (y, h, x) \in \bigcup_{i, j \in I_\lambda, s, t \in \mathbf{S}_\lambda} Y(i, s) \times H(s, t) \times X(t, j)\}$, hence, $A_\lambda^\circ = A_\lambda^\circ e_\lambda A_\lambda^\circ$. The subalgebra A_λ of $e_\lambda A_\lambda^\circ e_\lambda$ has basis $\bigcup_{s, t \in \mathbf{S}_\lambda} H(s, t)$. It follows that ϕ_λ sends a basis to a basis, so it is an isomorphism. \square

The freeness assumption in Lemma 5.34 may seem restrictive, but one can always pass to an idempotent expansion so that this is the case. In fact, we can do this in such a way that the algebras A_λ are *basic*, thereby giving A the structure of a based stratified algebra rather than merely an algebra with a triangular basis:

Theorem 5.35 (Based stratified algebras from triangular decompositions). *Suppose that A has a triangular decomposition as in Definition 5.31. Let $A^\circ = \bigoplus_{i, j \in \hat{I}} \hat{e}_i A^\circ \hat{e}_j$ be an idempotent expansion of $A^\circ = \bigoplus_{i, j \in I} e_i A^\circ e_j$ such that*

- (i) $\hat{I} \cap \Lambda = \emptyset$;
- (ii) \hat{I} contains a subset \mathbf{B} indexing a full set $\{\hat{e}_b \mid b \in \mathbf{B}\}$ of pairwise non-conjugate primitive idempotents in A° ;
- (iii) there is a function $q : \hat{I} \rightarrow I$ with $|q^{-1}(i)| < \infty$ and $e_i = \sum_{j \in q^{-1}(i)} \hat{e}_j$ for $i \in I$.

Then $A = \bigoplus_{i, j \in \hat{I}} \hat{e}_i A \hat{e}_j$ has a triangular decomposition with the given Borel subalgebras, taking the function from (TD2) now to be $\rho := \partial \circ q : \hat{I} \rightarrow \Lambda$. Moreover, $\hat{e}_i A^{\sharp} \hat{e}_j$ are \mathbf{B} -free as right and left A° -modules, respectively. Hence, we can apply the construction of Lemma 5.34 to $A = \bigoplus_{i, j \in \hat{I}} \hat{e}_i A \hat{e}_j$ to make A into a based stratified algebra in the sense of Definition 5.26 with $\rho : \mathbf{B} \rightarrow \Lambda$ defined by restriction.

Proof. The fact that we have in hand a triangular decomposition of $A = \bigoplus_{i, j \in \hat{I}} \hat{e}_i A \hat{e}_j$ is immediately clear from the nature of Definition 5.31. Since $1_\lambda A^{\sharp} \hat{e}_j$ is a finite-dimensional projective left A_λ° , Lemma 2.18 implies that it is \mathbf{B} -free as a left A_λ -module. Hence $A^{\sharp} \hat{e}_j = \bigoplus_{\lambda \in \Lambda} 1_\lambda A^{\sharp} \hat{e}_j$ is \mathbf{B} -free as a left module. Similarly, we get that $\hat{e}_i A^{\flat}$ is \mathbf{B} -free as a right module. So now Lemma 5.34 can be applied and we obtain a triangular basis such that $A_\lambda \cong \hat{e}_\lambda A_\lambda^\circ \hat{e}_\lambda$ for $\hat{e}_\lambda := \sum_{b \in \mathbf{B}_\lambda} \hat{e}_b$. By the choice of the idempotents $\{\hat{e}_b \mid b \in \mathbf{B}\}$, $\hat{e}_\lambda A_\lambda^\circ \hat{e}_\lambda$ is the basic algebra that is Morita equivalent to A_λ° , checking the remaining axiom (S7) needed in order to have a based stratified algebra. \square

Corollary 5.36. *If A has a triangular decomposition in which the Cartan subalgebra A° is semisimple, then there is an idempotent refinement $A = \bigoplus_{i, j \in \hat{I}} e_i A e_j$ of A with the structure of a based quasi-hereditary algebra in the sense of Definition 5.1.*

Proof. The construction in the theorem produces an idempotent refinement of A that is a based stratified algebra with stratification function $\rho : \mathbf{B} \rightarrow \Lambda$. Let $\Gamma := \mathbf{B}$ with partial order \leq on Γ defined by $a \leq b$ if and only if $a = b$ or $\rho(a) < \rho(b)$. Since A_λ is basic and semisimple, we have for $a, b \in \mathbf{B}_\lambda$ that $H(a, b)$ is empty unless $a = b$ and $H(a, a)$ may be chosen to be $\{\hat{e}_a\}$. It follows that A is actually a based quasi-hereditary algebra with weight poset (Γ, \leq) and the basis which we have constructed. \square

Remark 5.37. The construction used to prove Theorem 5.35 suggests yet another variation on all of these definitions, which is weaker than having a triangular decomposition but stronger than having a triangular basis. For A like in Definition 5.31 we say that it has a *Cartan decomposition* if there is the following additional data:

- (CD1) A poset (Λ, \leq) which is upper finite in the locally finite-dimensional case and interval finite in the essentially finite-dimensional case.
- (CD2) A function $\partial : I \rightarrow \Lambda$ with finite fibers $I_\lambda := \partial^{-1}(\lambda)$.
- (CD3) A locally unital subalgebra A° and (A°, A°) -subbimodules A^{\flat} and A^{\sharp} of A .

The axioms are:

- (CD4) A^b is a projective right A° -module and A^\sharp is a projective left A° -module.
 (CD5) The natural multiplication map $A^b \otimes_{A^\circ} A^\sharp \rightarrow A$ is a linear isomorphism.
 (CD6) For $i, j \in I$, $e_i A^\circ e_j$ is zero unless $\partial(i) = \partial(j)$, $e_i A^b e_j$ and $e_j A^\sharp e_i$ are zero unless $\partial(i) \leq \partial(j)$, and $e_i A^b e_j = e_i A^\circ e_j = e_i A^\sharp e_j$ when $\partial(i) = \partial(j)$.

The proof of Theorem 5.35 carry over to such algebras essentially unchanged. However we do not know of any compelling examples, whereas as we noted in Remarks 5.30, 5.33 and 5.32 there are plenty of important examples of algebras with triangular bases and with triangular decompositions, justifying both of those definitions.

If A is a finite-dimensional (resp., locally finite-dimensional, resp. essentially finite-dimensional) algebra with a triangular decomposition, then we can apply Theorems 5.35 and 5.23 to deduce that $A\text{-mod}_{\text{fd}}$ (resp., $A\text{-mod}_{\text{lfid}}$, resp., $A\text{-mod}_{\text{efid}}$) is a finite (resp., upper finite, resp., essentially finite) fully stratified category. We end the section by making this structure more explicit. We first define some global standardization and costandardization functors.

- The axioms imply that A is a projective right A^\sharp -module and that there is a locally unital projection homomorphism $A^\sharp \rightarrow A^\circ$. Let

$$j_! : A^\circ\text{-mod}_{\text{fd}} \rightarrow A\text{-mod} \quad (5.13)$$

be the exact functor defined by first inflating along this projection homomorphism $A^\sharp \rightarrow A^\circ$ and then applying the exact induction functor $A \otimes_{A^\sharp} ? : A^\sharp\text{-mod} \rightarrow A\text{-mod}$. The fact that it takes finite-dimensional modules to finite-dimensional or locally finite-dimensional modules (as appropriate for the case) follows because as functors to $A^b\text{-mod}$ we have that $A \otimes_{A^\sharp} ? \cong A^b \otimes_{A^\circ} ?$ due to (TD5).

- The axioms imply that A is a projective left A^b -module and that there is a locally unital projection homomorphism $A^b \rightarrow A^\circ$. Let

$$j_* : A^\circ\text{-mod}_{\text{fd}} \rightarrow A\text{-mod} \quad (5.14)$$

be the exact functor defined by first inflating along the projection $A^b \rightarrow A^\circ$ then applying the exact coinduction functor $\bigoplus_{i \in I} \text{Hom}_{A^b}(Ae_i, -) : A^b\text{-mod} \rightarrow A\text{-mod}$. It takes finite-dimensional modules to finite-dimensional or locally finite-dimensional modules (as appropriate for the case) follows because as a functor to $A^\sharp\text{-mod}$ it is isomorphic to $\bigoplus_{i \in I} \text{Hom}_{A^\circ}(A^\sharp e_i, ?)$.

The following theorem can be proved by mimicking standard arguments from Lie theory; see [CouZ] noting that (A^b, A°) and (A°, A^\sharp) are *Borelic pairs* in the sense defined there. We will deduce it instead from the work already done in Theorems 5.35 and 5.23.

Theorem 5.38 (Fully stratified categories from algebras with a triangular decomposition). *Suppose that A has a triangular decomposition of one of the three types as above. Let $\{L^\circ(b) \mid b \in \mathbf{B}\}$ be a full set of pairwise inequivalent irreducible left A° -modules. Let $\rho : \mathbf{B} \rightarrow \Lambda$ be the function sending $b \in \mathbf{B}$ to the unique $\lambda \in \Lambda$ such that $L^\circ(b)$ is an irreducible A_λ° -module. Let $\bar{\Delta}(b) := j_! L^\circ(b)$ and $\bar{\nabla}(b) := j_* L^\circ(b)$; cf. (5.13)–(5.14). Then*

$$\{L(b) := \text{hd } \bar{\Delta}(b) \cong \text{soc } \bar{\nabla}(b) \mid b \in \mathbf{B}\}$$

is a complete set of pairwise inequivalent irreducible left A -modules. Moreover, the category $\mathcal{R} := A\text{-mod}_{\text{fd}}$ (resp., $A\text{-mod}_{\text{lfid}}$, resp., $A\text{-mod}_{\text{efid}}$) is a finite (resp., upper finite, resp., essentially finite) fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$. Its strata may be identified with the categories $A_\lambda^\circ\text{-mod}_{\text{fd}}$ ($\lambda \in \Lambda$) with standardization and costandardization functors defined by the restrictions of $j_!$ and j_ , respectively.*

Proof. As explained by Theorem 5.35, we can pass to an idempotent refinement if necessary to assume without loss of generality that the set I indexing the distinguished idempotents is disjoint from Λ and contains \mathbf{B} as a subset in such a way that $L^\circ(b) \cong \text{hd}(A^\circ e_b)$ for each $b \in \mathbf{B}$. The function $\rho : \mathbf{B} \rightarrow \Lambda$ is then the restriction of $\partial : I \rightarrow \Lambda$. Now Theorem 5.35 gives bases making A into a based stratified algebra. We deduce

that \mathcal{R} is a finite (resp., upper finite, resp., essentially finite) fully stratified category with stratification $(\mathbf{B}, L, \rho, \Lambda, \leq)$ by applying Theorem 5.23. However for this the strata and the labelling function L are produced in a different way to the formulation here, so we need to argue a little further to see that the standardization and costandardization functors here and the ones from earlier may be identified. Using the isomorphism (5.12), the quotient functor $j^\lambda : A_{\leq \lambda}\text{-mod} \rightarrow A_\lambda\text{-mod}$ in the setup of (5.4) may be identified with the functor $j : A_{\leq \lambda}\text{-mod} \rightarrow e_\lambda A_\lambda^\circ e_\lambda\text{-mod}$ obtained by restriction to A° then multiplication by the idempotent e_λ . Since A_λ° and $e_\lambda A_\lambda^\circ e_\lambda$ are Morita equivalent, we can instead use the algebra A_λ° to realize the stratum, and then this quotient functor gets replaced by the functor obtained by restriction to A° then multiplication by 1_λ . It remains to observe that the restrictions of $j_!$ and j_* to $A_\lambda^\circ\text{-mod}$ are left and right adjoint to this functor, respectively. \square

Corollary 5.39. *Suppose that A has a triangular decomposition of one of the three types and that its Cartan subalgebra A° is semisimple. Let $\{L^\circ(\gamma) \mid \gamma \in \Gamma\}$ be a full set of pairwise inequivalent irreducible left A° -modules. Let $\rho : \Gamma \rightarrow \Lambda$ be the function sending γ to the unique λ such that $L^\circ(\gamma)$ is an irreducible A_λ° -module. Then $\mathcal{R} := A\text{-mod}_{\text{fd}}$ (resp., $A\text{-mod}_{\text{fd}}$, resp. $A\text{-mod}_{\text{fd}}$) is a finite (resp., upper finite, resp., essentially finite) highest weight category with weight poset (Γ, \leq) for \leq defined by $\beta \leq \gamma$ if either $\beta = \gamma$ or $\rho(\beta) < \rho(\gamma)$. Its standard and costandard modules are $\Delta(\gamma) := j_! L^\circ(\gamma)$ and $\nabla(\gamma) := j_* L^\circ(\gamma)$ for $\gamma \in \Gamma$.*

Proof. This follows from the theorem and Corollary 5.36. \square

Remark 5.40. We end by mentioning one last variation on the definitions in this subsection. We say that a triangular decomposition of A as in Definition 5.31 is a *symmetric triangular decomposition* if in addition there is given a locally unital algebra anti-involution $\sigma : A \rightarrow A$ which leaves A° invariant and interchanges A^\sharp and A^\flat , such that for each $\lambda \in \Lambda$ the subalgebras $e_\lambda A^\lambda e_\lambda$ are σ_λ -symmetric in the sense of Definition 4.51, where σ_λ denotes the restriction of σ . Then there is an enhanced version of Theorem 5.35 making A into a symmetrically based stratified algebra, and an enhanced version of Theorem 5.38 making \mathcal{R} into a fully stratified category with a Chevalley duality $?^\circledast$. We omit the details.

6. EXAMPLES

In this section, we explain several examples. For the ones in §§6.5–6.7 we give very few details but have tried to indicate the relevant ingredients from the existing literature.

6.1. A finite-dimensional example via quiver and relations. Let A and B be the basic finite-dimensional algebras defined by the following quivers:

$$\begin{aligned}
 A \ (1 > 2) : & \quad s \begin{array}{c} \curvearrowright \\ \circ \\ \curvearrowright \end{array} 1 \xrightarrow{y} 2 \begin{array}{c} \curvearrowleft \\ \circ \\ \curvearrowleft \end{array} t & \quad \text{with relations } s^2 = 0, t^2 = 0, ty = 0, \\
 B \ (1 < 2) : & \quad z \begin{array}{c} \curvearrowright \\ \circ \\ \curvearrowleft \end{array} 1 \begin{array}{c} \xrightarrow{u} \\ \circ \\ \xleftarrow{v} \end{array} 2 & \quad \text{with relations } z^2 = 0, uv = 0, vuzv = 0.
 \end{aligned}$$

The algebra A has basis $e_1, s; e_2, t; y, ys$ and B has basis $e_1, z, vu, vuz, zvu, zvvuz; e_2, uzv; v, zv; u, uz, uzvu, uzvvuz$. The irreducible A - and B -modules are indexed by the set $\{1, 2\}$. We are going to consider $A\text{-mod}_{\text{fd}}$ and $B\text{-mod}_{\text{fd}}$ with the stratifications defined by the orders $1 > 2$ and $1 < 2$, respectively.

We first look at $A\text{-mod}_{\text{fd}}$. As usual, we denote its irreducibles by $L(i)$, indecomposable projectives by $P(i)$, standards by $\Delta(i)$, etc.. The indecomposable projectives and

injectives look as follows (where we abbreviate the irreducible module $L(i)$ just by i):

$$P(1) = \begin{array}{c} 1 \\ \swarrow s \quad \searrow y \\ 1 \quad 2 \\ \downarrow \psi \\ 2 \end{array}, \quad P(2) = \begin{array}{c} 2 \\ \downarrow t \\ 2 \end{array}, \quad I(1) = \begin{array}{c} 1 \\ \downarrow s \\ 1 \end{array}, \quad I(2) = \begin{array}{c} 1 \\ \downarrow y \\ 2 \quad 1 \\ \swarrow t \quad \searrow \psi \\ 2 \end{array}.$$

It follows easily that $A\text{-mod}_{\text{fd}}$ is a fibered highest weight category in the sense of Definition 3.7 with the structure of the standards and costandards as follows:

$$\Delta(1) = P(1), \quad \bar{\Delta}(1) = \begin{array}{c} 1 \\ \downarrow \psi \\ 2 \end{array}, \quad \Delta(2) = P(2), \quad \bar{\Delta}(2) = L(2),$$

$$\nabla(1) = I(1), \quad \bar{\nabla}(1) = L(1), \quad \nabla(2) = \begin{array}{c} 2 \\ \downarrow t \\ 2 \end{array}, \quad \bar{\nabla}(2) = L(2).$$

This can also be seen from Theorem 5.23 on noting that A is a based properly stratified algebra in the sense of Definition 5.21 with $Y(2,1) = \{y\}, X(1,2) = \emptyset$ and $H(1) = \{e_1, s\}, H(2) = \{e_2, t\}$. The basic local algebras realizing the strata are $\mathbb{k}[s]/(s^2)$ and $\mathbb{k}[t]/(t^2)$. Next we look at the tilting modules in $A\text{-mod}_{\text{fd}}$. If one takes the sign function $\varepsilon = (\varepsilon_1, \varepsilon_2)$ to be either $(+, +)$ or $(-, +)$ then one finds that the indecomposable ε -tilting modules are:

$$T_+(1) = P(1) = \begin{array}{c} \bar{\Delta}(1) \\ | \\ \bar{\Delta}(1) \end{array} = \begin{array}{c} \nabla(1) \\ \swarrow \quad \searrow \\ \bar{\nabla}(2) \quad \bar{\nabla}(2) \end{array}, \quad T_+(2) = P(2) = \Delta(2) = \begin{array}{c} \bar{\nabla}(2) \\ | \\ \bar{\nabla}(2) \end{array}.$$

These cases are not very interesting since the Ringel dual category is just $A\text{-mod}_{\text{fd}}$ again. Assume henceforth that $\varepsilon = (-, -)$ or $(+, -)$. Then the indecomposable ε -tilting modules are:

$$T_-(1) = \begin{array}{c} 1 \quad 2 \\ \downarrow \psi \quad \downarrow y \quad \downarrow t \\ 1 \quad 2 \quad 2 \\ \swarrow y \quad \swarrow t \\ 2 \end{array} = \begin{array}{c} \bar{\Delta}(2) \quad \bar{\Delta}(2) \\ \swarrow \quad \searrow \\ \Delta(1) \end{array} = \begin{array}{c} \nabla(1) \\ \swarrow \quad \searrow \\ \nabla(2) \quad \nabla(2) \end{array}, \quad T_-(2) = P(2).$$

To see this, one just has to check that these modules are indecomposable with the appropriate Δ_ε - and ∇_ε -flags. This analysis reveals that $A\text{-mod}_{\text{fd}}$ is *not* tilting-rigid.

The minimal projective resolution of $T_-(1)$ takes the form

$$\cdots \longrightarrow P(2) \oplus P(2) \longrightarrow P(2) \oplus P(2) \longrightarrow P(1) \oplus P(2) \oplus P(2) \longrightarrow T_-(1) \longrightarrow 0.$$

In particular, it is not of finite projective dimension, as follows also from Lemma 4.38 since $T_-(1) \not\cong T_+(1)$. Observe also that there is a non-split short exact sequence $0 \rightarrow X \rightarrow T_-(1) \rightarrow X \rightarrow 0$ where

$$X = \begin{array}{c} 1 \quad 2 \\ \downarrow \psi \quad \downarrow y \quad \downarrow t \\ 2 \end{array}.$$

Now let $T := T_-(1) \oplus T_-(2)$. We claim that $\text{End}_A(T)^{\text{op}}$ is the algebra B defined above. To prove this, one takes $z : T_-(1) \rightarrow T_-(1)$ to be an endomorphism whose image and kernel is the submodule X of $T_-(1)$, $u : T_-(2) \rightarrow T_-(1)$ to be a homomorphism which includes $T_-(2)$ as a submodule of $X \subseteq T_-(1)$, and $v : T_-(1) \rightarrow T_-(2)$ to be a homomorphism with kernel containing X and image $L(2) \subseteq T_-(2)$. Hence, $B\text{-mod}_{\text{fd}}$ is

the Ringel dual of $A\text{-mod}_{\text{fd}}$ relative to T . Note also that the algebra B is based $(+, +)$ - and $(-, +)$ -quasi-hereditary but it is *not* based $(+, -)$ - or $(-, -)$ -quasi-hereditary (cf. Definition 5.13).

One can also analyze $B\text{-mod}_{\text{fd}}$ directly. Its projective modules have the following structure:

$$\begin{array}{c}
 1 \\
 \swarrow \quad \searrow \\
 2 \qquad 1 \\
 \downarrow \quad \downarrow \\
 1 \qquad 2 \\
 \downarrow \quad \downarrow \\
 1 \qquad 1 \\
 \downarrow \quad \downarrow \\
 2 \qquad 1 \\
 \qquad \downarrow \\
 \qquad 2
 \end{array}
 , \qquad
 \begin{array}{c}
 2 \\
 \downarrow \\
 1 \\
 \downarrow \\
 1 \\
 \downarrow \\
 1 \\
 \downarrow \\
 2
 \end{array}
 ,$$

Continuing with $\varepsilon = (-, -)$ or $\varepsilon = (+, -)$, it is easy to check directly from this that $B\text{-mod}_{\text{fd}}$ is $(-\varepsilon)$ -highest weight, as we knew already due to Theorem 4.10. However, it is not ε -highest weight for either of these choices of ε , so it is *not* fibered highest weight.

We leave it to the reader to compute explicitly the indecomposable $(-\varepsilon)$ -tilting modules $T'_+(1)$ and $T'_+(2)$ in $B\text{-mod}_{\text{fd}}$. Their structure reflects the structure of the injectives $I(1)$ and $I(2)$ in $A\text{-mod}_{\text{fd}}$. Let $T' := T'_+(1) \oplus T'_+(2) \cong T^*$. By the double centralizer property from Corollary 4.11, we have that $A = \text{End}_B(T')^{\text{op}}$, as may also be checked directly. By Theorem 4.16, the functor $\mathbb{R} \text{Hom}_B(T', ?) : D^b(B\text{-mod}_{\text{fd}}) \rightarrow D^b(A\text{-mod}_{\text{fd}})$ is an equivalence. Note though that $\mathbb{R} \text{Hom}_A(T, ?) : D^b(A\text{-mod}_{\text{fd}}) \rightarrow D^b(B\text{-mod}_{\text{fd}})$ is *not* one; this follows using [Kel, Th. 4.1] since $T_-(1)$ does not have finite projective dimension.

6.2. An explicit semi-infinite example. In this subsection, we give a baby example involving a lower finite highest weight category. Let C be the coalgebra with basis

$$\{c_{i,j}^{(\ell)} \mid i, j, \ell \in \mathbb{Z}, 0 \leq i, j \leq \ell\},$$

count defined by $\epsilon(c_{i,j}^{(\ell)}) := \delta_{i,\ell} \delta_{j,\ell}$, and comultiplication $\delta : C \rightarrow C \otimes C$ defined by

$$\begin{aligned}
 c_{i,j}^{(i)} &\mapsto \sum_{\substack{k=0 \\ i \neq j(2)}}^j c_{i,k}^{(i)} \otimes c_{k,j}^{(j)} + \sum_{\substack{k=j \\ k \equiv i(2)}}^i c_{i,k}^{(i)} \otimes c_{k,j}^{(k)}, & c_{i,j}^{(j)} &\mapsto \sum_{\substack{k=0 \\ i \neq j(2)}}^i c_{i,k}^{(i)} \otimes c_{k,j}^{(j)} + \sum_{\substack{k=i \\ k \equiv j(2)}}^j c_{i,k}^{(k)} \otimes c_{k,j}^{(j)}, \\
 c_{i,j}^{(\ell)} &\mapsto c_{i,\ell}^{(\ell)} \otimes c_{\ell,j}^{(\ell)} + \sum_{\substack{k=0 \\ i \neq \ell(2)}}^i c_{i,k}^{(i)} \otimes c_{k,j}^{(\ell)} + \sum_{\substack{k=i \\ k \equiv \ell(2)}}^{\ell-1} c_{i,k}^{(k)} \otimes c_{k,j}^{(\ell)} + \sum_{\substack{k=0 \\ j \neq \ell(2)}}^j c_{i,k}^{(\ell)} \otimes c_{k,j}^{(j)} + \sum_{\substack{k=j \\ k \equiv \ell(2)}}^{\ell-1} c_{i,k}^{(\ell)} \otimes c_{k,j}^{(k)}
 \end{aligned}$$

for $i, j \geq 0$ and $\ell > \max(i, j)$. We will show that $\mathcal{R} := \text{comod}_{\text{fd}}\text{-}C$ is a lower finite highest weight category with weight poset $\Lambda := \mathbb{N}$ ordered in the natural way. Then we will determine the costandard, standard and indecomposable injective and tilting objects explicitly, and describe the Ringel dual category \mathcal{R}' . To do this, we mimic some arguments for reductive groups which we learnt from [Jan1].

We will need comodule induction functors, which we review briefly. For any coalgebra C with comultiplication δ , a right C -comodule V with structure map $\eta_R : V \rightarrow V \otimes C$, and a left C -comodule W with structure map $\eta_L : W \rightarrow C \otimes W$, the *cotensor product* $V \square_C W$ is the subspace of the vector space $V \otimes W$ that is the equalizer of the diagram

$$V \otimes W \begin{array}{c} \xrightarrow{\eta_R \otimes \text{id}} \\ \xrightarrow{\text{id} \otimes \eta_L} \end{array} V \otimes C \otimes W.$$

In particular, $\eta_R : V \rightarrow V \otimes C$ is an isomorphism from V to the subspace $V \square_C C$, and similarly $\eta_L : W \xrightarrow{\sim} C \square_C W$. Now suppose that $\pi : C \rightarrow C'$ is a coalgebra homomorphism and V is a right C' -comodule. Viewing C as a left C' -comodule with structure map $\delta_L := (\pi \otimes \text{id}) \circ \delta : C \rightarrow C' \otimes C$, we define the induced comodule to be

$$\text{ind}_{C'}^C V := V \square_{C'} C.$$

This is a subcomodule of the right C -comodule $V \otimes C$ (with structure map $\text{id} \otimes \delta$). In fact, $\text{ind}_{C'}^C : \text{comod-}C' \rightarrow \text{comod-}C$ defines a functor which is right adjoint to the exact restriction functor $\text{res}_{C'}^C$, so it is left exact and sends injectives to injectives.

Now let C be the coalgebra defined above, and consider the natural quotient maps $\pi^b : C \rightarrow C^b$ and $\pi^\sharp : C \rightarrow C^\sharp$, where C^b and C^\sharp are the quotients of C by the coideals spanned by $\{c_{i,j}^{(\ell)} \mid \ell > j\}$ or $\{c_{i,j}^{(\ell)} \mid \ell > i\}$, respectively. These coalgebras have bases denoted $\{c_{i,j} := \pi^b(c_{i,j}^{(j)}) \mid 0 \leq i \leq j\}$ and $\{c_{i,j} := \pi^\sharp(c_{i,j}^{(i)}) \mid i \geq j \geq 0\}$, and comultiplications δ^b and δ^\sharp satisfying

$$\delta^b(c_{i,j}) = c_{i,i} \otimes c_{i,j} + \sum_{\substack{k=i+1 \\ k \equiv j(2)}}^j c_{i,k} \otimes c_{k,j}, \quad \delta^\sharp(c_{i,j}) = c_{i,j} \otimes c_{j,j} + \sum_{\substack{k=j+1 \\ k \equiv i(2)}}^i c_{i,k} \otimes c_{k,j}, \quad (6.1)$$

respectively. Also let $C^\circ \cong \bigoplus_{i \geq 0} \mathbb{k}$ be the semisimple coalgebra with basis $\{c_i \mid i \geq 0\}$ and comultiplication $\delta^\circ : c_i \mapsto c_i \otimes c_i$. Note C° is a quotient of both C^b and C^\sharp via the obvious maps sending $c_{i,j} \mapsto \delta_{i,j} c_i$; hence, it is also a quotient of C . It may also be identified with a subcoalgebra of both C^b and C^\sharp via the maps sending $c_i \mapsto c_{i,i}$.

Let $L^\circ(i)$ be the one-dimensional irreducible right C° -comodule spanned by $c_{i,i}$. Since C° is semisimple with these as its irreducible comodules, any irreducible right C° -comodule V decomposes as $V = \bigoplus_{i \in I} V_i$ with the ‘‘weight spaces’’ V_i being a direct sum of copies of $L^\circ(i)$. Similarly, any left C° -comodule V decomposes as $V = \bigoplus_{i \in I} {}_i V$. This applies in particular to left and right C^b, C^\sharp or C -comodules, since these may be viewed as C° -comodules by restriction.

Since C° is a subcoalgebra of C^b , the irreducible comodule $L^\circ(i)$ may also be viewed as an irreducible right C^b -comodule. We denote this instead by $L^b(i)$; it is the subcomodule of C^b spanned by the vector $c_{i,i}$. For $i \geq 0$, let $I(i) := {}_i C \cong \text{ind}_{C^\circ}^C L^\circ(i)$, let $\nabla(i)$ be the subcomodule of $I(i)$ spanned by the vectors $\{c_{i,j}^{(i)} \mid 0 \leq j \leq i\}$, and let $L(i)$ be the one-dimensional irreducible subcomodule of $\nabla(i)$ spanned by the vector $c_{i,i}^{(i)}$. Now we proceed in several steps.

Claim 1: Viewed as a functor to vector spaces, the induction functor $\text{ind}_{C^b}^C$ is isomorphic to the functor $V \mapsto V \square_{C^\circ} C^\sharp \cong \bigoplus_{i \geq 0} V_i \otimes {}_i C^\sharp$. Hence, this functor is exact. To prove this, let $\delta_{LR} := (\pi^b \otimes \pi^\sharp) \circ \delta : C \rightarrow C^b \square_{C^\circ} C^\sharp$. As $\delta_{LR}(c_{i,j}^{(\ell)}) = c_{i,\ell} \otimes c_{\ell,j}$ and these vectors for all $\ell \geq \max(i, j)$ give a basis for $C^b \square_{C^\circ} C^\sharp$, this map is a linear isomorphism. Moreover, the following diagram commutes:

$$\begin{array}{ccc} C & \xrightarrow{\delta_L} & C^b \otimes C \\ \delta_{LR} \downarrow & & \downarrow \text{id} \otimes \delta_{LR} \\ C^b \square_{C^\circ} C^\sharp & \xrightarrow{\delta^b \otimes \text{id}} & C^b \otimes C^b \square_{C^\circ} C^\sharp. \end{array}$$

The vertical maps are isomorphisms. Using the definition of $\text{ind}_{C^b}^C$, it follows for any right C^b -comodule V with structure map η that the induced module $\text{ind}_{C^b}^C V$ is isomorphic as a vector space (indeed, as a right C^\sharp -comodule) to the equalizer of the diagram

$$V \otimes C^b \square_{C^\circ} C^\sharp \begin{array}{c} \xrightarrow{\eta \otimes \text{id} \otimes \text{id}} \\ \xrightarrow{\text{id} \otimes \delta^b \otimes \text{id}} \end{array} V \otimes C^b \otimes C^b \square_{C^\circ} C^\sharp.$$

Since $\text{ind}_{C^b}^{C^b} V \cong V$, this is naturally isomorphic to $V \square_{C^\circ} C^\sharp$. As $C^\sharp \cong \bigoplus_{i \geq 0} iC^\sharp$, we get finally that $V \square_{C^\circ} C^\sharp \cong \bigoplus_{i \geq 0} V_i \otimes_i C^\sharp$.

Claim 2: For $i \geq 0$, the right C^b -comodule ${}_i C^b \cong \text{ind}_{C^\circ}^{C^b} L^\circ(i)$ has an exhaustive ascending filtration $0 < V_0 < V_1 < \dots$ such that $V_0 \cong L^b(i)$ and $V_r/V_{r-1} \cong L^b(i+2r-1) \oplus L^b(i+2r)$ for $r \geq 1$. Also, the modules $\{L^b(i) \mid i \geq 0\}$ give a full set of pairwise inequivalent irreducible right C^b -comodules. The first statement follows from (6.1), defining V_0 to be the subspace spanned by $c_{i,i}$, and V_r is spanned by $c_{i,i+2r-1}, c_{i,i+2r}$. To prove the second statement, take any irreducible C^b -comodule L . Take a non-zero homomorphism $\text{res}_{C^\circ}^{C^b} L \rightarrow L^\circ(i)$ for some i . Then use adjointness of $\text{res}_{C^\circ}^{C^b}$ and $\text{ind}_{C^\circ}^{C^b}$ to obtain an embedding $L \hookrightarrow {}_i C^b$. Hence, $L \cong L^b(i)$ as a C^b -comodule.

Claim 3: We have that $\nabla(i) \cong \text{ind}_{C^b}^{C^b} L^b(i)$ and it is uniserial with composition factors $L(i), L(i-2), L(i-4), \dots, L(a), L(b), \dots, L(i-3), L(i-1)$ (for $(a, b) \in \{(0, 1), (1, 0)\}$ depending on parity of i) in order from bottom to top:

$$\nabla(i) = \begin{array}{c} i-1 \\ | \\ i-3 \\ \vdots \\ i-2 \\ | \\ i \end{array} \tag{6.2}$$

The restriction of $\delta_L : C \rightarrow C^b \otimes C$ to $\nabla(i)$ gives an embedding of $\nabla(i)$ into $\text{ind}_{C^b}^{C^b} L^b(i)$. This embedding is an isomorphism since we know $\text{ind}_{C^b}^{C^b} L^b(i)$ has the same dimension $(i+1)$ as $\nabla(i)$ thanks to Claim 1. The determinant of the subcomodule structure is straightforward using the definition of $\delta(c_{i,j}^{(i)})$ for $0 \leq j \leq i$.


Claim 4: The injective C -comodule $I(i)$ has an exhaustive filtration $0 < I_0 < I_1 < \dots$ such that $I_0 \cong \nabla(i)$ and $I_r/I_{r-1} \cong \nabla(i+2r-1) \oplus \nabla(i+2r)$ for $r \geq 1$:

$$I(i) = \begin{array}{ccc} & \begin{array}{c} \vdots \\ \nabla(i+3) \\ | \\ \nabla(i+1) \end{array} & \begin{array}{c} \vdots \\ \nabla(i+4) \\ | \\ \nabla(i+2) \end{array} \\ & \diagdown & \diagup \\ & \nabla(i) & \end{array} \tag{6.3}$$

This follows from Claims 1, 2 and 3.

Claim 5: The C -comodules $\{L(i) \mid i \geq 0\}$ give a full set of pairwise inequivalent irreducibles. Moreover, $I(i)$ is the injective hull of $L(i)$. By Claim 3, the last part of Claim 2, and an adjunction argument, any irreducible C -comodule embeds into $\nabla(i)$ for some i , hence, it is isomorphic to $L(i)$. The module $I(i)$ is injective, and it has irreducible socle $L(i)$ by another adjunction argument. Hence, it is the injective hull of $L(i)$.

Claim 6: The category $\mathcal{R} := \text{comod}_{\text{id}} C$ is a lower finite highest weight category with costandard objects $\nabla(i)$ ($i \geq 0$). It also possesses a Chevalley duality. We use the criterion from Corollary 3.61. From Claim 4, it follows that the largest submodule of $I(i)$ that belongs to $\mathcal{R}_{\leq i}$ is $\nabla(i)$, which is finite-dimensional. This shows that $\mathcal{R}_{\leq i}$ has enough injectives with the injective hull of $L(i)$ being $\nabla(i)$. We also know already that $[\nabla(i) : L(i)] = 1$, and the property $(I\widehat{\nabla}^{\text{asc}})$ follows from Claim 4. Hence, \mathcal{R} is a lower finite highest weight category. Finally, the Chevalley duality is defined using the evident coalgebra antiautomorphism of C which maps $c_{i,j}^{(\ell)} \mapsto c_{j,i}^{(\ell)}$.



Remark 6.2. The above example can be changed slightly to obtain an essentially finite example with weight poset $\Lambda := \mathbb{Z}$ ordered by the opposite of the natural ordering. To do this, let D be the essentially finite-dimensional locally unital algebra defined by the following quiver:

$$D : \cdots -1 \begin{array}{c} \xrightarrow{y_{-1}} \\ \xleftarrow{x_{-1}} \end{array} 0 \begin{array}{c} \xrightarrow{y_0} \\ \xleftarrow{x_0} \end{array} 1 \begin{array}{c} \xrightarrow{y_1} \\ \xleftarrow{x_1} \end{array} 2 \cdots \quad \text{with relations } y_{i+1}y_i = x_i x_{i+1} = x_i y_i = 0.$$

Like for A , this algebra has a triangular decomposition, so $D\text{-mod}_{\text{fd}}$ is an essentially finite highest weight category. Recalling that the construction of tilting modules in the essentially finite case explained in §4.5 is by passing to an upper finite truncation, the indecomposable tilting module $T(0)$ for D has the same structure as for A ; see (6.6). This module is infinite-dimensional; thus $D\text{-mod}_{\text{fd}}$ is *not* tilting-bounded. Note also that this algebra D can be obtained from the general construction from Remark 5.33, starting from the obvious triangular decomposition of the \mathbb{Z} -graded algebra $\bar{A} = \mathbb{k}\langle x, y \mid x^2 = y^2 = 0, xy = 0 \rangle$ with x in degree 1 and y in degree -1 ; cf. [BT, Ex. 5.12].

6.3. Category \mathcal{O} for affine Lie algebras. Perhaps the first naturally-occurring examples of finite highest weight categories came from the blocks of the BGG category \mathcal{O} for a semisimple Lie algebra. This context also provides natural examples of finite fibered highest weight categories; see [Maz1] for a survey. To get examples of *semi-infinite* highest weight categories, one can consider instead blocks of the category \mathcal{O} for an affine Kac-Moody Lie algebra. We briefly recall the setup referring to [Kac], [Car] for more details.

Let \mathfrak{g} be a finite-dimensional semisimple Lie algebra over \mathbb{C} and

$$\mathfrak{g} := \mathring{\mathfrak{g}} \otimes_{\mathbb{C}} \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c \oplus \mathbb{C}d$$

be the corresponding affine Kac-Moody algebra. Fix also a Cartan subalgebra \mathfrak{h} contained in a Borel subalgebra \mathfrak{b} of $\mathring{\mathfrak{g}}$. There are corresponding subalgebras \mathfrak{h} and \mathfrak{b} of \mathfrak{g} , namely,

$$\mathfrak{h} := \mathring{\mathfrak{h}} \oplus \mathbb{C}c \oplus \mathbb{C}d, \quad \mathfrak{b} := \left(\mathring{\mathfrak{b}} \otimes_{\mathbb{C}} \mathbb{C}[t] + \mathring{\mathfrak{g}} \otimes_{\mathbb{C}} t\mathbb{C}[t] \right) \oplus \mathbb{C}c \oplus \mathbb{C}d.$$

Let $\{\alpha_i \mid i \in I\} \subset \mathfrak{h}^*$ and $\{h_i \mid i \in I\} \subset \mathfrak{h}$ be the simple roots and coroots of \mathfrak{g} and $(\cdot | \cdot)$ be the normalized invariant form on \mathfrak{h}^* , all as in [Kac, Ch. 7–8]. The *basic imaginary root* $\delta \in \mathfrak{h}^*$ is the positive root corresponding to the canonical central element $c \in \mathfrak{h}$ under $(\cdot | \cdot)$. The linear automorphisms of \mathfrak{h}^* defined by $s_i : \lambda \mapsto \lambda - \lambda(h_i)\alpha_i$ generate the Weyl group W of \mathfrak{g} . Let $\rho \in \mathfrak{h}^*$ be the element satisfying $\rho(h_i) = 1$ for all $i \in I$ and $\rho(d) = 0$. Then define the shifted action of W on \mathfrak{h}^* by $w \cdot \lambda = w(\lambda + \rho) - \rho$ for $w \in W$, $\lambda \in \mathfrak{h}^*$.

We define the *level* of $\lambda \in \mathfrak{h}^*$ to be $(\lambda + \rho)(c) \in \mathbb{C}$. It is *critical* if it equals the level of $\lambda = -\rho$, i.e., it is zero¹². We usually restrict our attention to integral weights λ , that is,

¹²Many authors define the level to be $\lambda(c)$, in which case the critical level is $-\check{h}$, where \check{h} is the dual Coxeter number.

weights $\lambda \in \mathfrak{h}^*$ such that $\lambda(h_i) \in \mathbb{Z}$ for all $i \in I$. The level of an integral weight is either *positive*, *negative* or *critical* (= zero). For any $\lambda \in \mathfrak{h}^*$, we define

$$\lambda' := -\lambda - 2\rho. \quad (6.7)$$

Since $w \cdot (-\lambda - 2\rho) = -w \cdot \lambda - 2\rho$, weights λ and μ are in the same orbit under the shifted action of W if and only if so are λ' and μ' . Note also that the level of λ is positive (resp., critical) if and only if the level of λ' is negative (resp., critical). A crucial fact is that the orbit $W \cdot \lambda$ of an integral weight λ of positive level contains a unique weight λ_{\max} such that $\lambda_{\max} + \rho$ is dominant; e.g., see [Kum, Ex. 13.1.E8a, Prop. 1.4.2]. By [Kum, Cor. 1.3.22], this weight is maximal in its orbit with respect to the usual dominance ordering \leq on weights, i.e., $\mu \leq \lambda$ if $\lambda - \mu \in \bigoplus_{i \in I} \mathbb{N}\alpha_i$. If λ is integral of negative level, we deduce from this discussion that its orbit contains a unique minimal weight λ_{\min} .

For $\lambda \in \mathfrak{h}^*$, let $\Delta(\lambda)$ be the Verma module with highest weight λ and $L(\lambda)$ be its unique irreducible quotient. Although Verma modules need not be of finite length, the composition multiplicities $[\Delta(\lambda) : L(\mu)]$ are always finite. There is also the *dual Verma module* $\nabla(\lambda)$ which is the restricted dual $\Delta(\lambda)^\#$ of $\Delta(\lambda)$, i.e., the sum of the duals of the weight spaces of $\Delta(\lambda)$ with the \mathfrak{g} -action twisted by the Chevalley antiautomorphism. All of the modules just introduced are objects in the category \mathcal{O} consisting of all \mathfrak{g} -modules M which are semisimple over \mathfrak{h} with finite-dimensional weight spaces and such that the set of weights of M is contained in the lower set generated by a finite subset of \mathfrak{h}^* ; see [Kum, §2.1]. There is also a larger category $\hat{\mathcal{O}}$ consisting of the \mathfrak{g} -modules M which are semisimple over \mathfrak{h} and locally finite-dimensional over \mathfrak{b} .

Let \sim be the equivalence relation on \mathfrak{h}^* generated by $\lambda \sim \mu$ if there exists a positive root γ and $n \in \mathbb{Z}$ such that $2(\lambda + \rho|\gamma) = n(\gamma|\gamma)$ and $\lambda - \mu = n\gamma$. For a \sim -equivalence class Λ , let \mathcal{O}_Λ (resp., $\hat{\mathcal{O}}_\Lambda$) be the full subcategory of \mathcal{O} (resp., $\hat{\mathcal{O}}$) consisting of all $M \in \mathcal{O}$ (resp., $M \in \hat{\mathcal{O}}$) such that $[M : L(\lambda)] \neq 0 \Rightarrow \lambda \in \Lambda$. In view of the linkage principle from [KK, Th. 2], these subcategories may be called the *blocks* of \mathcal{O} and of $\hat{\mathcal{O}}$, respectively. In particular, by [DGK, Th. 4.2], any $M \in \mathcal{O}$ decomposes uniquely as a direct sum $M = \bigoplus_{\Lambda \in \mathfrak{h}^*/\sim} M_\Lambda$ with $M_\Lambda \in \mathcal{O}_\Lambda$. Note though that \mathcal{O} is not the coproduct of its blocks in the strict sense since it is possible to find $M \in \mathcal{O}$ such that M_Λ is non-zero for infinitely many different Λ . The situation is more satisfactory for $\hat{\mathcal{O}}$: $\hat{\mathcal{O}}$ is the product of its blocks since by [Soe, Th. 6.1] the functor


$$\prod_{\Lambda \in \mathfrak{h}^*/\sim} \hat{\mathcal{O}}_\Lambda \rightarrow \hat{\mathcal{O}}, \quad (M_\Lambda)_{\Lambda \in \mathfrak{h}^*/\sim} \mapsto \bigoplus_{\Lambda \in \mathfrak{h}^*/\sim} M_\Lambda \quad (6.8)$$

is an equivalence of categories. Note also that $[\Delta(\lambda) : L(\mu)] \neq 0$ implies that the level of λ equals that of μ , since the scalars by which c acts on $L(\lambda)$ and $L(\mu)$ must agree. Consequently, we can talk simply about the level of a block.

A general combinatorial description of the \sim -equivalence classes Λ can be found for instance in [Fie3, Lem. 3.9]. For simplicity, we restrict ourselves from now on to integral blocks. In non-critical levels, one gets exactly the W -orbits $W \cdot \lambda$ of the integral weights of non-critical level. In critical level, one needs to incorporate also the translates by $\mathbb{Z}\delta$. From this description, it follows that the poset (Λ, \leq) underlying an integral block \mathcal{O}_Λ is upper finite with unique maximal element λ_{\max} if \mathcal{O}_Λ is of positive level, and lower finite with unique minimal element λ_{\min} if \mathcal{O}_Λ is of negative level. In case of the critical level, the poset is neither upper finite nor lower finite, but it is always interval finite.

Example 6.3. Here we give some explicit examples of posets which can occur for $\mathfrak{g} = \mathfrak{sl}_2$, the Kac-Moody algebra for the Cartan matrix $\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$. The labelling set for the principal block is $W \cdot 0 = \{\lambda_k, \mu_k \mid k \geq 0\}$ where $\lambda_k := -\frac{1}{2}k(k+1)\alpha_0 - \frac{1}{2}k(k-1)\alpha_1$ and $\mu_k := -\frac{1}{2}k(k-1)\alpha_0 - \frac{1}{2}k(k+1)\alpha_1$. This is a block of positive level with maximal element $\lambda_0 = \mu_0 = 0$. Applying the map (6.7), we deduce that $W \cdot (-2\rho) = \{\lambda'_k, \mu'_k \mid k \geq 0\}$. This is the labelling set for a block of negative level with minimal element $\lambda'_0 = \mu'_0 = -2\rho$. Finally, we have that $W \cdot (\alpha_0 - \rho) \sqcup W \cdot (\alpha_1 - \rho) = \{\bar{\lambda}_k, \bar{\mu}_k \mid k \in \mathbb{Z}\}$

where $\bar{\lambda}_k := (k + 1)\alpha_0 + k\alpha_1 - \rho$ and $\bar{\mu}_k := k\alpha_0 + (k + 1)\alpha_1 - \rho$. This is the labelling set for a block of critical level, and it is neither upper nor lower finite.



Recall the definitions of upper finite and lower finite highest weight categories from Definitions 3.34 and 3.50, respectively.

Theorem 6.4. *Let \mathcal{O}_Λ be an integral block of \mathcal{O} of non-critical level. Then it is an upper finite or lower finite highest weight category according to whether the level is positive or negative, respectively. In both cases, the standard and costandard objects are the Verma modules $\Delta(\lambda)$ and the dual Verma modules $\nabla(\lambda)$, respectively, for $\lambda \in \Lambda$. The partial order \leq on Λ is the dominance order.*

Proof. First, we prove the result for an integral block \mathcal{O}_Λ of positive level. As explained above, the poset Λ is upper finite in this case. Let λ_{\max} be its unique maximal weight.

Claim 1: *In the positive level case, \mathcal{O}_Λ is the full subcategory of $\widehat{\mathcal{O}}_\Lambda$ consisting of all modules M such that $[M : L(\lambda)] < \infty$ for all $\lambda \in \Lambda$.* To prove this, given $M \in \mathcal{O}_\Lambda$, it is obvious that all of its composition multiplicities are finite since M has finite-dimensional weight spaces. Conversely, suppose that all of the composition multiplicities of $M \in \widehat{\mathcal{O}}_\Lambda$ are finite. All weights of M lie in the lower set generated by λ_{\max} . Moreover, for $\lambda \leq \lambda_{\max}$, the dimension of the λ -weight space of M is

$$\dim M_\lambda = \sum_{\mu \in \Lambda} [M : L(\mu)] \dim L(\mu)_\lambda.$$

Since the poset is upper finite, there are only finitely many $\mu \in \Lambda$ such that the λ -weight space $L(\mu)_\lambda$ is non-zero, and these weight spaces are finite-dimensional, so we deduce that $\dim M_\lambda < \infty$. This proves the claim.

Now we observe that the Verma module $M(\lambda_{\max})$ with maximal possible highest weight is projective in $\widehat{\mathcal{O}}_\Lambda$. From this and a standard argument involving translation functors through walls (see e.g. [Nei]) and the combinatorics from [Fie1, §4] (see also the introduction of [Fie2]), it follows that there are projective modules $P_\lambda \in \widehat{\mathcal{O}}_\Lambda$ with (finite) Δ -flags as in the axiom $(\widehat{P\Delta})$. Since each $\Delta(\lambda)$ belongs to \mathcal{O}_Λ , we actually have that $P_\lambda \in \mathcal{O}_\Lambda$. All that is left to complete the proof of the theorem in the positive level case is to show that \mathcal{O}_Λ is a Schurian category. Let $A := \left(\bigoplus_{\lambda, \mu \in \Lambda} \text{Hom}_{\mathfrak{g}}(P_\lambda, P_\mu) \right)^{\text{op}}$. Since the multiplicities $[P_\mu : L(\lambda)]$ are finite, A is a locally finite-dimensional locally unital algebra. Using Lemma 2.4, we deduce that $\widehat{\mathcal{O}}_\Lambda$ is equivalent to the category $A\text{-mod}$ of all left A -modules. As explained in the discussion after (2.22), $A\text{-mod}_{\text{lfid}}$ is the full subcategory of $A\text{-mod}$ consisting of all modules with finite composition multiplicities. Combining this with Claim 1, we deduce that the equivalence between $\widehat{\mathcal{O}}_\Lambda$ and $A\text{-mod}$ restricts to an equivalence between \mathcal{O}_Λ and $A\text{-mod}_{\text{lfid}}$. Hence, \mathcal{O}_Λ is a Schurian category.

We turn our attention to an integral block \mathcal{O}_Λ of negative level. In this case, we know already that the poset Λ is lower finite with a unique minimal element λ_{\min} .

Claim 2: *In the negative level case, the category \mathcal{O}_Λ is the full subcategory of $\widehat{\mathcal{O}}_\Lambda$ consisting of all modules of finite length.* For this, it is obvious that any module in $\widehat{\mathcal{O}}_\Lambda$ of finite length belongs to \mathcal{O}_Λ . Conversely, any object in \mathcal{O}_Λ is of finite length thanks to the formula [Kum, 2.1.11(1)], taking λ therein to be λ_{\min} .

From Claim 2 and Lemma 2.1, it follows that $\mathcal{R} := \mathcal{O}_\Lambda$ is a locally finite Abelian category. By [Fie1, Th. 2.7] the Serre subcategory \mathcal{R}^\downarrow of \mathcal{R} associated to Λ^\downarrow is a finite highest weight category for each finite lower set Λ^\downarrow of Λ . We deduce that \mathcal{R} is a lower finite highest weight category according to Definition 3.50. \square

Let \mathcal{O}_Λ be an integral block of non-critical level. The following assertions about projective and injective modules follow from Theorem 6.4 and the general theory from §§2.1–2.3; see also [Soe, Rem. 6.5].

- In the positive level case, when \mathcal{O}_Λ is a Schurian category, $\widehat{\mathcal{O}}_\Lambda$ has enough projectives and injectives. Moreover, the projective covers of the irreducible modules are the modules $\{P(\lambda) \mid \lambda \in \Lambda\}$ constructed in the proof of Theorem 6.4, and these belong to \mathcal{O}_Λ . Their restricted duals $I(\lambda) := P(\lambda)^\#$ are the indecomposable injective modules in $\widehat{\mathcal{O}}_\Lambda$, and also belong to \mathcal{O}_Λ .
- The situation is completely different in the negative level case, as we need to pass to $\widehat{\mathcal{O}}_\Lambda$, which is the ind-completion of the finite Abelian category \mathcal{O}_Λ , before we can talk about injective modules. In $\widehat{\mathcal{O}}_\Lambda$, the irreducible module $L(\lambda)$ ($\lambda \in \Lambda$) has an injective hull $I(\lambda)$ in $\widehat{\mathcal{O}}_\Lambda$, which possesses a (possibly infinite) ascending ∇ -flag in the sense of Definition 3.52. However, $\widehat{\mathcal{O}}_\Lambda$ usually does not have any projectives at all (although one could construct such modules in the pro-completion of \mathcal{O}_Λ as done e.g. in [Fie2]).

The following results about tilting modules are consequences of the general theory developed in §4.1 and §4.3. They already appeared in an equivalent form in [Soe].

- In the negative level case, tilting modules are objects in \mathcal{O}_Λ admitting both a (finite) Δ -flag and a (finite) ∇ -flag. The isomorphism classes of indecomposable tilting modules in \mathcal{O}_Λ are parametrized by their highest weights. They may also be constructed by applying translation functors to the Verma module $\Delta(\lambda_{\min})$.
- In the positive level case, tilting modules are objects in \mathcal{O}_Λ which admit both a (possibly infinite) ascending Δ -flag and a (possibly infinite) descending ∇ -flag in the sense of Definition 3.35. Again, the isomorphism classes of indecomposable tilting modules are parametrized by their highest weights.

In both cases, our characterization of the indecomposable tilting module $T(\lambda)$ of highest weight λ is slightly different from the one given in [Soe, Def. 6.3]. From our definition, one sees immediately that $T(\lambda)^\# \cong T(\lambda)$.

Remark 6.5. Elsewhere in the literature dealing with positive level, it is common to pass to a different category of modules, e.g., to the Whittaker category in [BY] or to truncated versions of \mathcal{O} in [SVV, §3], before contemplating tilting modules.

Our next result is concerned with the Ringel duality between integral blocks of positive and negative level. This depends crucially on a special case of the Arkhipov-Soergel equivalence from [Ark], [Soe]. Let S be Arkhipov's semi-regular bimodule, which is the bimodule S_γ of [Soe] with $\gamma := 2\rho$ as in [Soe, Lem. 7.1]. For $\lambda \in \mathfrak{h}^*$, let $T(\lambda)$ be the indecomposable tilting module from [Soe, Def. 6.3] (which is the same as in the previous paragraph for integral λ of positive or negative level). Also let $P(\lambda)$ be a projective cover of $L(\lambda)$ in $\widehat{\mathcal{O}}$ whenever such an object exists; cf. [Soe, Rem. 6.5(2)].

Theorem 6.6 (Arkhipov-Soergel equivalence). *Tensoring with the semi-regular bimodule defines an equivalence $S \otimes_{U(\mathfrak{g})}^? : \Delta(\mathcal{O}) \rightarrow \nabla(\mathcal{O})$ between the exact subcategories of \mathcal{O} consisting of objects with (finite) Δ - and ∇ -flags, respectively. Moreover:*

- (1) $S \otimes_{U(\mathfrak{g})} \Delta(\lambda) \cong \nabla(\lambda')$.
- (2) $S \otimes_{U(\mathfrak{g})} P(\lambda) \cong T(\lambda')$ (assuming $P(\lambda)$ exists).

Corollary 6.7. *Assume that \mathcal{O}_Λ is an integral block of negative level. Let \mathcal{O}'_Λ be the Ringel dual of \mathcal{O}_Λ relative to some choice of $T = \bigoplus_{i \in I} T_i$ as in Definition 4.24, and let F be the Ringel duality functor from (4.14). Also let $\Lambda' := \{\lambda' \mid \lambda \in \Lambda\}$. Then there is an equivalence of categories $E : \mathcal{O}'_\Lambda \rightarrow \mathcal{O}_{\Lambda'}$ such that $E \circ F : \nabla(\mathcal{O}_\Lambda) \rightarrow \Delta(\mathcal{O}_{\Lambda'})$ is a quasi-inverse to the Arkhipov-Soergel equivalence $S \otimes_{U(\mathfrak{g})}^? : \Delta(\mathcal{O}_{\Lambda'}) \rightarrow \nabla(\mathcal{O}_\Lambda)$.*

Proof. Note to start with that $\mathcal{O}_{\Lambda'}$ is an integral block of positive level. Moreover, the map $(\Lambda, \geq) \rightarrow (\Lambda', \leq), \lambda \mapsto \lambda'$ is an order isomorphism.

Choose a quasi-inverse D to $S \otimes_{U(\mathfrak{g})}^? : \Delta(\mathcal{O}_{\Lambda'}) \rightarrow \nabla(\mathcal{O}_\Lambda)$, and set $P_i := DT_i$. By Theorem 6.6(2), $(P_i)_{i \in I}$ is a projective generating family for $\mathcal{O}_{\Lambda'}$. Moreover, recalling that \mathcal{O}'_Λ is the category $A\text{-mod}_{\text{fhd}}$ where $A := \left(\bigoplus_{i,j \in I} \text{Hom}_{\mathcal{O}_\Lambda}(T_i, T_j) \right)^{\text{op}}$, the functor D induces an isomorphism via which we can identify A with $\left(\bigoplus_{i,j \in I} \text{Hom}_{\mathcal{O}_{\Lambda'}}(P_i, P_j) \right)^{\text{op}}$.

As explained in the proof of Theorem 6.4, the functor

$$H := \bigoplus_{i \in I} \text{Hom}_{\mathcal{O}_{\Lambda'}}(P_i, ?) : \mathcal{O}_{\Lambda'} \rightarrow A\text{-mod}_{\text{fhd}}$$

is an equivalence of categories. Moreover, we have that

$$H \circ D = \bigoplus_{i \in I} \text{Hom}_{\mathcal{O}_{\Lambda'}}(P_i, D?) \cong \bigoplus_{i \in I} \text{Hom}_{\mathcal{O}_\Lambda}(S \otimes_{U(\mathfrak{g})} P_i, ?) \cong \bigoplus_{i \in I} \text{Hom}_{\mathcal{O}_\Lambda}(T_i, ?) = F.$$

Letting E be a quasi-inverse equivalence to H , it follows that $E \circ F \cong D$. \square

Remark 6.8. In the setup of Corollary 6.7, the Arkhipov-Soergel equivalence extends to an equivalence $S \otimes_{U(\mathfrak{g})}^? : \Delta^{\text{asc}}(\mathcal{O}_{\Lambda'}) \rightarrow \nabla^{\text{asc}}(\mathcal{O}_\Lambda)$, which is a quasi-inverse to $E \circ F : \nabla^{\text{asc}}(\mathcal{O}_\Lambda) \rightarrow \Delta^{\text{asc}}(\mathcal{O}_{\Lambda'})$. These functors interchange the indecomposable injectives in $\widehat{\mathcal{O}}_\Lambda$ with the indecomposable tiltings in $\mathcal{O}_{\Lambda'}$.

Finally we discuss the situation for an integral critical block \mathcal{O}_Λ . As we have already explained, in this case the poset Λ is neither upper nor lower finite. In fact, these blocks do not fit into the framework of this article at all, since the Verma modules have infinite length and there are no projectives. One sees this already for the Verma module $\Delta(-\rho)$ for $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, which has composition factors $L(-\rho - m\delta)$ for $m \geq 0$, each appearing with multiplicity equal to the number of partitions of m ; see e.g. [AF1, Th. 4.9(1)]. However, there is an autoequivalence $\Sigma := L(\delta) \otimes^? : \widehat{\mathcal{O}}_\Lambda \rightarrow \widehat{\mathcal{O}}_\Lambda$, which makes it possible to pass to the *restricted category* $\widehat{\mathcal{O}}_\Lambda^{\text{res}}$, which we define next.

Let A_n be the vector space of natural transformations $\Sigma^n \rightarrow \text{Id}$. This gives rise to a graded algebra $A := \bigoplus_{n \in \mathbb{Z}} A_n$. Then the restricted category $\widehat{\mathcal{O}}_\Lambda^{\text{res}}$ is the full subcategory of $\widehat{\mathcal{O}}_\Lambda$ consisting of all modules which are annihilated by the induced action of A_n for $n \neq 0$; cf. [AF1, §4.3]. The irreducible modules in the restricted category are the same as in $\widehat{\mathcal{O}}_\Lambda$ itself. There are also the *restricted Verma modules*

$$\Delta(\lambda)^{\text{res}} := \Delta(\lambda) \Big/ \sum_{\eta \in A_{\neq 0}} \text{im}(\eta_{\Delta(\lambda)} : \Sigma^\eta \Delta(\lambda) \rightarrow \Delta(\lambda)) \quad (6.9)$$

from [AF1, §4.4]. In other words, $\Delta(\lambda)^{\text{res}}$ is the largest quotient of $\Delta(\lambda)$ that belongs to the restricted category. Similarly, the *restricted dual Verma module* $\nabla(\lambda)^{\text{res}}$ is the largest submodule of $\nabla(\lambda)$ that belongs to the restricted category.

The restricted category $\widehat{\mathcal{O}}_{\Lambda}^{\text{res}}$ is no longer indecomposable: by [AF2, Th. 5.1] it decomposes further as

$$\widehat{\mathcal{O}}_{\Lambda}^{\text{res}} = \prod_{\bar{\Lambda} \in \Lambda/W} \widehat{\mathcal{O}}_{\bar{\Lambda}}^{\text{res}} \quad (6.10)$$

where Λ/W denotes the orbits of W under the dot action. For instance, the poset Λ for the critical level displayed in Example 6.3 splits into two orbits $W \cdot (\alpha_0 - \rho)$ and $W \cdot (\alpha_1 - \rho)$ (i.e., one removes the edges labelled by δ). In the most singular case, $\widehat{\mathcal{O}}_{-\rho}^{\text{res}}$ is a product of simple blocks; in particular, $\Delta^{\text{res}}(-\rho) = L(-\rho) = \nabla^{\text{res}}(-\rho)$.

Conjecture 6.9 (Critical block conjecture). Let $\widehat{\mathcal{O}}_{\bar{\Lambda}}^{\text{res}}$ be a regular integral critical block in the sense of [AF2]. Let $\mathcal{O}_{\bar{\Lambda}}^{\text{res}} := \text{Fin}(\widehat{\mathcal{O}}_{\bar{\Lambda}}^{\text{res}})$ be the full subcategory consisting of all modules of finite length. Then $\mathcal{O}_{\bar{\Lambda}}^{\text{res}}$ is an essentially finite highest weight category with standard and costandard objects $\Delta(\lambda)^{\text{res}}$ and $\nabla(\lambda)^{\text{res}}$ for $\lambda \in \bar{\Lambda}$. Moreover, the indecomposable projective modules in $\mathcal{O}_{\bar{\Lambda}}^{\text{res}}$ are also its indecomposable tilting modules, and therefore $\mathcal{O}_{\bar{\Lambda}}^{\text{res}}$ is tilting-bounded and Ringel self-dual.

This conjecture is true for the basic example of a critical block from Example 6.3 thanks to [Fie3, Th. 6.6]; the same category arises as the principal block of category \mathcal{O} for $\mathfrak{gl}_{|1|}(\mathbb{C})$ discussed in §6.7 below. The conjecture is also consistent with the so-called *Feigin-Frenkel conjecture* [AF1, Conj. 4.7], which says that composition multiplicities of restricted Verma modules are related to the periodic Kazhdan-Lusztig polynomials from [Lus] (and Jantzen's generic decomposition patterns from [Jan2]). These polynomials depend on the relative position of the given pair of weights and, when not too close to walls, they vanish for weights that are far apart. This is consistent with the conjectured existence of indecomposable projectives of finite length in regular blocks of the restricted category.

Remark 6.10. It seems to us that the Feigin-Frenkel conjecture might have an explanation in terms of a sequence of equivalences of categories similar to [FG, (7)]. Ultimately this should connect $\mathcal{O}_{\bar{\Lambda}}^{\text{res}}$ with representations of the quantum group analog of Jantzen's thickened Frobenius kernel G_1T . Assuming that ℓ (the order of the root of unity) is odd and bigger than or equal to the Coxeter number, the latter are known by [AJS, §17] to be essentially finite highest weight categories controlled by the periodic Kazhdan-Lusztig polynomials. Also, in these categories, tilting modules are projective, hence, the Ringel self-duality would be an obvious consequence.

6.4. Rational representations. As we noted in Remark 3.62, the definition of lower finite highest weight category originated in the work of Cline, Parshall and Scott [CPS1]. As well as the BGG category \mathcal{O} already mentioned, their work was motivated by the representation theory of a reductive algebraic group G in positive characteristic, as developed for example in [Jan1]: the symmetric tensor¹³ category $\mathcal{R}ep(G)$ of finite-dimensional rational representations of G is a lower finite highest weight category. Tilting modules for G were studied in [Don3], although our formulation of semi-infinite Ringel duality from §4.4 is not mentioned explicitly there: Donkin instead took the approach pioneered in [Don2] of truncating to a finite lower set before taking Ringel duals. In fact, now, there is monoidal structure in play and the story is even richer.

To give more details, we fix a maximal torus T contained in an opposite pair of Borel subgroups B^+ and B^- of G . Then the weight poset Λ is the set $X^+(T)$ of dominant characters of T with respect to B^+ . We denote the natural duality on $\mathcal{R}ep(G)$ by $V \mapsto V^*$ (with action defined via $g \mapsto g^{-1}$). The costandard objects are the induced modules $H^0(\lambda) := H^0(G/B^-, \mathcal{L}_{\lambda})$ and the standard objects are the Weyl modules $V(\lambda) := H^0(G/B^+, \mathcal{L}_{\lambda}^*)^*$. For the partial order \leq , one can use the usual dominance ordering on $X^+(T)$, or the more refined Bruhat order of [Jan1, §II.6.4]. This makes

¹³Locally finite Abelian, monoidal, rigid, $\text{End}(\mathbf{1}) = \mathbb{k}$.

$\mathcal{R}ep(G)$ into a lower finite highest weight category by [Jan1, Prop. II.4.18] and [Jan1, Prop. II.6.13]. In fact, in the case of $\mathcal{R}ep(G)$, all of the general results about ascending ∇ -flags found in §3.5 were known already before the time of [CPS1], e.g., they are discussed in Donkin’s book [Don1] (and called there *good filtrations*).

Let $\mathcal{T}ilt(G)$ be the full subcategory of $\mathcal{R}ep(G)$ consisting of all tilting modules. A key theorem in this setting is that tensor products of tilting modules are tilting; this is the Donkin-Mathieu-Wang theorem [Don1], [Mat], [Wan]. Thus, $\mathcal{T}ilt(G)$ is a symmetric pseudo-tensor¹⁴ category. Let $(T_i)_{i \in I}$ be a monoidal generator for $\mathcal{T}ilt(G)$, i.e., each T_i is a tilting module and every indecomposable tilting module is isomorphic to a summand of a tensor product $T_{\mathbf{i}} := T_{i_1} \otimes \cdots \otimes T_{i_n}$ for some $n \geq 0$ and $\mathbf{i} = (i_1, \dots, i_n) \in I^n$. Then define \mathcal{A} to be the category with objects $\mathbf{I} := \bigsqcup_{n \geq 0} I^n$ and morphisms defined from $\text{Hom}_{\mathcal{A}}(\mathbf{j}, \mathbf{i}) := \text{Hom}_G(T_{\mathbf{j}}, T_{\mathbf{i}})$, composition being induced by the opposite of composition in $\mathcal{R}ep(G)$. The category \mathcal{A} is naturally a strict symmetric monoidal category, with the tensor product of objects being by concatenation of sequences. The evident monoidal functor $\mathcal{A} \rightarrow \mathcal{T}ilt(G)^{\text{op}}$ extends to the Karoubi envelope of \mathcal{A} , and the resulting functor $\text{Kar}(\mathcal{A}) \rightarrow \mathcal{T}ilt(G)^{\text{op}}$ is a symmetric monoidal equivalence.

Forgetting the monoidal structure, one can think instead in terms of the locally finite-dimensional locally unital algebra $A = \bigoplus_{i,j \in I} e_i A e_j$ that is the path algebra of \mathcal{A} in the sense of Remark 2.3. It becomes convenient to identify $T = \bigoplus_{i \in I} T_i$ and $T^{\otimes} = \bigoplus_{i \in I} T_i^*$ with the tensor algebras

$$T = T(V), \quad T^{\otimes} = T(V^*) \quad \text{where} \quad V := \bigoplus_{i \in I} T_i. \quad (6.11)$$

Note that T is naturally a right A -module and T^{\otimes} is a left A -module. Since T is a tilting generator for $\mathcal{R}ep(G)$ in the sense of Definition 4.24, $A\text{-mod}_{\text{fd}}$ is the Ringel dual of $\mathcal{R}ep(G)$ with respect to T . Theorem 4.25 implies that $A\text{-mod}_{\text{fd}}$ is an upper finite highest weight category with poset $(X^+(T), \geq)$. Moreover, by Corollary 4.29, T^{\otimes} is a tilting generator for $A\text{-mod}_{\text{fd}}$ with $\text{Coend}_A(T^{\otimes}) \cong \mathbb{k}[G]$ as coalgebras.

At this point, the monoidal structure on the category \mathcal{A} comes back into the picture since the A -module T comes from a faithful symmetric monoidal functor (“fiber functor”) $T : \mathcal{A} \rightarrow (\text{Vec}_{\text{fd}})^{\text{op}}$. Consequently, by classical arguments of Tannaka duality (e.g., see [DM, §2] and [EGNO, §5.4]), $\text{Coend}_A(T^{\otimes})$ can be endowed with the structure of a commutative Hopf algebra which reconstructs the coordinate algebra of G . To explain this in more detail, we use the setup of (2.13), so now we are identifying the coalgebra $\text{Coend}_A(T^{\otimes})$ with

$$C := T \otimes_A T^{\otimes} = T(V) \otimes_A T(V^*). \quad (6.12)$$

Then the algebra structure on C is induced by the natural multiplication on the tensor product of algebras $T(V) \otimes T(V^*)$, that is,

$$(v \otimes u) \cdot (v' \otimes u') := (v \otimes v') \otimes (u \otimes u') \quad (6.13)$$

for $v, v' \in T(V)$ and $u, u' \in T(V^*)$. If we pick a basis $v_1^{(i)}, \dots, v_{d(i)}^{(i)}$ for each T_i and let $u_1^{(i)}, \dots, u_{d(i)}^{(i)}$ be the dual basis for T_i^* , then the elements

$$\{c_{r,s}^{(i)} := v_s^{(i)} \otimes u_r^{(i)} \mid i \in I, 1 \leq r, s \leq d(i)\} \quad (6.14)$$

generate C as an algebra. The coalgebra structure satisfies

$$\delta(c_{r,s}^{(i)}) = \sum_{t=1}^{d(i)} c_{r,s}^{(i)} \otimes c_{t,s}^{(i)}, \quad \varepsilon(c_{r,s}^{(i)}) = \delta_{r,s}. \quad (6.15)$$

Now the reconstruction theorem can be formulated as follows.

¹⁴Additive Karoubian, monoidal, rigid, $\text{End}(\mathbf{1}) = \mathbb{k}$.

Theorem 6.11 (Tannakian reconstruction). *The above construction makes the coalgebra $C = \text{Coend}_A(T^{\otimes})$ into a commutative Hopf algebra which is isomorphic to the coordinate algebra $\mathbb{k}[G]$ via the unique algebra homomorphism sending $c_{r,s}^{(i)} \in C$ to the matrix coefficient function $\tilde{c}_{r,s}^{(i)} \in \mathbb{k}[G]$ defined by $gv_s^{(i)} = \sum_{r=1}^{d(i)} \tilde{c}_{r,s}^{(i)}(g)v_r^{(i)}$ for $g \in G$.*

Proof. For $\mathbf{i} = (i_1, \dots, i_n) \in \mathbf{I}^n$ and $\mathbf{r} = (r_1, \dots, r_n), \mathbf{s} = (s_1, \dots, s_n) \in \mathbb{Z}^n$ with $1 \leq r_k, s_k \leq d(i_k)$ for each k , let $c_{\mathbf{r},\mathbf{s}}^{(\mathbf{i})} := (v_{r_1}^{(i_1)} \otimes \dots \otimes v_{r_n}^{(i_n)}) \otimes (u_{s_1}^{(i_1)} \otimes \dots \otimes u_{s_n}^{(i_n)}) \in C$. These are the elements in the formula (4.17), and they span C . The coalgebra isomorphism $C \xrightarrow{\sim} \mathbb{k}[G]$ from Corollary 4.29(i) sends $c_{\mathbf{r},\mathbf{s}}^{(\mathbf{i})} \in C$ to $\tilde{c}_{r_1,s_1}^{(i_1)} \dots \tilde{c}_{r_n,s_n}^{(i_n)} \in \mathbb{k}[G]$. So to be an algebra isomorphism, we must have that $c_{\mathbf{r},\mathbf{s}}^{(\mathbf{i})} = c_{r_1,s_1}^{(i_1)} \dots c_{r_n,s_n}^{(i_n)}$, which is exactly the definition of multiplication given above. \square

Theorem 6.11 recovers a classical result: it is a special case of [DM, Th. 2.11], which implies that $\mathbb{k}[G]$ is isomorphic to $\text{Coend}(F)$ where $F : \mathcal{R}ep(G) \rightarrow \mathcal{V}ec_{\text{fd}}$ is the forgetful functor. To deduce Theorem 6.11 from this statement, one also needs to observe that $\text{Coend}(F) \cong \text{Coend}_A(T)$; this holds because the algebraic group G is isomorphic to its image in its representation on $V = \bigoplus_{i \in I} T_i$ by weight considerations.

Remark 6.12. To get a full set of relations between the generators (6.14) of C , one just needs to take the equations $vx \otimes u = v \otimes xu$ for $x : \mathbf{i} \rightarrow \mathbf{j}$ running over a system of monoidal generators for \mathcal{A} and all $v \in T_{\mathbf{i}}, u \in T_{\mathbf{j}}^*$.

Remark 6.13. Theorem 5.10 can often be applied in this context to give A (or some idempotent expansion of A) the structure of an upper finite (perhaps symmetrically) based quasi-hereditary algebra.

The first example comes from $G = SL_2$. For this, we may take $I := \{|\}$ and $T_{|}$ to be the natural two-dimensional representation V of G with its standard basis v_1, v_2 ; we also use u_1, u_2 to denote the dual basis of V^* . The module V is a monoidal generator for $\mathcal{T}ilt(G)$ by weight considerations. Note also that V possesses an invariant symplectic form such that $(v_1, v_2) = 1$, hence, $V \cong V^*$. The object set $\mathbf{I} = \{|\otimes^n \mid n \in \mathbb{N}\}$ in the above setup may be identified with \mathbb{N} . Hence, $T = \bigoplus_{n \geq 0} T_n$ is the tensor algebra $T(V) = \bigoplus_{n \geq 0} T^n(V)$ and T^{\otimes} is $T(V^*)$. As is well known, the monoidal category \mathcal{A} in this case is the *Temperley-Lieb category* $\mathcal{T}L(-2)$; see e.g. [GW]. It is easy to verify that

$$C = T(V) \otimes_A T(V^*) \cong \mathbb{k}[c_{1,1}, c_{1,2}, c_{2,1}, c_{2,2}]/(\det - 1)$$

where $c_{r,s} = v_s \otimes u_r$ as above and $\det = c_{1,1}c_{2,2} - c_{2,1}c_{1,2}$. Of course this is $\mathbb{k}[SL_2]$.

This example becomes more interesting if we replace the Temperley-Lieb category $\mathcal{T}L(-2)$ with its q -analog $\mathcal{T}L(-q - q^{-1})$ for $q \in \mathbb{k}^\times$. Recall that this is generated as a strictly pivotal monoidal category by one object $|$ and two morphisms $\cup : 0 \rightarrow 2$ and $\cap : 2 \rightarrow 0$ subject to $\bigcirc = -q - q^{-1}$. It is braided with braiding defined by

$$\begin{array}{c} \diagdown \\ \diagup \end{array} := \begin{array}{c} \cup \\ \cap \end{array} + q \left| \right|, \quad \begin{array}{c} \diagdown \\ \diagup \end{array} = \begin{array}{c} \cup \\ \cap \end{array} + q^{-1} \left| \right|. \quad (6.16)$$

As mentioned in Remark 5.4, the natural diagram basis makes the path algebra A of $\mathcal{A} := \mathcal{T}L(-q - q^{-1})$ into an upper finite based quasi-hereditary algebra with weight poset (\mathbb{N}, \geq) . Hence, $A\text{-mod}_{\text{fd}}$ is an upper finite highest weight category.

Next let V be a two-dimensional vector space with basis v_1, v_2 and $(\cdot, \cdot) : V \times V \rightarrow \mathbb{k}$ be the bilinear form with $(v_1, v_2) = 1, (v_2, v_1) = -q^{-1}$ and $(v_1, v_1) = (v_2, v_2) = 0$. A relation check shows that there is a monoidal functor $T : \mathcal{A} \rightarrow (\mathcal{V}ec_{\text{fd}})^{\text{op}}$ such that $T(|) = V$ and

$$T(\cup) : V \otimes V \rightarrow \mathbb{k}, \quad v_i \otimes v_j \mapsto (v_i, v_j), \quad (6.17)$$

$$T(\cap) : \mathbb{k} \rightarrow V \otimes V, \quad 1 \mapsto v_2 \otimes v_1 - qv_1 \otimes v_2. \quad (6.18)$$

Equivalently, the tensor algebra $T = T(V)$ is a right A -module, and its dual $T^\otimes = T(V^*)$ is a left A -module. Then we define C as in (6.12). The coend construction makes C into a cobraided Hopf algebra, hence, $\text{comod}_{\text{fd}}\text{-}C$ is a braided tensor category. Now one can check directly using the homological criterion for ∇ -flags from Theorem 3.39 that T^\otimes is a tilting generator for $A\text{-mod}_{\text{fd}}$. Hence, $\text{comod}_{\text{fd}}\text{-}C$ is the Ringel dual of the upper finite highest weight category $A\text{-mod}_{\text{fd}}$, so it is a lower finite highest weight category thanks to Theorem 4.27.

Let us compute C explicitly. Let u_1, u_2 be the basis for V^* dual to v_1, v_2 . Then C is generated by $\{c_{r,s} := v_s \otimes u_r \mid r, s = 1, 2\}$, and the comultiplication and counit are defined by $\delta(c_{r,s}) = c_{r,1} \otimes c_{1,s} + c_{r,2} \otimes c_{2,s}$, $\varepsilon(c_{r,s}) = \delta_{r,s}$. By Remark 6.12, the following equations give a full set of relations for the algebra C :

$$(v_i \otimes v_j) \otimes (\cup 1) = (v_i \otimes v_j \cup) \otimes 1, \\ (1 \cap) \otimes (u_i \otimes u_j) = 1 \otimes (\cap u_i \otimes u_j).$$

To expand these, note that the left A -module $T^\otimes = T(V^*)$ comes from the monoidal functor $T^\otimes : \mathcal{A} \rightarrow \mathcal{V}ec_{\text{fd}}$ defined by $T^\otimes(\cup) = V^*$ and

$$T^\otimes(\cap) : V^* \otimes V^* \rightarrow \mathbb{k}, \quad u_i \otimes u_j \mapsto (v_j, v_i)^{-1}, \tag{6.19}$$

$$T^\otimes(\cup) : \mathbb{k} \rightarrow V^* \otimes V^*, \quad 1 \mapsto u_1 \otimes u_2 - q^{-1}u_2 \otimes u_1. \tag{6.20}$$

Using this, the relations become $c_{1,i}c_{2,j} - q^{-1}c_{2,i}c_{1,j} = (v_i, v_j)$ and $c_{i,2}c_{j,1} - qc_{i,1}c_{j,2} = (v_j, v_i)^{-1}$, hence, we get

$$\begin{pmatrix} c_{2,2} & -qc_{1,2} \\ -q^{-1}c_{2,1} & c_{1,1} \end{pmatrix} \begin{pmatrix} c_{1,1} & c_{1,2} \\ c_{2,1} & c_{2,2} \end{pmatrix} = \begin{pmatrix} c_{1,1} & c_{1,2} \\ c_{2,1} & c_{2,2} \end{pmatrix} \begin{pmatrix} c_{2,2} & -qc_{1,2} \\ -q^{-1}c_{2,1} & c_{1,1} \end{pmatrix} = I_2.$$

So C is generated by $c_{1,1}, c_{1,2}, c_{2,1}, c_{2,2}$ subject to the relations needed to ensure that

$$\begin{pmatrix} c_{1,1} & c_{1,2} \\ c_{2,1} & c_{2,2} \end{pmatrix}^{-1} = \begin{pmatrix} c_{2,2} & -qc_{1,2} \\ -q^{-1}c_{2,1} & c_{1,1} \end{pmatrix}. \tag{6.21}$$

Equivalently, C is generated by $c_{1,1}, c_{1,2}, c_{2,1}, c_{2,2}$ subject to the relations

$$c_{i,2}c_{i,1} = qc_{i,1}c_{i,2}, \quad c_{2,j}c_{1,j} = qc_{1,j}c_{2,j}, \\ c_{1,2}c_{2,1} = c_{2,1}c_{1,2}, \quad c_{2,2}c_{1,1} = c_{1,1}c_{2,2} + (q - q^{-1})c_{1,2}c_{2,1},$$

and $\det_q := c_{1,1}c_{2,2} - q^{-1}c_{1,2}c_{2,1} = 1$. Thus, we have recovered the well-known quantized coordinate algebra $\mathbb{k}_q[SL_2]$, and $\text{comod}_{\text{fd}}\text{-}C$ is the category of *rational representations of quantum SL_2* .

When at a root of unity over the ground field is \mathbb{C} , the indecomposable projectives and injectives in the category of rational representations of quantum SL_2 (or indeed the quantum group corresponding to a reductive group) are all finite-dimensional, i.e., the category is essentially finite Abelian. Tiltings are also finite-dimensional, indeed, the category is tilting-bounded in the sense of Definition 4.20. The structure of the principal block can be worked out explicitly (e.g., see [AT, Th. 3.12, Def. 3.3]): it is Morita equivalent to the locally unital algebra that is the path algebra of the quiver

$$0 \begin{array}{c} \xrightarrow{x_0} \\ \xleftarrow{y_0} \end{array} 1 \begin{array}{c} \xrightarrow{x_1} \\ \xleftarrow{y_1} \end{array} 2 \begin{array}{c} \xrightarrow{x_2} \\ \xleftarrow{y_2} \end{array} 3 \cdots \quad \text{with relations } x_{i+1}x_i = y_iy_{i+1} = x_iy_i - y_{i+1}x_{i+1} = 0.$$

The appropriate partial order on the weight poset \mathbb{N} is the natural order $0 < 1 < \dots$. The indecomposable projectives have the following structure:

$$P(0) = \begin{array}{c} 0 \\ \downarrow \\ 1 \\ \downarrow \\ 0 \end{array}, \quad P(1) = \begin{array}{c} 1 \\ \swarrow \quad \searrow \\ 0 \quad 2 \\ \nwarrow \quad \nearrow \\ \quad 1 \end{array}, \quad P(2) = \begin{array}{c} 2 \\ \swarrow \quad \searrow \\ 1 \quad 3 \\ \nwarrow \quad \nearrow \\ \quad 2 \end{array}, \quad P(3) = \begin{array}{c} 3 \\ \swarrow \quad \searrow \\ 2 \quad 4 \\ \nwarrow \quad \nearrow \\ \quad 3 \end{array}, \dots$$

The tilting objects are $T(0) := L(0)$ and $T(n) := P(n - 1)$ for $n \geq 1$. From this, it is easy to see that the Ringel dual is described by the same quiver with one additional relation, namely, that $y_0x_0 = 0$ (and of course the partial order is reversed).

6.5. Tensor product categorifications. Until quite recently, most of the naturally-occurring examples were highest weight categories (like the ones described in the previous two subsections). But the work of Webster [Web1], [Web2] and Losev and Webster [LW] has brought to prominence a very general source of examples that are fully stratified but seldom highest weight.

Fundamental amongst these new examples are the categorifications of tensor products of irreducible highest weight modules of symmetrizable Kac-Moody Lie algebras. Rather than attempting to repeat the definition of these here, we refer the reader to [LW]. All of these examples are finite fully stratified categories possessing a Chevalley duality. They are also tilting-rigid; the proof of this depends on an argument involving translation/projective functors. Consequently, the Ringel dual is again a finite fully stratified category that is tilting-rigid. In fact, the Ringel dual category is always another tensor product categorification¹⁵ (reverse the order of the tensor product). In the earlier article [Web2], Webster also wrote down explicit finite-dimensional algebras which give realization of these categories. In view of Theorem 5.25, all of Webster's algebras admit bases making them into symmetrically based stratified algebras, although these bases are usually hard to construct explicitly.

In [Web1], Webster also introduced some more general tensor product categorifications, including ones which categorify the tensor product of an integrable lowest weight module tensored with an integrable highest weight module. The latter are particularly important since they may be realized as *generalized cyclotomic quotients* of the Kac-Moody 2-category. They are upper finite fully stratified categories. In type A, they can also be realized as generalized cyclotomic quotients of the (degenerate or quantum) Heisenberg category; see [BSW, Th. B]. In the latter realization, they should possess explicit triangular bases, generalizing the ones for the cyclotomic quotients of central charge zero discussed in [GRS].

6.6. Deligne categories. Another source of upper finite highest weight categories comes from various Deligne categories. The definition of these categories is diagrammatic in nature. For example, in characteristic zero, the Deligne category $\underline{\text{Rep}}(GL_\delta)$ is the Karoubi envelope of the oriented Brauer category $\mathcal{OB}(\delta)$. This case was studied in the PhD thesis of Reynolds [Rey] based on the observation that it admits a symmetric split triangular decomposition; see also [Bru] which treats the HOMFLY-PT skein category at the same time. Rui and Song [RS] have analysed the Brauer category and the Kauffman skein category by similar techniques. Similar ideas have been developed independently by Sam and Snowden [SS], who also consider other types of Deligne category.

The category of locally finite-dimensional representations of the Deligne category $\underline{\text{Rep}}(GL_\delta)$ can also be interpreted as a special case of the lowest weight tensored highest weight tensor product categorifications discussed in the previous subsection; see the introduction of [Bru]. The Ringel dual in this example is equivalent to the Abelian envelope $\underline{\text{Rep}}^{ab}(GL_\delta)$ of Deligne's category constructed by Entova, Hinich and Serganova [EHS], which is a monoidal lower finite highest weight category. In [Ent], it is shown that $\underline{\text{Rep}}^{ab}(GL_\delta)$ categorifies a highest weight tensored lowest weight representation, which is the dual result to the one from [Bru]. This example will be discussed further in the sequel to this article, where we give an explicit description of the blocks of $\underline{\text{Rep}}^{ab}(GL_\delta)$ via Khovanov's arc coalgebra (an interesting explicit example of a based quasi-hereditary coalgebra), thereby proving a conjecture formulated in the introduction of [BS2].

¹⁵This was noted in Remark 3.10 of the arXiv version of [LW] but the authors removed this remark in the published version.

These and the other classical families of Deligne categories $\text{Rep}(O_\delta)$, $\text{Rep}(P)$ and $\text{Rep}(Q)$ are being investigated actively along similar lines by several groups of authors and there has been considerable recent progress; e.g., see [Cou4], [SS]. There are also many interesting connections here with rational representations of the corresponding families of classical supergroups.

6.7. Representations of Lie superalgebras. Finally, we mention briefly an interesting source of essentially finite highest weight categories: the analogs of the BGG category \mathcal{O} for classical Lie superalgebras. A detailed account in the case of the Lie superalgebra $\mathfrak{gl}_{m|n}(\mathbb{C})$ can be found in [BLW]. Its category \mathcal{O} gives an essentially finite highest weight category which is neither lower finite nor upper finite. Moreover, it is tilting-bounded as in Definition 4.20, so that the Ringel dual category is also an essentially finite highest weight category.

There is one very easy special case: the principal block of category \mathcal{O} for $\mathfrak{gl}_{1|1}(\mathbb{C})$ is equivalent to the category of finite-dimensional modules over the essentially finite-dimensional locally unital algebra which is the path algebra of the following quiver:

$$\cdots -1 \begin{array}{c} \xrightarrow{x_{-1}} \\ \xleftarrow{y_{-1}} \end{array} 0 \begin{array}{c} \xrightarrow{x_0} \\ \xleftarrow{y_0} \end{array} 1 \begin{array}{c} \xrightarrow{x_1} \\ \xleftarrow{y_1} \end{array} 2 \cdots \text{ with relations } x_{i+1}x_i = y_i y_{i+1} = x_i y_i - y_{i+1} x_{i+1} = 0,$$

see e.g. [BS1, p. 380]. This is very similar to the $U_q(\mathfrak{sl}_2)$ -example from §6.4, but now the poset \mathbb{Z} (ordered naturally) is neither lower nor upper finite. From the category \mathcal{O} perspective, this example is rather misleading since its projective, injective and tilting objects coincide, hence, it is Ringel self-dual.

One gets similar examples from $\mathfrak{osp}_{m|2n}(\mathbb{C})$, as discussed for example in [BW] and [ES]. The simplest non-trivial case of $\mathfrak{osp}_{3|2}(\mathbb{C})$ produces the path algebra of a D_∞ quiver (replacing than the A_∞ quiver above); see [ES, §II]. The “strange” families $\mathfrak{p}_n(\mathbb{C})$ and $\mathfrak{q}_n(\mathbb{C})$ also exhibit similar structures. The former has not yet been investigated systematically (although basic aspects of the finite-dimensional finite-dimensional representations and category \mathcal{O} were recently studied in [B+9] and [CC], respectively). It is an interesting example of a naturally-occurring highest weight category which does not admit a Chevalley duality. For $\mathfrak{q}_n(\mathbb{C})$, we refer to [BD2] and the references therein. In fact, the integral blocks for $\mathfrak{q}_n(\mathbb{C})$ are fibered highest weight categories; this observation is due to Frisk [Fri2].

REFERENCES

- [ADL] I. Ágoston, V. Dlab and E. Lukács, Stratified algebras, *Math. Rep. Acad. Sci. Canada* **20** (1998), 22–28.
- [AHLU] I. Ágoston, D. Happel, E. Lukács and L. Unger, Standardly stratified algebras and tilting, *J. Algebra* **226** (2000), 144–160.
- [And] H. H. Andersen, Tilting modules and cellular categories, *J. Pure Appl. Algebra*, **224** (2020), 106366, 29 pp..
- [AJS] H. H. Andersen, J. C. Jantzen and W. Soergel, Representations of quantum groups at a pth root of unity and of semisimple groups in characteristic p , *Astérisque* **220** (1994).
- [AST] H. H. Andersen, C. Stroppel and D. Tubbenhauer, Cellular structures using U_q -tilting modules, *Pacific J. Math.* **292** (2018), 21–59.
- [AT] H. H. Andersen and D. Tubbenhauer, Diagram categories for U_q -tilting modules at roots of unity, *Transform. Groups* **22** (2017), 29–89.
- [AF1] T. Arakawa and P. Fiebig, On the restricted Verma modules at the critical level, *Trans. Amer. Math. Soc.* **364** (2012), 4683–4712.
- [AF2] ———, The linkage principle for restricted critical level representations of affine Kac-Moody algebras, *Compositio Math.* **148** (2012), 1787–1810.
- [Ark] S. Arkhipov, Semi-infinite cohomology of associative algebras and bar duality, *Int. Math. Res. Notices* 1997, 833–863.
- [B+9] M. Balagovic, Z. Daugherty, I. Entova-Aizenbud, I. Halacheva, J. Hennig, M. S. Im, G. Letzter, E. Norton, V. Serganova and C. Stroppel, Translation functors and decomposition numbers for the periplectic Lie superalgebra $\mathfrak{p}(n)$, *Math. Res. Lett.* **26** (2019), 643–710.

- [BW] H. Bao and W. Wang, A new approach to Kazhdan-Lusztig theory of type B via quantum symmetric pairs, *Astérisque* 2018, no. 402, vii+134 pp..
- [BBD] A. Beilinson, J. Bernstein and P. Deligne, Faisceaux pervers, *Astérisque* **100** (1982), 5–171.
- [BR] A. Beligiannis and I. Reiten, Homological and homotopical aspects of torsion theories, *Mem. Amer. Math. Soc.* **188** (2007), no. 883, viii+207 pp..
- [BT] G. Bellamy and U. Thiel, Highest weight theory for finite-dimensional graded algebras with triangular decomposition, *Advances Math.* **330** (2018), 361–419.
- [BY] R. Bezrukavnikov and Z. Yun, On Koszul duality for Kac-Moody groups, *Represent. Theory* **17** (2013), 1–98.
- [Bru] J. Brundan, Representations of oriented skein categories; [arXiv:1712.08953](https://arxiv.org/abs/1712.08953).
- [BD1] J. Brundan and N. Davidson, Categorical actions and crystals, *Contemp. Math.* **684** (2017), 116–159.
- [BD2] ———, Type C blocks of super category \mathcal{O} , *Math. Z.* **293** (2019), 867–901.
- [BLW] J. Brundan, I. Losev and B. Webster, Tensor product categorifications and the super Kazhdan-Lusztig conjecture, *Int. Math. Res. Notices* 2017, 6329–6410.
- [BSW] J. Brundan, A. Savage and B. Webster, Heisenberg and Kac-Moody categorification, *Selecta Math.* **26** (2020), 74.
- [BS1] J. Brundan and C. Stroppel, Highest weight categories arising from Khovanov’s diagram algebra IV: the general linear supergroup, *J. Eur. Math. Soc.* **14** (2012), 373–419.
- [BS2] ———, Gradings on walled Brauer algebras and Khovanov’s arc algebra, *Advances Math.* **231** (2012), 709–773.
- [CaeZ] B. Zhu and S. Caenepeel, On good filtration dimensions for standardly stratified algebras, *Comm. Algebra* **32** (2004), 1603–1614.
- [Car] R. W. Carter, *Lie Algebras of Finite and Affine Type*, Cambridge Studies in Advanced Mathematics **96**, Cambridge University Press, 2005.
- [Che] X. Chen, Gorenstein Homological Algebra of Artin Algebras; [arXiv:1712.04587](https://arxiv.org/abs/1712.04587).
- [CC] C.-W. Chen and K. Coulembier, The primitive spectrum and category \mathcal{O} for the periplectic Lie superalgebra, *Canad. J. Math.* **72** (2020), 625–655.
- [CPS1] E. Cline, B. Parshall and L. Scott, Finite dimensional algebras and highest weight categories, *J. Reine Angew. Math.* **391** (1988), 85–99.
- [CPS2] ———, Stratifying endomorphism algebras, *Mem. Amer. Math. Soc.* **124** (1996), 1–119.
- [Cou1] K. Coulembier, Tensor ideals, Deligne categories and invariant theory, *Selecta Math.* **24** (2018), 4659–4710.
- [Cou2] ———, The classification of blocks in BGG category \mathcal{O} , *Math. Z.* **295** (2020), 821–837.
- [Cou3] ———, Some homological properties of ind-completions and highest weight categories, *J. Algebra* **562** (2020), 341–367.
- [Cou4] ———, Monoidal abelian envelopes; [arXiv:2003.10105](https://arxiv.org/abs/2003.10105).
- [CouZ] K. Coulembier and R.-B. Zhang, Borelic pairs for stratified algebras, *Advances Math.* **345** (2019), 53–115.
- [CM] T. Cruz and R. Marczinzik, On properly stratified Gorenstein algebras; [arXiv:2101.11899](https://arxiv.org/abs/2101.11899).
- [DM] P. Deligne and J. Milne, Tannakian categories, in: “Hodge cycles, motives, and Shimura varieties”, Lecture Notes in Mathematics, 900, Springer-Verlag, 1982.
- [DGK] V. Deodhar, O. Gabber and V. Kac, Structure of some categories of representations of infinite-dimensional Lie algebras, *Advances Math.* **45** (1982), 92–116.
- [Dla1] V. Dlab, Quasi-hereditary algebras revisited, *An. Șt. Univ. Ovidius Constanța* **4** (1996), 43–54.
- [Dla2] ———, Properly stratified algebras, *C. R. Acad. Sci. Paris* **331** (2000), 191–196.
- [DR] V. Dlab and C. Ringel, The module theoretical approach to quasi-hereditary algebras, in: “Representations of Algebras and Related Topics (1990)”, pp. 200–224, London Math. Soc. Lecture Note Ser. **168**, Cambridge University Press, 1992.
- [Don1] S. Donkin, *Rational Representations of Algebraic Groups*, LNM **1140**, Springer, 1985.
- [Don2] ———, On Schur algebras and related algebras I, *J. Algebra* **104** (1986), 310–328.
- [Don3] ———, On tilting modules for algebraic groups, *Math. Z.* **212** (1993), 39–60.
- [Don4] ———, *The q-Schur Algebra*, London Math. Soc., 1998.
- [DuR] J. Du and H. Rui, Based algebras and standard bases for quasi-hereditary algebras, *Trans. Amer. Math. Soc.* **350** (1998), 3207–3235.
- [ES] M. Ehrig and C. Stroppel, On the category of finite-dimensional representations of $OSp(r|2n)$: part I, in: “Representation Theory—Current Trends and Perspectives”, 109–170, *EMS Ser. Congr. Rep.*, Eur. Math. Soc., Zurich, 2017.
- [ELau] B. Elias and A. Lauda, Trace decategorification of the Hecke category, *J. Algebra* **449** (2016), 615–634.
- [ELos] B. Elias and I. Losev, Modular representation theory in type A via Soergel bimodules; [arXiv:1701.00560](https://arxiv.org/abs/1701.00560).
- [EW] B. Elias and G. Williamson, Soergel calculus, *Represent. Theory* **20** (2016), 295–374.

- [Ent] I. Entova-Aizenbud, Categorical actions and multiplicities in the Deligne category $\text{Rep}(GL_t)$ *J. Algebra* **504** (2018), 391–431.
- [EHS] I. Entova-Aizenbud, V. Hinich and V. Serganova, Deligne categories and the limit of categories $\text{Rep}(GL(m|n))$, *Int. Math. Res. Not.* **2020**, 4602–4666.
- [EP] K. Erdmann and A. Parker, On the global and ∇ -filtration dimensions of quasi-hereditary algebras, *J. Pure Appl. Algebra* **194** (2004), 95–111.
- [EGNO] P. Etingof, S. Gelaki, D. Nikshych and V. Ostrik, *Tensor Categories*, Amer. Math. Soc., 2015.
- [Fie1] P. Fiebig, Centers and translation functors for the category \mathcal{O} over Kac-Moody algebras, *Math. Z.* **243** (2003), 689–717.
- [Fie2] ———, The combinatorics of category \mathcal{O} over symmetrizable Kac-Moody algebras, *Transform. Groups* **11** (2006), 29–49.
- [Fie3] ———, On the subgeneric restricted blocks of affine category \mathcal{O} at the critical level, in: “Symmetries, Integrable Systems and Representations”, pp. 65–84, *Springer Proc. Math. Stat.* **40**, 2013.
- [FG] E. Frenkel and D. Gaitsgory, Localization of \mathfrak{g} -modules on the affine Grassmannian, *Ann. Math.* **170** (2009), 1339–1381.
- [FKS] I. Frenkel, M. Khovanov and C. Stroppel, A categorification of finite-dimensional irreducible representations of quantum \mathfrak{sl}_2 and their tensor products, *Selecta Math.* **12** (2006), 379–431.
- [Fre] P. Freyd, Abelian categories, *Reprints in Theory and Applications of Categories* **3** (2003), 23–164.
- [Fri1] A. Frisk, Dlab’s theorem and tilting modules for stratified algebras, *J. Algebra* **314** (2007) 507–537.
- [FM] A. Frisk and V. Mazorchuk, Properly stratified algebras and tilting, *Proc. London Math. Soc.* **92** (2006), 29–61.
- [Fri2] ———, Typical blocks of the category \mathcal{O} for the queer Lie superalgebra, *J. Algebra Appl.* **6** (2007), 731–778.
- [Gab] P. Gabriel, Des catégories Abéliennes, *Bull. Soc. Math. France* **90** (1962), 323–448.
- [GRS] M. Gao, H. Rui and L. Song, Representations of weakly triangular categories; [arXiv:2012.02945](https://arxiv.org/abs/2012.02945).
- [GHK] N. Gubareni, M. Hazewinkel and V. Kirichenko, *Algebras, Rings and Modules*, vol. 2, Mathematics and Its Applications (Springer), **586**, Springer, Dordrecht, 2007.
- [GW] F. Goodman and H. Wenzl, The Temperley-Lieb algebra at roots of unity, *Pacific J. Math.* **161** (1993), 307–334.
- [GL] J. Graham and G. Lehrer, Cellular algebras, *Invent. Math.* **123** (1996), 1–34.
- [Gre] J. A. Green, Combinatorics and the Schur algebra, *J. Pure Appl. Algebra* **88** (1993), 89–106.
- [Hap] D. Happel, *Triangulated Categories in the Representation Theory of Finite Dimensional Algebras*, London Math. Soc., 1988.
- [HU] D. Happel and L Unger, Modules of finite projective dimension and cocovers, *Math. Ann.* **306** (1996), 445–457.
- [Jan1] J. C. Jantzen, *Representations of Algebraic Groups*, Academic Press, Orlando, 1987.
- [Jan2] ———, Über das Dekompositionsverhalten gewisser modularer Darstellungen halbeinfacher Gruppen und ihrer Lie-Algebren, *J. Algebra* **49** (1977), 441–469.
- [HN] R. Holmes and D. Nakano, Brauer-type reciprocity for a class of graded associative algebras, *J. Algebra* **144** (1991), 117–126.
- [Kac] V. Kac, *Infinite-dimensional Lie Algebras*, Cambridge University Press, 1985.
- [KK] V. Kac and D. Kazhdan, Structure of representations with highest weight of infinite dimensional Lie algebras, *Advances Math.* **34** (1979), 97–108.
- [KS] M. Kashiwara and P. Shapira, *Sheaves and Categories*, Springer, 2000.
- [Kel] B. Keller, Derived categories and tilting, in: “Handbook of Tilting Theory”, pp. 49–104, London Math. Soc. Lecture Notes **332**, Cambridge University Press, 2007.
- [Kle] A. Kleshchev, Affine highest weight categories and affine quasihereditary algebras, *Proc. London Math. Soc.*, **110**, (2015), 841–882.
- [KM] A. Kleshchev and R. Muth, Based quasi-hereditary algebras, *J. Algebra* **558** (2020), 504–522.
- [KX] S. König, and C. Xi, Affine cellular algebras, *Adv. Math.* **229** (2012), 139–182.
- [Kum] S. Kumar, *Kac-Moody Groups, their Flag Varieties and Representation Theory*, Progress in Mathematics **204**, Birkhäuser Boston, 2002.
- [LW] I. Losev and B. Webster, On uniqueness of tensor products of irreducible categorifications, *Selecta Math.* **21** (2015), 345–377.
- [Lus] G. Lusztig, Hecke algebras and Jantzen’s generic decomposition patterns, *Advances Math.* **37** (1980), 121–164.
- [Mac] S. Mac Lane, *Categories for the Working Mathematician*, Springer, 1978.
- [Mad] D. Madsen, Quasi-hereditary algebras and the category of modules with standard filtration, *São Paulo J. Math. Sci.* **11** (2017), 68–80.

- [MZ] F. Marko and A. Zubkov, Pseudocompact algebras and highest weight categories, *Algebr. Represent. Theor.* **16** (2013), 689–728.
- [Mat] O. Mathieu, Filtrations of G -modules, *Ann. Sci. ENS* **23** (1990), 625–644.
- [Maz1] V. Mazorchuk, Stratified algebras arising in Lie theory, in: “Representations of Finite Dimensional Algebras and Related Topics in Lie Theory and Geometry”, pp. 245–260, *Fields Inst. Commun.* **40**, Amer. Math. Soc., Providence, RI, 2004.
- [Maz2] ———, Koszul duality for stratified algebras II, *J. Aust. Math. Soc.* **89** (2010), 23–49.
- [MO] V. Mazorchuk and S. Ovsienko, Finitistic dimension of properly stratified algebras, *Advances Math.* **186** (2004), 251–265.
- [MP] V. Mazorchuk and A. Parker, On the relation between finitistic and good filtration dimensions, *Comm. Algebra* **32** (2004), 1903–1916.
- [MS] V. Mazorchuk and C. Stroppel, Categorification of (induced) cell modules and the rough structure of generalised Verma modules, *Advances Math.* **219** (2008), 1363–1426.
- [Mit] B. Mitchell, Rings with several objects, *Advances Math.* **8** (1972), 1–161.
- [Nav] G. Navarro, Simple comodules and localization in coalgebras, in: “New techniques in Hopf algebras and graded ring theory”, pp. 141–164, KVAB, Brussels, 2007.
- [Nei] W. Neidhardt, Translation to and fro over Kac-Moody algebras, *Pacific J. Math.* **139** (1989), 107–153.
- [PS] L. Positselski and J. Štoviček, ∞ -tilting theory, *Pacific J. Math.* **301** (2019), 297–334.
- [Rei] I. Reiten, Tilting theory and homologically finite subcategories with applications to quasihereditary algebras, in: “Handbook of Tilting Theory”, pp. 179–214, London Math. Soc. Lecture Notes **332**, Cambridge University Press, 2007.
- [Rey] A. Reynolds, *Representations of Oriented Brauer Categories*, PhD thesis, University of Oregon, 2015.
- [RW] S. Riche and G. Williamson, Tilting modules and the p -canonical basis, *Astérisque* **397** (2018).
- [Ric] J. Rickard, Equivalences of derived categories for symmetric algebras, *J. Algebra* **257** (2002), 460–481.
- [Rin] C. Ringel, The category of modules with good filtrations over a quasi-hereditary algebra has almost split sequences, *Math. Z.* **208** (1991), 209–225.
- [RS] H. Rui and L. Song, Representations of Brauer category and categorification, *J. Algebra* **557** (2020), 1–36.
- [SS] S. Sam and A. Snowden, The representation theory of Brauer categories I: triangular categories; [arXiv:2006.04328](https://arxiv.org/abs/2006.04328).
- [SVV] P. Shan, M. Varagnolo, and E. Vasserot, Koszul duality of affine Kac-Moody algebras and cyclotomic rational double affine Hecke algebras, *Advances Math.* **262** (2014), 370–435.
- [Sim] D. Simson, Coalgebras, comodules, pseudocompact algebras and tame comodule type, *Colloq. Math.* **90** (2001), 101–150.
- [Soe] W. Soergel, Character formulas for tilting modules over Kac-Moody algebras, *Represent. Theory* **2** (1998), 432–448.
- [Str] C. Stroppel, Untersuchungen zu den parabolischen Kazhdan-Lusztig-Polynomen für affine Weyl-Gruppen, Diploma thesis, University of Freiburg, 1997.
- [Tak] M. Takeuchi, Morita theorems for categories of comodules, *J. Fac. Sci. Univ. Tokyo* **24** (1977), 629–644.
- [Wak] T. Wakamatsu, On modules with trivial self-extensions, *J. Algebra* **114** (1988), 106–114.
- [Wan] J.-P. Wang, Sheaf cohomology on G/B and tensor products of Weyl modules, *J. Algebra* **77** (1982), 162–185.
- [Web1] B. Webster, Canonical bases and higher representation theory, *Compositio Math.* **151** (2015), 121–166.
- [Web2] ———, Knot invariants and higher representation theory, *Mem. Amer. Math. Soc.* **250** (2017), 1–141.
- [Weil] C. Weibel, *An Introduction to Homological Algebra*, CUP, 1994.
- [Wes] B. Westbury, Invariant tensors and cellular categories, *J. Algebra* **321** (2009), 3563–3567.
- [Xi] C. Xi, Cellular and affine cellular algebras, *Adv. Math (China)* **39**, 257–270.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403, USA
 Email address: brundan@uoregon.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BONN, 53115 BONN, GERMANY
 Email address: stroppel@math.uni-bonn.de