

**CAPACITIES IN WIENER SPACE,
QUASI-SURE LOWER FUNCTIONS,
AND KOLMOGOROV'S ε -ENTROPY**

DAVAR KHOSHNEVISAN, DAVID A. LEVIN, AND PEDRO J. MÉNDEZ-HERNÁNDEZ

ABSTRACT. We propose a set-indexed family of capacities $\{\text{cap}_G\}_{G \subseteq \mathbf{R}_+}$ on the classical Wiener space $C(\mathbf{R}_+)$. This family interpolates between the Wiener measure $(\text{cap}_{\{0\}})$ on $C(\mathbf{R}_+)$ and the standard capacity $(\text{cap}_{\mathbf{R}_+})$ on Wiener space. We then apply our capacities to characterize all quasi-sure lower functions in $C(\mathbf{R}_+)$. In order to do this we derive the following capacity estimate (Theorem 2.3) which may be of independent interest: There exists a constant $a > 1$ such that for all $r > 0$,

$$\frac{1}{a} K_G(r^6) e^{-\pi^2/(8r^2)} \leq \text{cap}_G\{f^* \leq r\} \leq a K_G(r^6) e^{-\pi^2/(8r^2)}.$$

Here, K_G denotes the Kolmogorov ε -entropy of G , and $f^* := \sup_{[0,1]} |f|$.

1. INTRODUCTION

Let $C(\mathbf{R}_+)$ denote the collection of all continuous functions $f : \mathbf{R}_+ \rightarrow \mathbf{R}$. We endow $C(\mathbf{R}_+)$ with its usual topology of uniform convergence on compacts as well as the corresponding Borel σ -algebra \mathcal{B} . In keeping with the literature, elements of \mathcal{B} are called *events*.

Denote by μ the Wiener measure on $(C(\mathbf{R}_+), \mathcal{B})$. Recall that an event Λ is said to hold almost surely [a.s.] if $\mu(\Lambda) = 1$.

Next we define $U := \{U_s\}_{s \geq 0}$ to be the *Ornstein–Uhlenbeck process* on $C(\mathbf{R}_+)$. The process U is characterized by the following requirements:

- (1) It is a stationary infinite-dimensional diffusion with value in $C(\mathbf{R}_+)$;
- (2) Its invariant measure is μ . This implies that for any fixed $s \geq 0$, $\{U_s(t)\}_{t \geq 0}$ is a standard linear Brownian motion.
- (3) For any given $t \geq 0$, $\{U_s(t)\}_{s \geq 0}$ is a standard Ornstein–Uhlenbeck process on \mathbf{R} ; i.e., it satisfies the stochastic differential equation,

$$(1.1) \quad dU_s(t) = -U_s(t) ds + \sqrt{2} dX_s \quad \forall s \geq 0,$$

where X is a Brownian motion.

Following P. Malliavin (1979), we say that an event Λ holds *quasi-surely* [q.s.] if

$$(1.2) \quad \mathbb{P} \{U_s \in \Lambda \text{ for all } s \geq 0\} = 1.$$

Because $t \mapsto U_s(t)$ is a Brownian motion, any event Λ that holds q.s. also holds a.s. The converse is not always true. For example, define Λ_0 to be the collection of all

Date: September 25, 2004.

1991 Mathematics Subject Classification. 60J45, 60J65, 28C20.

Key words and phrases. Capacity in Wiener space, lower functions, Kolmogorov entropy.

The research of D. Kh. is partially supported by a grant from the NSF.

functions $f \in C(\mathbf{R}_+)$ that satisfy $f(1) \neq 0$ (Fukushima, 1984). Evidently, Λ_0 holds a.s. because with probability one Brownian motion at time one is not at the origin. On the other hand, Λ_0 does not hold q.s. because $\{U_s(1)\}_{s \geq 0}$ is point-recurrent. So the chances are 100% that $U_s(1) = 0$ for some $s \geq 0$.

Despite the preceding disclaimer, a number of interesting classical events of full Wiener measure do hold q.s. A notable example is a theorem of M. Fukushima (1984). We can state it, somewhat informally, as follows:

(1.3) The Law of the Iterated Logarithm (LIL) of Khintchine (1933) holds q.s.

It might help to recall Khintchine's theorem: For μ -every $f \in C(\mathbf{R}_+)$,

$$(1.4) \quad \limsup_{t \rightarrow \infty} \frac{f(t)}{\sqrt{2t \ln \ln t}} = 1.$$

Thus we are led to the precise formulation of (1.3): With probability one, the continuous function $f := U_s$ satisfies (1.4), simultaneously for all $s \geq 0$.

For another example consider "the other LIL" which was discovered by K. L. Chung (1948). Chung's LIL states that for μ -almost every $f \in C(\mathbf{R}_+)$,

$$(1.5) \quad \liminf_{t \rightarrow \infty} \frac{\sup_{u \in [0, t]} |f(u)|}{\sqrt{t / \ln \ln t}} = \frac{\pi}{\sqrt{8}}.$$

Fukushima's method can be adapted to prove that

(1.6) Chung's LIL holds q.s.

To be more precise: With probability one, the continuous function $f := U_s$ satisfies (1.5) simultaneously for all $s \geq 0$.

T. S. Mountford (1992) has derived the quasi-sure integral test corresponding to (1.3). One of the initial aims of this article was to complement Mountford's theorem by finding a precise quasi-sure integral test for (1.6). Before presenting this work, let us introduce the notion of "relative capacity."

For all Borel sets $G \subseteq \mathbf{R}_+$ and $\Lambda \in C(\mathbf{R}_+)$ define

$$(1.7) \quad \text{cap}_G(\Lambda) := \int_0^\infty \mathbb{P} \{U_s \in \Lambda \text{ for some } s \in G \cap [0, \sigma]\} e^{-\sigma} d\sigma.$$

We think of $\text{cap}_G(\Lambda)$ as the *capacity of Λ relative to the coordinates in G* . The special case $\text{cap}_{\mathbf{R}_+}$ is well known and well studied (Fukushima, 1984); $\text{cap}_{\mathbf{R}_+}$ is called *the capacity on Wiener space*. According to (1.2), an event Λ holds q.s. iff its complement has zero $\text{cap}_{\mathbf{R}_+}$ -capacity.

The case where $G := \{s\}$ is a singleton is even better studied because of the simple fact that $\text{cap}_{\{s\}}$ is a multiple of the Wiener measure. Thus, $G \mapsto \text{cap}_G(\Lambda)$ interpolates from the Wiener measure ($G = \{0\}$) to the standard capacity on Wiener space ($G = \mathbf{R}_+$). This "interpolation" property was announced in the Abstract.

Now let $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ be decreasing and measurable, and define

$$(1.8) \quad \mathcal{L}(H) := \left\{ f \in C(\mathbf{R}_+) : \liminf_{t \rightarrow \infty} \left[\sup_{u \in [0, t]} |f(u)| - H(t)\sqrt{t} \right] > 0 \right\}.$$

A decreasing measurable function $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ is called an *a.s.-lower function* if $\mathcal{L}(H)$ holds a.s.; i.e., μ -almost every $f \in C(\mathbf{R}_+)$ is in $\mathcal{L}(H)$. Likewise, H is

called a *q.s.-lower function* if $\mathcal{L}(H)$ holds q.s. [The literature actually calls the function $t \mapsto H(t)\sqrt{t}$ an a.s.[q.s]-lower function if $\mathcal{L}(H)$ holds a.s.[q.s.], but we find our parameterization here convenient.]

To understand the utility of these definitions better, consider the special case that $H(t) = \sqrt{c/\ln \ln t}$ for a fixed $c > 0$ ($t \geq 0$). In this case, Chung's LIL (1.5) states that $\mathcal{L}(H)$ holds a.s. if $c < \pi/\sqrt{8}$; its complement holds a.s. if $c > \pi/\sqrt{8}$. In fact, a precise P-a.s. integral test is known (Chung, 1948); see Corollary 1.3 below.

We aim to characterize exactly when $(\mathcal{L}(H))^{\mathbb{G}}$ has positive cap_G -capacity. Define K_G to be the *Kolmogorov ε -entropy* of G (Dudley, 1973; Tihomirov, 1963); i.e., for any $\varepsilon > 0$, $k = K_E(\varepsilon)$ is the maximal number of points $x_1, \dots, x_k \in E$ such that whenever $i \neq j$, $|x_i - x_j| \geq \varepsilon$.

Theorem 1.1. *Choose and fix a decreasing measurable function $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$, and a bounded Borel set $G \subset \mathbf{R}_+$. Then, $\text{cap}_G((\mathcal{L}(H))^{\mathbb{G}}) = 0$ if and only if there exists a decomposition $G = \cup_{n=1}^{\infty} G_n$ in terms of closed sets $\{G_n\}_{n=1}^{\infty}$, such that*

$$(1.9) \quad \int_1^{\infty} \frac{K_{G_n}(H^6(s))}{sH^2(s)} \exp\left(-\frac{\pi^2}{8H^2(s)}\right) ds < \infty \quad \forall n \geq 1.$$

Theorem 1.1 yields the following definite refinement of (1.5).

Corollary 1.2. *Choose and fix a decreasing measurable function $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$. Then, $\mathcal{L}(H)$ holds q.s. if and only if*

$$(1.10) \quad \int_1^{\infty} \exp\left(-\frac{\pi^2}{8H^2(s)}\right) \frac{ds}{sH^8(s)} < \infty.$$

Theorem 1.1 also contains the original almost-sure integral test of Chung (1948). To prove this, simply plug $G = \{u\}$ in Theorem 1.1. Then, $K_{\{u\} \cap J}(\varepsilon)$ is one if $u \in J$ and zero otherwise. Thus we obtain the following.

Corollary 1.3 (Chung (1948)). *Choose and fix a decreasing measurable function $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$. Then $\mathcal{L}(H)$ holds a.s. if and only if*

$$(1.11) \quad \int_1^{\infty} \exp\left(-\frac{\pi^2}{8H^2(s)}\right) \frac{ds}{sH^2(s)} < \infty.$$

To put the preceding in perspective define

$$(1.12) \quad H_{\nu}(t) := \frac{\pi}{\sqrt{8(\ln_+ \ln_+ t + \nu \ln_+ \ln_+ \ln_+ t)}} \quad \forall t, \nu > 0.$$

[$1/0 := \infty$] Then, we can deduce from Corollaries 1.2 and 1.3 that $\mathcal{L}(H_{\nu})$ occurs q.s. iff $\nu > 5$, whereas $\mathcal{L}(H_{\nu})$ occurs a.s. iff $\nu > 2$. In particular, $\mathcal{L}(H_{\nu})$ occurs a.s. but not q.s. if $\nu \in [2, 5)$. The following is another interesting consequence of Theorem 1.1.

Corollary 1.4. *Let $G \subseteq [0, 1]$ be a non-random Borel set. Then,*

$$(1.13) \quad \begin{aligned} \dim_{\varphi} G > \frac{\nu - 2}{3} &\implies \text{cap}_G\left((\mathcal{L}(H_{\nu}))^{\mathbb{G}}\right) > 0, \text{ whereas} \\ \dim_{\varphi} G < \frac{\nu - 2}{3} &\implies \text{cap}_G\left((\mathcal{L}(H_{\nu}))^{\mathbb{G}}\right) = 0. \end{aligned}$$

Here, $\dim_{\varphi} G$ denotes the *packing dimension* (Mattila, 1995) of the set G .

Throughout this paper, uninteresting constants are denoted by a, b, α, A , etc. Their values may change from line to line.

Acknowledgements. We wish to thank Professor Zhan Shi for generously sharing with us the English translation of Lifshits and Shi (2003).

2. BROWNIAN SHEET, AND CAPACITY IN WIENER SPACE

We will be working with a special construction of the process U . This construction is due to D. Williams (Meyer, 1982, Appendix).

Let $B := \{B(s, t)\}_{s, t \geq 0}$ denote a two-parameter Brownian sheet. This means that B is a centered, continuous, Gaussian process with

$$(2.1) \quad \text{Cov}(B(s, t), B(s', t')) = \min(s, s') \times \min(t, t') \quad \forall s, s', t, t' \geq 0.$$

The Ornstein–Uhlenbeck process $U = \{U_s\}_{s \geq 0}$ on $C(\mathbf{R}_+)$ is precisely the infinite-dimensional process that is defined by

$$(2.2) \quad U_s(t) = \frac{B(e^s, t)}{e^{s/2}} \quad \forall s, t \geq 0.$$

Indeed, one can check directly that U is a $C(\mathbf{R}_+)$ -valued, stationary, symmetric diffusion. And for every $t \geq 0$, $\{U_s(t)\}_{s \geq 0}$ solves the stochastic differential equation (1.1) of the Ornstein–Uhlenbeck type. Furthermore, the invariant measure of U is the Wiener measure.

The following well-known result is a useful localization tool.

Lemma 2.1. *For all bounded Borel sets $G \subseteq \mathbf{R}_+$ and $\Lambda \in \mathcal{B}$, $\text{cap}_G(\Lambda) > 0$ iff with positive probability there exists $s \in G$ such that $U_s \in \Lambda$.*

Remark 2.2. The previous lemma continues to hold even when G is unbounded.

Proof. Without loss of much generality, we may—and will—assume that $G \subseteq [0, q]$ for some $q > 0$. Let $p_G(\Lambda)$ denote the probability that there exists $s \in G$ such that $U_s \in \Lambda$. Evidently, $\text{cap}_G(\Lambda) \leq p_G(\Lambda)$. Furthermore, $\text{cap}_G(\Lambda) = \int_0^q \mathbb{P}\{\exists s \in G \cap [0, \tau] : U_s \in \Lambda\} e^{-\tau} d\tau + e^{-q} p_G(\Lambda)$, whence the bounds,

$$(2.3) \quad e^{-q} p_G(\Lambda) \leq \text{cap}_G(\Lambda) \leq p_G(\Lambda).$$

The lemma follows. □

Define

$$(2.4) \quad f^* := \sup_{u \in [0, 1]} |f(u)| \quad \forall f \in C(\mathbf{R}_+).$$

The following is the main step in the proof of Theorem 1.1. It was announced earlier in the Abstract.

Theorem 2.3. *There exists $a > 1$ such that for all $r \in (0, 1)$ and all Borel sets $G \subseteq [0, 1]$,*

$$(2.5) \quad \frac{1}{a} \mathbf{K}_G(r^6) e^{-\pi^2/(8r^2)} \leq \text{cap}_G \{f^* \leq r\} \leq a \mathbf{K}_G(r^6) e^{-\pi^2/(8r^2)}.$$

Remark 2.4. The constant a depends on G only through the fact that G is a subset of $[0, 1]$. Therefore, there exists $a > 1$ such that simultaneously for all Borel sets $F, G \subseteq [0, 1]$,

$$(2.6) \quad \frac{1}{a} \frac{\mathbf{K}_F(r^6)}{\mathbf{K}_G(r^6)} \leq \frac{\text{cap}_F \{f^* \leq r\}}{\text{cap}_G \{f^* \leq r\}} \leq a \frac{\mathbf{K}_F(r^6)}{\mathbf{K}_G(r^6)} \quad \forall r \in (0, 1).$$

Remark 2.5. It turns out that for any fixed $\varepsilon > 0$, $\text{cap}_{\mathbf{R}_+}$ and $\text{cap}_{[0,\varepsilon]}$ are equivalent. To prove this, we can assume without loss of generality that $\varepsilon \in (0, 1)$. [This is because $\varepsilon \mapsto \text{cap}_{[0,\varepsilon]}(\Lambda)$ is increasing.] Now, on one hand, $\text{cap}_{[0,\varepsilon]}(\Lambda) \leq \text{cap}_{\mathbf{R}_+}(\Lambda)$. On the other hand,

$$(2.7) \quad \begin{aligned} \text{cap}_{\mathbf{R}_+}(\Lambda) &\leq \int_0^\infty \sum_{0 \leq j \leq \sigma/\varepsilon} \mathbb{P} \{ \exists s \in [j\varepsilon, (j+1)\varepsilon] : U_s \in \Lambda \} e^{-\sigma} d\sigma \\ &\leq \mathbb{P} \{ \exists s \in [0, \varepsilon] : U_s \in \Lambda \} \int_0^\infty \frac{\sigma+1}{\varepsilon} e^{-\sigma} d\sigma, \end{aligned}$$

by stationarity. In the notation of Lemma 2.1, the last term is $(2/\varepsilon)p_{[0,\varepsilon]}(\Lambda) \leq (2e/\varepsilon)\text{cap}_{[0,\varepsilon]}(\Lambda)$; cf. (2.3). Thus,

$$(2.8) \quad \frac{\varepsilon}{2e} \text{cap}_{\mathbf{R}_+}(\Lambda) \leq \text{cap}_{[0,\varepsilon]}(\Lambda) \leq \text{cap}_{\mathbf{R}_+}(\Lambda) \quad \forall \Lambda \in \mathcal{B}.$$

This proves amply the claimed equivalence of $\text{cap}_{[0,\varepsilon]}$ and $\text{cap}_{\mathbf{R}_+}$.

According to the eigenfunction expansion of Chung (1948),

$$(2.9) \quad \mu \{ f^* \leq r \} \sim \frac{4}{\pi} e^{-\pi^2/(8r^2)} \quad (r \rightarrow 0).$$

Therefore, thanks to (2.3), Theorem 2.3 is equivalent to our next result.

Theorem 2.6. *Recall that $U_s^* = \sup_{t \in [0,1]} |U_s(t)|$ [eq. (2.4)]. Then, there exists a constant $a > 1$ such that for all $r \in (0, 1)$ and all Borel sets $G \subseteq [0, 1]$,*

$$(2.10) \quad \frac{1}{a} \text{K}_G(r^6) \mu \{ f^* \leq r \} \leq \mathbb{P} \left\{ \inf_{s \in G} U_s^* \leq r \right\} \leq a \text{K}_G(r^6) \mu \{ f^* \leq r \}.$$

We will derive this particular reformulation of Theorem 2.3. The following result plays a key role in our analysis.

Proposition 2.7 (Lifshits and Shi (2003, Proposition 2.1)). *Let $\{X_t\}_{t \geq 0}$ denote planar Brownian motion. For every $r > 0$ and $\lambda \in (0, 1]$ define*

$$(2.11) \quad \mathcal{D}_\lambda^r = \left\{ (x, y) \in \mathbf{R}^2 : |x| \leq r, \left| x\sqrt{1-\lambda} + y\sqrt{\lambda} \right| \leq r \right\}.$$

Then there exists an $a \in (0, 1/2)$ such that for all $r > 0$ and $\lambda \in (0, 1]$,

$$(2.12) \quad \mathbb{P} \{ X_t \in \mathcal{D}_\lambda^r \quad \forall t \in [0, 1] \} \leq \frac{1}{a} \mu \{ f^* \leq r \} e^{-a\lambda^{1/3}/r^2}.$$

Lemma 2.8. *There exists a constant $a \in (0, 1)$ such that for all $1 \geq S > s > 0$,*

$$(2.13) \quad \mathbb{P} \{ U_s^* \leq r, U_S^* \leq r \} \leq \frac{1}{a} \mu \{ f^* \leq r \} e^{-a(S-s)^{1/3}/r^2} \quad \forall r \in (0, 1).$$

Proof. Define $\lambda = 1 - e^{-(S-s)}$. Then owing to (2.2) we can write

$$(2.14) \quad U_S(t) = U_s(t)\sqrt{1-\lambda} + \frac{B(e^S, t) - B(e^s, t)}{\sqrt{e^S - e^s}} \sqrt{\lambda} := U_s(t)\sqrt{1-\lambda} + V(t)\sqrt{\lambda}.$$

By the Markov properties of the Brownian sheet, $X_t := (U_s(t), V(t))$ defines a planar Brownian motion. Moreover, $\mathbb{P} \{ U_s^* \leq r, U_S^* \leq r \} = \mathbb{P} \{ X_t \in \mathcal{D}_\lambda^r, \forall t \in [0, 1] \}$. By Taylor's expansion, $1 - e^{-x} \geq (x/2)$ ($x \in [0, 1]$). Therefore, Proposition 2.7 completes the proof. \square

Proof of Theorem 2.6: Lower Bound. Let $k = K_G(r^6)$, and choose maximal Kolmogorov points $s(1) < \dots < s(k)$ such that $s(i+1) - s(i) \geq r^6$. Evidently, whenever $j > i$ we have $s(j) - s(i) \geq (j-i)r^6$. Now define

$$(2.15) \quad N_r = \sum_{i=1}^k \mathbf{1}_{\{U_{s(i)}^* \leq r\}}.$$

According to Lemma 2.8,

$$(2.16) \quad \begin{aligned} \mathbb{E}[N_r^2] &= k\mu\{f^* \leq r\} + 2 \sum_{i=1}^{k-1} \sum_{j=i+1}^k \mathbb{P}\{U_{s(i)}^* \leq r, U_{s(j)}^* \leq r\} \\ &\leq k\mu\{f^* \leq r\} + \frac{2}{a}\mu\{f^* \leq r\} \sum_{i=1}^{k-1} \sum_{j=i+1}^k \exp\left(-\frac{a(s(j) - s(i))^{1/3}}{r^2}\right) \\ &\leq k\mu\{f^* \leq r\} + \frac{2}{a}\mu\{f^* \leq r\} \sum_{i=1}^{k-1} \sum_{j=i+1}^k \exp\left(-a(j-i)^{1/3}\right) \\ &\leq Ak\mu\{f^* \leq r\}. \end{aligned}$$

Note that A is a positive and finite constant that does not depend on r . Also note that $\mathbb{E}[N_r] = k\mu\{f^* \leq r\}$. This and the Paley–Zygmund inequality (Khoshnevisan, 2002, Lemma 1.4.1, p. 72) together reveal that

$$(2.17) \quad \mathbb{P}\left\{\inf_{s \in G} U_s^* \leq r\right\} \geq \mathbb{P}\{N_r > 0\} \geq \frac{(\mathbb{E}[N_r])^2}{\mathbb{E}[N_r^2]} \geq \frac{k}{A}\mu\{f^* \leq r\}.$$

The definition of k implies the lower bound in Theorem 2.6. \square

Before proving the upper bound of Theorem 2.6 in complete generality, we first derive the following weak form:

Proposition 2.9. *There exists a finite constant $a > 1$ such that for all $r \in (0, 1)$, $\mathbb{P}\{\inf_{s \in [0, r^6]} U_s^* \leq r\} \leq a\mu\{f^* \leq r\}$.*

Proof. Recall (2.15), and define

$$(2.18) \quad L(s; r) = \int_0^s \mathbf{1}_{\{U_\nu^* \leq r\}} d\nu \quad \forall s, r > 0.$$

Let $\mathcal{F} := \{\mathcal{F}_s\}_{s \geq 0}$ denote the augmented filtration generated by the infinite-dimensional process $\{U_s\}_{s \geq 0}$. The latter process is Markov with respect to \mathcal{F} . Moreover,

$$(2.19) \quad \mathbb{E}[L(2r^6; r + r^3) \mid \mathcal{F}_s] \geq \int_s^{2r^6} \mathbb{P}\{U_\nu^* \leq r + r^3 \mid \mathcal{F}_s\} d\nu \cdot \mathbf{1}_{\{U_s^* \leq r\}}.$$

As in (2.14), if $\nu > s$ are fixed, then we can write

$$(2.20) \quad \begin{aligned} U_\nu(t) &= U_s(t)e^{-(\nu-s)/2} + \frac{B(e^\nu, t) - B(e^s, t)}{\sqrt{e^\nu - e^s}} \sqrt{1 - e^{-(\nu-s)}} \\ &:= U_s(t)e^{-(\nu-s)/2} + V(t)\sqrt{1 - e^{-(\nu-s)}}. \end{aligned}$$

We emphasize, once again, that (U_s, V) is a planar Brownian motion. In addition, V is independent of \mathcal{F}_s , and $U_\nu^* \leq U_s^* + V^* \sqrt{1 - \exp\{-(\nu - s)\}}$. Consequently, as long as $0 \leq s \leq r^6$ and $s < \nu < 2r^6$,

$$(2.21) \quad U_\nu^* \leq U_s^* + \frac{r^3}{\sqrt{2}} V^*.$$

[We have used the inequality $1 - e^{-z} \leq z/2$ valid for all $z \in (0, 1)$.] Therefore, for all $0 \leq s \leq r^6$,

$$(2.22) \quad \begin{aligned} M(s) &= \mathbb{E} [L(2r^6; r + r^3) \mid \mathcal{F}_s] \\ &\geq \int_s^{2r^6} \mathbb{P} \{V^* \leq \sqrt{2}\} d\nu \cdot \mathbf{1}_{\{U_s^* \leq r\}} \\ &= \mu \{f^* \leq \sqrt{2}\} (2r^6 - s) \cdot \mathbf{1}_{\{U_s^* \leq r\}} \\ &\geq \mu \{f^* \leq \sqrt{2}\} r^6 \cdot \mathbf{1}_{\{U_s^* \leq r\}}. \end{aligned}$$

Because $\{M(s)\}_{s \geq 0}$ is a martingale, we can apply Doob's maximal inequality to obtain the following:

$$(2.23) \quad \begin{aligned} \mathbb{P} \left\{ \inf_{s \in [0, r^6]} U_s^* \leq r \right\} &\leq \mathbb{P} \left\{ \sup_{s \in [0, r^6]} M(s) \geq \mu \{f^* \leq \sqrt{2}\} r^6 \right\} \\ &\leq \frac{\mathbb{E} [L(2r^6; r + r^3)]}{\mu \{f^* \leq \sqrt{2}\} r^6} = \frac{2\mu \{f^* \leq r + r^3\}}{\mu \{f^* \leq \sqrt{2}\}}. \end{aligned}$$

Thanks to (2.9),

$$(2.24) \quad \frac{\mu \{f^* \leq r + r^3\}}{\mu \{f^* \leq r\}} \sim \exp \left(-\frac{\pi^2}{8} \left[\frac{1}{(r + r^3)^2} - \frac{1}{r^2} \right] \right) \rightarrow e^{\pi^2/4}. \quad (r \rightarrow 0).$$

Thus, the left-hand side is bounded ($r \in (0, 1)$), and the proposition follows. \square

Proof of Theorem 2.6: Upper Bound. Define $n = n(r)$ to be $\lfloor r^{-6} \rfloor$, and define $I(j; n)$ to be the interval $[j/n, (j+1)/n)$ ($j = 0, \dots, n$). Then, by stationarity and Proposition 2.9,

$$(2.25) \quad \mathbb{P} \left\{ \inf_{s \in G} U_s^* \leq r \right\} \leq \sum_{\substack{0 \leq j \leq n: \\ I(j; n) \cap G \neq \emptyset}} \mathbb{P} \left\{ \inf_{s \in I(j; n)} U_s^* \leq r \right\} \leq a\mu \{f^* \leq r\} M_n(G),$$

where $M_n(G) = \#\{0 \leq j \leq n : I(j; n) \cap G \neq \emptyset\}$ defines the *Minkowski content* of G . In the companion to this paper (2004, Proposition 2.7) we proved that $M_n(G) \leq 3K_G(1/n)$. By monotonicity, the latter is at most $3K_G(r^6)$, whence the theorem. \square

3. PROOF OF THEOREM 1.1 AND COROLLARIES 1.2 AND 1.4

We begin with some preliminary discussions. Define

$$(3.1) \quad \psi_H(G) := \int_1^\infty \frac{K_G(H^6(s))}{sH^2(s)} \exp \left(-\frac{\pi^2}{8H^2(s)} \right) ds, \quad \sigma(r) := \mu \{f^* \leq r\}.$$

Following Erdős (1942), define

$$(3.2) \quad \mathbf{e}_n = e^{n/\ln_+ n}, \quad H_n = H(\mathbf{e}_n) \quad \forall n \geq 1.$$

The “critical” function in (1.11) is $H^2(t) = \pi^2/(8 \ln_+ \ln_+ t)$. This, the fact that $\pi/\sqrt{8} \in (1, 2)$, and a familiar argument (Erdős, 1942, equations 1.2 and 3.4), together allow us to assume without loss of generality that

$$(3.3) \quad \frac{1}{\sqrt{\ln_+ n}} \leq H_n \leq \frac{2}{\sqrt{\ln_+ n}} \quad \forall n \geq 1.$$

From this we can conclude the existence of a constant $a > 1$ such that

$$(3.4) \quad \frac{1}{a} H_n^2 \mathbf{e}_{n+1} \leq \mathbf{e}_{n+1} - \mathbf{e}_n \leq a H_{n+1}^2 \mathbf{e}_n \quad \forall n \geq 1.$$

According to our companion work (2004, eq. 2.8), for all $r > 0$ sufficiently small,

$$(3.5) \quad K_G(\varepsilon) \leq 6K_G(2\varepsilon).$$

Because $\mathbf{e}_{n+1} \sim \mathbf{e}_n$ as $n \rightarrow \infty$, (2.9), (3.4), and (3.5) together imply that

$$(3.6) \quad \sum_{n=1}^{\infty} K_G(H_n^6) \sigma(H_n) < \infty \quad \text{iff} \quad \psi_H(G) < \infty.$$

The following is the key step toward proving Theorem 1.1.

Proposition 3.1. *Let $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ be decreasing and measurable. Then for all non-random Borel sets $G \subseteq [0, 1]$,*

$$(3.7) \quad \liminf_{t \rightarrow \infty} \left(\inf_{s \in G} \sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t} \right) = \begin{cases} +\infty, & \text{if } \psi_H(G) < \infty, \\ -\infty, & \text{if } \psi_H(G) = \infty. \end{cases}$$

First we assume this proposition and derive Theorem 1.1. Then, we will tidy things up by proving the technical Proposition 3.1.

Let us recall (3.1).

Definition 3.2. We say that $\Psi_H(G) < \infty$ if we can decompose G as $G = \cup_{n=1}^{\infty} G_n$ —where G_1, G_2, \dots are closed—such that for all $n \geq 1$, $\psi_H(G_n) < \infty$. Else, we say that $\Psi_H(G) = \infty$.

Let us first rephrase Theorem 1.1 in the following convenient, and equivalent, form.

Proposition 3.3. *Let $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ be decreasing and measurable and $G \subseteq [0, 1]$ be non-random and Borel. If $\Psi_H(G) < \infty$, then*

$$(3.8) \quad \inf_{s \in G} \liminf_{t \rightarrow \infty} \left(\sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t} \right) = \infty \quad \text{P-a.s.}$$

Else, the left-hand side is P-a.s. equal to $-\infty$.

Proof of Theorem 1.1 in the form of Proposition 3.3. First suppose $\Psi_H(G)$ is finite. We can write $G = \cup_{n=1}^{\infty} G_n$, where the G_n 's are closed and $\psi_H(G_n) < \infty$ for all $n \geq 1$. Then, according to Proposition 3.1,

$$(3.9) \quad \begin{aligned} & \inf_{s \in G_n} \liminf_{t \rightarrow \infty} \left[\sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t} \right] \\ & \geq \liminf_{t \rightarrow \infty} \inf_{s \in G_n} \left[\sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t} \right] = \infty. \end{aligned}$$

This proves that $\inf_{s \in G} \liminf_{t \rightarrow \infty} (\sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t}) = \infty$ a.s. [P].

For the converse portion suppose $\Psi_H(G) = \infty$, and choose arbitrary non-random closed sets $\{G_n\}_{n=1}^\infty$ such that $\cup_{n=1}^\infty G_n = G$. By definition, $\psi_H(G_n) = \infty$ for some $n \geq 1$. Define for all $T \geq 1$,

$$(3.10) \quad \mathcal{S}_T := \left\{ s \in [0, 1] : \inf_{t \geq T} \frac{\sup_{u \in [0, t]} |U_s(u)|}{H(t)\sqrt{t}} \leq 1 \right\}.$$

Evidently, \mathcal{S}_T is a random set for each $T \geq 0$. Moreover, the continuity of the Brownian sheet implies that with probability one, \mathcal{S}_T is closed for all T ; hence, so is $\mathcal{S}_T \cap G_n$. Because $\psi_H(G_n) = \infty$, Proposition 3.1 implies that almost surely, $\mathcal{S}_T \cap G_n \neq \emptyset$. Since $\{\mathcal{S}_T \cap G_n\}_{T=1}^\infty$ is a decreasing sequence of non-void compact sets, they have non-void intersection. That is, $(\cap_{T=1}^\infty \mathcal{S}_T) \cap G_n \neq \emptyset$ a.s. [P]. Replace H by $H - H^3$ to complete the proof of Proposition 3.3. \square

Now we derive Proposition 3.1. This completes our proof of Theorem 1.1. Our proof is divided naturally into two halves.

Proof of Proposition 3.1: First Half. Throughout this portion of the proof, we assume that $\psi_H(G) < \infty$.

Because $\mathbf{e}_{n+1} \sim \mathbf{e}_n$ as $n \rightarrow \infty$, Theorem 2.6 and Brownian scaling together imply that

$$(3.11) \quad \begin{aligned} \mathbb{P} \left\{ \inf_{s \in G} \sup_{u \in [0, \mathbf{e}_{n-1}]} |U_s(u)| \leq H_n \sqrt{\mathbf{e}_n} \right\} &= \mathbb{P} \left\{ \inf_{s \in G} U_s^* \leq H_n \sqrt{\mathbf{e}_n / \mathbf{e}_{n-1}} \right\} \\ &\leq aK_G \left(H_n^6 \left[\frac{\mathbf{e}_n}{\mathbf{e}_{n-1}} \right]^3 \right) \sigma \left(H_n \sqrt{\frac{\mathbf{e}_n}{\mathbf{e}_{n-1}}} \right). \end{aligned}$$

According to (3.5), $K_G(\dots) \leq 6K_G(H_n^6)$ for all n large. This and (3.4) together imply that for all n large,

$$(3.12) \quad \begin{aligned} \mathbb{P} \left\{ \inf_{s \in G} \sup_{u \in [0, \mathbf{e}_{n-1}]} |U_s(u)| \leq H_n \sqrt{\mathbf{e}_n} \right\} \\ \leq aK_G(H_n^6) \sigma \left(H_n \sqrt{1 + AH_n^2} \right) \\ \leq aK_G(H_n^6) \sigma \left(H_n [1 + AH_n^2] \right). \end{aligned}$$

In accord with (2.9), for any fixed $c \in \mathbf{R}$,

$$(3.13) \quad \sigma(r + cr^3) = O(\sigma(r)) \quad (r \rightarrow 0).$$

Thus, for all $n \geq 1$,

$$(3.14) \quad \mathbb{P} \left\{ \inf_{s \in G} \sup_{u \in [0, \mathbf{e}_{n-1}]} |U_s(u)| \leq H_n \sqrt{\mathbf{e}_n} \right\} \leq aK_G(H_n^6) \sigma(H_n).$$

Because we are assuming that $\psi_H(G)$ is finite, (3.6) and the Borel–Cantelli lemma together imply that almost surely, $\inf_{s \in G} \sup_{u \in [0, \mathbf{e}_{n-1}]} |U_s(u)| > H_n \sqrt{\mathbf{e}_n}$ for all but a finite number of n 's. It follows from this and a standard monotonicity argument that

$$(3.15) \quad \psi_H(G) < \infty \implies \liminf_{t \rightarrow \infty} \left[\inf_{s \in G} \sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t} \right] > 0 \text{ a.s. [P].}$$

But if $\psi_H(G)$ were finite then $\psi_{H+H^3}(G)$ is also finite; compare (3.5) and (3.13). Thanks to (3.3), $\lim_{t \rightarrow \infty} H^3(t)\sqrt{t} = \infty$. Therefore, the lim inf of the preceding display is infinity. This concludes the first half of our proof of Proposition 3.1. \square

In order to prove the second half of Proposition 3.1 we assume that $\psi_H(G) = \infty$, recall (3.1), and define

$$(3.16) \quad \begin{aligned} L_n &:= \left\{ \inf_{s \in G} \sup_{u \in [0, \mathbf{e}_n]} |U_s(u)| \leq H_n \sqrt{\mathbf{e}_n} \right\}, \\ f(z) &:= K_G(z^6) \sigma(z). \end{aligned}$$

Lemma 3.4. *Define for all $j \geq i$, $\lambda_{i,j} := \mathbf{e}_j / (\mathbf{e}_j - \mathbf{e}_i)$ and $\delta_{i,j} := H_j \sqrt{\lambda_{i,j}} + H_i \sqrt{\lambda_{i,j} - 1}$. Then, there exists $a > 1$ such that for all $j \geq i$, $P(L_j | L_i) \leq a K_G(\delta_{i,j}^6) \sigma(\delta_{i,j})$.*

Proof. Evidently, $P(L_j | L_i)$ is at most

$$(3.17) \quad \begin{aligned} &P \left\{ \inf_{s \in G} \sup_{u \in [\mathbf{e}_i, \mathbf{e}_j]} |U_s(u)| \leq H_j \sqrt{\mathbf{e}_j} \mid L_i \right\} \\ &= P \left\{ \inf_{s \in G} \sup_{u \in [\mathbf{e}_i, \mathbf{e}_j]} |U_s(u) - U_s(\mathbf{e}_i) + U_s(\mathbf{e}_i)| \leq H_j \sqrt{\mathbf{e}_j} \mid L_i \right\} \\ &\leq P \left\{ \inf_{s \in G} \sup_{u \in [\mathbf{e}_i, \mathbf{e}_j]} |U_s(u) - U_s(\mathbf{e}_i)| \leq H_j \sqrt{\mathbf{e}_j} + H_i \sqrt{\mathbf{e}_i} \right\}. \end{aligned}$$

We have appealed to the Markov properties of the Brownian sheet in the last line. Because $u \mapsto U_\bullet(u)$ is a $C(\mathbf{R}_+)$ -valued Brownian motion,

$$(3.18) \quad \begin{aligned} P(L_j | L_i) &\leq P \left\{ \inf_{s \in G} \sup_{u \in [0, \mathbf{e}_j - \mathbf{e}_i]} |U_s(u)| \leq H_j \sqrt{\mathbf{e}_j} + H_i \sqrt{\mathbf{e}_i} \right\} \\ &= P \left\{ \inf_{s \in G} U_s^* \leq \delta_{i,j} \right\}. \end{aligned}$$

Theorem 2.6 completes the proof. \square

Our forthcoming estimates of $P(L_j | L_i)$ rely on the following elementary bound; see, for example, our earlier work (2003, eq. 8.30): Uniformly for all integers $j > i$,

$$(3.19) \quad \mathbf{e}_j - \mathbf{e}_i \geq \mathbf{e}_i \left(\frac{j-i}{\ln i} \right) (1 + o(1)) \quad (i \rightarrow \infty).$$

Lemma 3.5. *There exist $i_0 \geq 1$ and a finite $a > 1$ such that for all $i \geq i_0$ and $j \geq i + \ln^{19}(j)$,*

$$(3.20) \quad P(L_j | L_i) \leq a P(L_j).$$

Proof. Thanks to (3.3) and (3.19), the following holds uniformly over all $j > i + \ln^{19}(j)$: $(\mathbf{e}_j / \mathbf{e}_i) \geq (1 + o(1)) H_j^{-36}$ ($i \rightarrow \infty$). Thus, uniformly over all $j > i + \ln^{19}(j)$,

$$(3.21) \quad \begin{aligned} \sqrt{\lambda_{i,j}} &= \frac{1}{\sqrt{1 - (\mathbf{e}_i / \mathbf{e}_j)}} \leq \frac{1}{\sqrt{1 - (1 + o(1)) H_j^{36}}} = 1 + O(H_j^3), \\ H_i \sqrt{\lambda_{i,j} - 1} &= O(H_j^3) \quad (i \rightarrow \infty). \end{aligned}$$

Lemma 3.4 guarantees then that uniformly over all $j > i + \ln^{19}(j)$, $\delta_{i,j} \leq H_j + O(H_j^3)$, and the big- O and little- o terms do not depend on the j 's in question. The lemma follows from this, equations (3.5) and (3.13), and Theorem 2.6. \square

Lemma 3.6. *There exist $i_1 \geq 1$ and $a \in (0, 1)$ such that for all $i \geq i_1$ and $j \in [i + \ln(i), i + \ln^{19}(j))$, $P(L_j | L_i) \leq (aj^a)^{-1}$.*

Proof. Equations (3.19) and (3.3) together imply that uniformly for all $j \geq i + \ln(i)$, $(\mathbf{e}_i/\mathbf{e}_j) \leq \frac{1}{2} + o(1)$ ($i \rightarrow \infty$). This is equivalent to the existence of a constant $A_{3.22}$ such that for all (i, j) in the range of the lemma,

$$(3.22) \quad \sqrt{\lambda_{i,j}} \vee \sqrt{\lambda_{i,j} - 1} \leq a.$$

Thanks to (3.3), we can enlarge the last constant a , if necessary, to ensure that for all (i, j) in the range of this lemma, $H_i \leq aH_j$. Therefore, Lemma 3.4 then implies that $\delta_{i,j} = O(H_j)$, and the big- O term does not depend on the range of j 's in question. Because $G \subseteq [0, 1]$,

$$(3.23) \quad K_G(\varepsilon) \leq K_{[0,1]}(\varepsilon) \sim 1/\varepsilon \quad (\varepsilon \rightarrow 0).$$

Thus, Lemma 3.4 ensures that $P(L_j | L_i) \leq a\delta_{i,j}^{-6}\sigma(\delta_{i,j})$. Near the origin, the function $\delta \mapsto \delta^{-6}\sigma(\delta)$ is increasing. Because we have proved that over the range of (i, j) of this lemma $\delta_{i,j} = O(H_j)$, equation (2.9) asserts the existence of a universal $\alpha > 1$ such that $P(L_j | L_i)$ is at most $\alpha H_j^{-6} \exp(-\alpha^{-1}H_j^{-2})$. Equation (3.3) then completes our proof. \square

Lemma 3.7. *There exist $i_2 \geq 1$ and $a > 1$ such that for all $i \geq i_2$ and $j \in (i, i + \ln i)$, $P(L_j | L_i) \leq ae^{-(j-i)/a}$.*

Proof. By (3.19), $(\mathbf{e}_i/\mathbf{e}_j) \leq 1 - (1 + o(1))(j - i) \ln^{-1}(i)$ ($i \rightarrow \infty$), where the little- o term does not depend on $j \in (i, i + \ln i)$. Similarly, $(\mathbf{e}_j/\mathbf{e}_i) \geq 1 + (1 + o(1))(j - i) \ln^{-1}(i)$. Thus, as $i \rightarrow \infty$,

$$(3.24) \quad \begin{aligned} \sqrt{\lambda_{i,j}} &= \frac{1}{\sqrt{1 - (\mathbf{e}_i/\mathbf{e}_j)}} \leq (1 + o(1)) \sqrt{\frac{\ln i}{j - i}} \leq \frac{2 + o(1)}{H_j \sqrt{j - i}}, \\ \sqrt{\lambda_{i,j} - 1} &= \frac{1}{\sqrt{(\mathbf{e}_j/\mathbf{e}_i) - 1}} \leq (1 + o(1)) \sqrt{\frac{\ln i}{j - i}} \leq \frac{2 + o(1)}{H_j \sqrt{j - i}}, \end{aligned}$$

by (3.3). Once again, the little- o terms are all independent of $j \in (i, i + \ln i)$. Because $H_i = O(H_j)$ uniformly for all (i, j) in the range considered here, Lemma 3.4 implies that uniformly for all $j \in (i, i + \ln i)$, $\delta_{i,j} = O(1/\sqrt{j - i})$. Equation (3.23) bounds the first term on the right-hand side; (2.9) bounds the second. This and (3.3) together prove the existence of a constant $\alpha > 1$ such that for all $i \geq i_2$ and all $j \in (i, i + \ln i)$, $P(L_j | L_i) \leq \alpha(j - i)^3 \exp\{-(j - i)/\alpha\}$. The lemma follows. \square

Proof of Proposition 3.1: Second Half. According to Theorem 2.6, for all n large enough, $P(L_n) \geq af(H_n)$. Because $\psi_H(G) = \infty$, the latter estimate and (3.6) together imply that

$$(3.25) \quad \sum_{i=1}^{\infty} P(L_i) = \infty.$$

Thus, our derivation is complete once we demonstrate the following:

$$(3.26) \quad \liminf_{n \rightarrow \infty} \frac{\sum_{i=1}^{n-1} \sum_{j=i}^n \mathbb{P}(L_i \cap L_j)}{(\sum_{i=1}^n \mathbb{P}(L_i))^2} < \infty.$$

See Chung and Erdős (1952). In fact, the preceding display holds with a lim sup in place of the lim inf. This fact follows from combining, using standard arguments, Lemmas 3.5 through 3.7.

Indeed, let $I := \max(3, i_1, i_2, i_3)$ and $s_n := \sum_{i=1}^n \mathbb{P}(L_i)$. Lemma 3.5 ensures that

$$(3.27) \quad \sum_{i=I}^{n-1} \sum_{\substack{j=i \\ j > i + \ln^{19}(j)}}^n \mathbb{P}(L_j \cap L_i) = O(s_n^2).$$

By Lemma 3.6,

$$(3.28) \quad \begin{aligned} \sum_{i=I}^{n-1} \sum_{\substack{j=i \\ j \in (i + \ln(i), i + \ln^{19}(j)]}}^n \mathbb{P}(L_j \cap L_i) &\leq \frac{1}{a} \sum_{i=I}^{n-1} \sum_{\substack{j=i \\ j \in (i + \ln(i), i + \ln^{19}(j)]}}^n j^{-a} \mathbb{P}(L_i) \\ &= \sum_{i=I}^n O\left(\frac{\ln^{19}(i)}{i^a}\right) \mathbb{P}(L_i) = O(s_n). \end{aligned}$$

The big- O terms do not depend on the variables (j, n) .

Finally, Lemma 3.7 implies that

$$(3.29) \quad \sum_{i=I}^{n-1} \sum_{\substack{j=i \\ j \in (i, i + \ln i]}}^n \mathbb{P}(L_j \cap L_i) \leq a \sum_{i=I}^n \sum_{j=i}^{\infty} \mathbb{P}(L_i) e^{(j-i)/a} = O(s_n).$$

We have already seen that $s_n \rightarrow \infty$. Thus, (3.27)–(3.29) imply (3.26), and hence the theorem. More precisely, we have proved so far that

$$(3.30) \quad \psi_H(G) = \infty \implies \liminf_{t \rightarrow \infty} \left[\inf_{s \in G} \sup_{u \in [0, t]} |U_s(u)| - H(t) \sqrt{t} \right] < 0 \text{ a.s. } [\mathbb{P}].$$

Replace H by $H + H^3$ to deduce that the preceding lim inf is in fact $-\infty$. This completes our proof of Proposition 3.1. \square

We conclude this section by proving the remaining Corollaries 1.2 and 1.4.

Proof of Corollary 1.2. By definition, $\mathcal{L}(H)$ holds q.s. iff $\text{cap}_{\mathbf{R}_+}((\mathcal{L}(H))^{\mathfrak{c}}) = 0$. Thanks to Theorem 1.1, this condition is equivalent to the existence of a non-random “closed-denumerable” decomposition $\mathbf{R}_+ = \cup_{n=1}^{\infty} G_n$ such that for all $n \geq 1$, $\psi_H(G_n) < \infty$. But one of the G_n ’s must contain a closed interval that has positive length. Therefore, by the translation-invariance of $G \mapsto K_G(r)$, there exists $\varepsilon \in (0, 1)$ such that $\psi_H([0, \varepsilon]) < \infty$.

Conversely, if $\psi_H([0, \varepsilon])$ is finite, then we can define G_n to be $[(n-1)\varepsilon, n\varepsilon]$ ($n \geq 1$) to find that $\psi_H(G_n) = \psi_H([0, \varepsilon]) < \infty$. Theorem 1.1 then proves that $\text{cap}_{\mathbf{R}_+}((\mathcal{L}(H))^{\mathfrak{c}}) = 0$ iff there exists $\varepsilon > 0$ such that $\psi_H([0, \varepsilon]) < \infty$. Because $K_{[0, \varepsilon]}(r) \sim \varepsilon/r$ ($r \rightarrow 0$), the corollary follows. \square

Proof of Corollary 1.4. We can change variables to deduce that $\psi_{H_\nu}(G)$ is finite iff $\int_1^\infty K_G(1/s)s^{-1-(\nu/3)} ds$ converges. This and Proposition 2.8 of our companion work (2004) together imply that

$$(3.31) \quad \inf\{\nu > 0 : \psi_{H_\nu}(G) < \infty\} = 2 + 3\overline{\dim}_{\mathcal{M}} G,$$

where $\overline{\dim}_{\mathcal{M}}$ denotes the (upper) Minkowski dimension (Mattila, 1995). By regularization (Mattila, 1995, p. 81),

$$(3.32) \quad \inf\{\nu > 0 : \Psi_{H_\nu}(G) < \infty\} = 2 + 3 \dim_{\mathcal{D}} G.$$

Theorem 1.1 now implies Corollary 1.4. \square

REFERENCES

- Chung, Kai Lai. 1948. *On the maximum partial sums of sequences of independent random variables*, Trans. Amer. Math. Soc. **64**, 205–233.
- Chung, K. L. and P. Erdős. 1952. *On the application of the Borel-Cantelli lemma*, Trans. Amer. Math. Soc. **72**, 179–186.
- Dudley, R. M. 1973. *A Course in Empirical Processes*, École d'été de probabilités Saint-Flour, XII–1982, Springer, Berlin, pp. 1–142.
- Erdős, Paul. 1942. *On the law of the iterated logarithm*, Ann. Math. **43(2)**, 419–436.
- Fukushima, Masatoshi. 1984. *Basic properties of Brownian motion and a capacity on the Wiener space*, J. Math. Soc. Japan **36(1)**, 161–176.
- Khintchine, A. Ya. 1933. *Asymptotische Gesetz der Wahrscheinlichkeitsrechnung*, Springer, Berlin.
- Khoshnevisan, Davar. 2002. *Multiparameter Processes: An Introduction to Random Fields*, Springer, New York.
- Khoshnevisan, Davar, David A. Levin, and Pedro J. Méndez–Hernández. 2003. *On dynamical Gaussian random walks*, Ann. Probab. (To appear).
- . 2004. *Exceptional times and invariance for dynamical random walks* (Preprint).
- Lifshits, M. A. and Z. Shi. 2003. *Lower functions of an empirical process and of a Brownian sheet*, Teor. Veroyatnost. i Primenen. **48(2)**, 321–339. (Russian, with Russian summary)
- Malliavin, Paul. 1979. *Régularité de lois conditionnelles et calcul des variations stochastiques*, C.R. Acad. Sci. Paris, Sér. A-B **289**.
- Mattila, Pertti. 1995. *Geometry of Sets and Measures in Euclidean Spaces: Fractals and Rectifiability*, Cambridge University Press, Cambridge.
- Meyer, P.-A. 1982. *Note sur les processus d'Ornstein–Uhlenbeck (Appendice: Un resultat de D. Williams)*, Sémin. de Probab. XVI, Lec. Notes in Math., vol. 920, Springer, pp. 95–133.
- Mountford, T. S. 1992. *Quasi-everywhere upper functions*, Sémin. de Probab. XXVI, Lect. Notes in Math., vol. 1526, Springer, pp. 95–106.
- Tihomirov, V. M. 1963. *The works of A. N. Kolmogorov on ε -entropy of function classes and superpositions of functions*, Uspehi Mat. Nauk **18(5 (113))**, 55–92.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF UTAH, 155 S. 1400 E., SALT LAKE CITY, UT 84112-0090

E-mail address: davar@math.utah.edu

URL: <http://www.math.utah.edu/~davar>

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF UTAH, 155 S. 1400 E., SALT LAKE CITY, UT 84112-0090

E-mail address: levin@math.utah.edu

URL: <http://www.math.utah.edu/~levin>

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF UTAH, 155 S. 1400 E., SALT LAKE CITY, UT 84112-0090

Current address: Escuela de Matematica, Universidad de Costa Rica, San Pedro de Montes de Oca, Costa Rica

E-mail address: mendez@math.utah.edu, pmendez@emate.ucr.ac.cr

URL: <http://www.math.utah.edu/~mendez>