

## WORKSHEET SOLUTIONS: THE FALLING CALCULUS BOOK

Names and student IDs: Solutions  $[\pi\pi\pi-\pi\pi-\pi\pi\pi\pi]$

The gravitational constant at the surface of the planet Yuggxth is exactly  $2 \text{ m/sec}^2$ . It has no atmosphere, so falling objects encounter no friction. A calculus book is dropped down a vertical hole at time  $t = 0 \text{ sec}$ , so its height at time  $t \text{ sec}$  is  $p(t) = -t^2 \text{ m}$  above the surface. (Nobody on Yuggxth knows any calculus.) Note:  $-t^2$ , because the calculus book is falling *down*.

We want to know the velocity of the calculus book at time  $t = 2 \text{ sec}$ . (This is analogous to the reading of the speedometer in a car.)

As a first approximation, let's find the average velocity over the time interval  $[2, 4]$  (in seconds). It is

$$\frac{p(4) - p(2)}{4 - 2} = \frac{-4^2 - (-2^2)}{4 - 2} = \frac{-12}{2} = -6 \text{ m/sec.}$$

The answer is negative because the calculus book is moving down and we measured height in meters *above* the surface. Thus, the calculus book is rising at  $-6 \text{ m/sec}$ , or falling at  $6 \text{ m/sec}$ . Keep the signs right!

(What happens if we measure height in meters *below* the surface?)

1. Use the same method to find the average velocity over the time interval  $[2, 3]$  (in seconds).

*Solution.*

$$\frac{p(3) - p(2)}{3 - 2} = \frac{-3^2 - (-2^2)}{3 - 2} = \frac{-5}{1} = -5 \text{ m/sec.}$$

□

2. Use the same method to find the average velocity over the time interval  $[2, 2.1]$  (in seconds). (Feel free to use a calculator on this one.)

*Solution.*

$$\frac{p(2.1) - p(2)}{2.1 - 2} = \frac{-2.1^2 - (-2^2)}{2.1 - 2} = \frac{-0.41}{0.1} = -4.1 \text{ m/sec.}$$

□

3. Use the same method to find the average velocity over the time interval  $[2, 2.01]$  (in seconds). (Feel free to use a calculator on this one.)

*Solution.*

$$\frac{p(2.01) - p(2)}{2.01 - 2} = \frac{-2.01^2 - (-2^2)}{2.01 - 2} = \frac{-0.0401}{0.01} = -4.01 \text{ m/sec.}$$

□

4. Use the same method to find the average velocity over the time interval  $[1, 2]$  (in seconds).

*Solution.*

$$\frac{p(1) - p(2)}{1 - 2} = \frac{-1^2 - (-2^2)}{1 - 2} = \frac{3}{-1} = -3 \text{ m/sec.}$$

□

5. Use the same method to find the average velocity over the time interval  $[1.9, 2]$  (in seconds). (Feel free to use a calculator on this one.)

*Solution.*

$$\frac{p(1.9) - p(2)}{1.9 - 2} = \frac{-1.9^2 - (-2^2)}{1.9 - 2} = \frac{0.39}{-0.1} = -3.9 \text{ m/sec.}$$

□

6. What goes wrong if you try to find the velocity at  $t = 2$  sec by taking both times to be  $t = 2$  sec?

*Solution.* You get the meaningless expression “ $\frac{0}{0}$ ”. (Important note: the expression “ $\frac{0}{0}$ ” can **never** appear in an equation.) □

7. Nevertheless, what do you think the velocity at  $t = 2$  sec actually is?

*Solution.* Looking at the results above, it sure seems that the correct velocity at  $t = 2$  sec is  $-4$  m/sec. □